

What is *Inoceramus peruanus* Brüggen, 1910 ?

ANNIE V. DHONDT¹, IRENEUSZ WALASZCZYK², NADEJDA TCHEGLIAKOVA³ & ETIENNE JAILLARD⁴

¹Royal Belgian Institute of Natural Sciences, Vautierstr. 29, B-1000 Brussels, Belgium.

E-mail: annie.dhondt@naturalsciences.be

²Department of Geology, Warsaw University, Al. Zwirki i Wigury, 93, PL-02-089 Warsaw, Poland.

E-mail: i.walaszczyk@uw.edu.pl

³Departamento de Geociencias, Facultad de Ciencias, Universidad Nacional de Colombia, Santa Fé de Bogota, Colombia. E-mail: nadejda@ciencias.unal.edu.co

⁴IRD-LGCA, Maison des Géosciences, BP 53, 1381 rue de la Piscine, F-38041 Grenoble Cedex.

E-mail: Etienne.Jaillard@ujf-grenoble.fr

ABSTRACT

DHONDT, A.V., WALASZCZYK, I., TCHEGLIAKOVA, N. & JAILLARD, E. 2004. What is *Inoceramus peruanus* Brüggen, 1910 ? *Acta Geologica Polonica*, **54** (3), 535-539. Warszawa.

The figured holotype of the Lower Turonian *Inoceramus peruanus* BRÜGGEN, 1910 from Peru was found in the collections of the Goldfuss Museum in Bonn. Its general shape is close to that of *Mytiloides goppelnensis* (BADILLET & SORNAY) and *M. kossmati* (HEINZ), and is a *Mytiloides*. The ornamentation consists of the double-rugae typical of *M. kossmati*. The presence of *M. peruanus* in Peru, Ecuador and Colombia is discussed.

Key-words: Cretaceous, South America Lower Turonian, Bivalvia, *Mytiloides peruanus*.

INTRODUCTION

Today, in Colombian and Peruvian palaeontological and geological literature the taxon *Inoceramus peruanus* BRÜGGEN, 1910 is frequently mentioned from Turonian-Coniacian strata. However, the taxon is rarely figured and never described.

Inoceramus peruanus BRÜGGEN, 1910 (p. 751, pl. 25, fig. 6) was first described from the Cuesta de Huanyamba (between Celendin and Sendamal, Peru) and the age BRÜGGEN suggested for it was "Lower Senonian".

STEINMANN (1929, p. 147) suggested a Turonian age for this Peruvian taxon and for the strata in which it was found, based on the presence of *Coilophoceras* (sic) *requieni* and *Inoceramus labiatus*. Unfortunately, in later literature the locality "Cuesta de Huanyamba" is not found. It is not indicated on the 1/100.000 map. Thus, today we have no way to locate it precisely.

BENAVIDES-CACERES (1956) mentioned *Inoceramus peruanus* from a few Peruvian localities, but they vary in age from the Early Turonian to the Late Coniacian (ibidem, p. 387, p. 397 – *Coilopoceras newelli* Zone and *Buchiceras bilobatum* Zone). None of the inoceramid specimens in the lists of BENAVIDES-CACERES (1956) were figured – therefore it is impossible to confirm what his concept of *Inoceramus peruanus* represents.

On the other hand, BENAVIDES-CACERES (1956, p. 473) also stated a strong similarity between *Coelopoceras* (sic) *requieni* in BRÜGGEN (1910) and his taxon *Coilopoceras jenksi* from the Lower Turonian Coñor Fm. *Inoceramus peruanus* BRÜGGEN, 1910 co-occurs with *Coelopoceras requieni* (D'ORBIGNY) in BRÜGGEN (1910) and STEINMANN (1929), making an early Turonian age probable.

In Colombia, BÜRGL & DUMIT TOBON (1954) and BÜRGL (1957) mentioned *Inoceramus peruanus* BRÜGGEN

from the “Coniaciano inferior and superior” in Girardot, Prov. Alto Magdalena, Alban, Guacheta and Ubaté (Prov. Bogota) Colombia. In these areas some strata containing “*Inoceramus peruanus*” are overlain with strata with *Didymotis* and are assigned to a middle Coniacian age by BÜRGL (1957). It seems likely that the age assignment used in these Colombian publications is not completely accurate.

Possibly because of general poor understanding of Turonian inoceramids from Colombia, F. ETAYO-SERNA asked SORNAY (1981) to study inoceramids from Quebrada La Modelia (Cundinamarca, Colombia) of the La Frontera Fm., Lower Turonian. Amongst the specimens SORNAY studied he described a new taxon *Mytiloides modeliaensis* (SORNAY, 1981).

VILLAMIL & ARANGO (1998) on the stratigraphy of the latest Cenomanian and early Turonian facies of Colombia, and VILLAMIL (1998) on a sequence stratigraphic model for basinal Cretaceous facies of Colombia, also discussed and illustrated inoceramid taxa. Among others they mentioned and illustrated *Mytiloides kossmati*, *M. columbianus*, *M. modeliaensis*, *M. subhercynicus* all from the Lower-Middle Turonian, but suggested that *M. modeliaensis* is younger than the three others. They did not mention the taxon *M. peruanus*. VILLAMIL & ARANGO did not describe any of the taxa they illustrated, and unfortunately their illustrations are not ideal. This makes interpretation of these taxa and comparison between them difficult.

HEINZ (1935) introduced the new name *Orpheoceramus columbianus* to replace his former identification “*Inoceramus plicatus*” D’ORBIGNY, 1842 (in HEINZ 1928). The “*Inoceramus plicatus*” specimen from the 1928 publication came from Cachira in Colombia and HEINZ considered it Turonian. In 1935, he suggested a tentatively Early Cretaceous age for this specimen, but that is certainly erroneous.

In JAILLARD & *al.* (1999), *Mytiloides* specimens from SW Ecuador (Carmelo Fm. in the Celica Basin, fig. 9) are mentioned. By re-studying these specimens we have come to the conclusion that some of the better preserved ones undoubtedly belong to *M. peruanus*.

Based on material from outside South America several authors discussed these taxa and suggested comparisons with taxa from other regions. HEINZ in BESAIRIE (1930) first listed the taxon *Inoceramus kossmati* from the Turonian of Madagascar, which he regarded as a variety of *Inoceramus naumanni* YOKOHAMA. He described it as a separate species in 1933, which he referred to the genus *Striatoceramus* (*S. kossmati*).

WALASZCZYK (1992) discussed the species *Mytiloides kossmati* (HEINZ, 1930) and indicated (p. 10) that *M. opalensis* (sensu SEITZ non BOESE, 1934), *M. goppelnensis* (BADILLET & SORNAY, 1980) and *M. modeliaensis*

(SORNAY, 1981) are synonymous with *M. kossmati* (HEINZ). He also considered *M. columbianus* (HEINZ, 1935) to be very close to *M. kossmati*. In the same paper, he suggested that *M. subhercynicus* (SEITZ, 1934) is not an independent taxon but includes extreme forms of several other taxa such as *M. kossmati* (HEINZ), *M. hercynicus* (PETRASCHECK, 1903) and *M. opalensis* (BOESE, 1923).

Based on the rich and collected bed-by-bed material from the Pueblo section, US Western Interior, WALASZCZYK & COBBAN in KENNEDY & *al.* (2000, pp. 322-323) re-discussed some of these Lower Turonian taxa. This resulted in a slightly different interpretation from that in WALASZCZYK (1992). Thus in 2000, *M. kossmati* includes *M. columbianus* (HEINZ, 1935) and *M. modeliaensis* (SORNAY, 1981), but *M. goppelnensis* (BADILLET & SORNAY, 1980) and *M. subhercynicus* (SEITZ, 1934) are considered as distinct taxa.

TAXONOMY

Mollusca
Class Bivalvia

Inoceramidae ZITTEL, 1881 (ICZN 473)

Mytiloides BRONGNIART in CUVIER, 1822

TYPE SPECIES: *Ostracites labiatus* VON SCHLOTHEIM, 1813

Mytiloides peruanus (BRÜGGEN, 1910)
(Text-figs ?1.3, 1.4-1.6, ?1.7)

- v.* 1910. *Inoceramus peruanus* n.sp., BRÜGGEN, p. 751, pl. 25, fig. 6
1928. *Inoceramus plicatus* D’ORB.; HEINZ, p. 63, pl. 4, fig. 4.
1929. *Inoceramus peruanus* BRÜGG.; STEINMANN, p. 147
non 1954. *Inoceramus peruanus* BRUEGGEN; BÜRGL & DUMIT TOBON, p. 32.
? 1957. *Inoceramus peruanus* BRÜGGEN; BÜRGL, p. 138, 139, 140, pl. 15, fig.

HOLOTYPE: Bonn University (Rheinische Friedrich-Wilhelms Universität, Paläontologisches Institut, Goldfuss Museum), Bonn, Germany: n° Brüggén 4.

LOCUS TYPICUS: Cuesta de Huanyamba (between Celendin and Sendamal), Peru.

STRATUM TYPICUM: In BRÜGGEN (1910) the age given is “Untersenen”; STEINMANN (1929) re-interpreted this age as “Turonian”.

ADDITIONAL MATERIAL: Six relatively poorly pre-




Fig. 1. IRSNB MI: Institut royal des Sciences naturelles de Belgique, Mesozoic Invertebrates ; 1 – *Mytiloides* cf. *ratonensis* (WALASZCZYK & COBBAN, 2000), IRSNB MI 11000 from the Girardot-Nariño section, 4 km NW of Girardot (Cundimarca, Colombia, Lat. 4°21' 31", Long. 74° 12'), Hondita Fm., Upper Turonian, × 1; 2 – *Mytiloides scupini* (HEINZ, 1930b), IRSNB MI 10999, from the Girardot-Nariño section, 4 km NW of Girardot (Cundimarca, Colombia, Lat. 4°21' 31", Long. 74° 12'), Hondita Fm., Upper Turonian; × 1; 3 – *Mytiloides* ?*peruanus* (BRÜGGEN, 1910), IRSNB MI 11001, from Yamana, NW of Catamayo (Ecuador) Carmelo Fm., Celica Basin: Jaillard Coll. 96-2, × 1; 4 – *Mytiloides peruanus* (BRÜGGEN, 1910) (Bonn Goldfuss Museum, Brügggen A1) from the upper Lower Turonian of Cuesta de Huanyamba (between Celendin and Sendamal), Peru, × 1.5; 5 – *Mytiloides peruanus* (BRÜGGEN, 1910) (Bonn Goldfuss Museum, Brügggen A3) from the upper Lower Turonian of Cuesta de Huanyamba (between Celendin and Sendamal), Peru, × 1.5; 6 – *Mytiloides peruanus* (BRÜGGEN, 1910) holotype (Bonn Goldfuss, Brügggen 4), from the upper Lower Turonian of Cuesta de Huanyamba (between Celendin and Sendamal), Peru, × 1.5; 7 – *Mytiloides* ?*peruanus* (BRÜGGEN, 1910) (Bonn Goldfuss Museum, Brügggen A4) from the upper Lower Turonian of Cuesta de Huanyamba (between Celendin and Sendamal), Peru, × 1.5

served specimens from the same locality and collection as the holotype – right valves Brü A1, Brü A4; left valves Brü A2, Brü A3, Brü A5, Brü A6. Four specimens from Yamana and El Carmelo, NW of Catamayo (Ecuador) Carmelo Fm., the Celica Basin: Jaillard Coll. 96-2.

DESCRIPTION: Shell small to medium-sized, prosocline, subcircular to subovate, moderately unequivalved with slight convexity, umbo small and not prominent.

Ornamentation consists of double-ridged rugae, very closely situated near the umbo and progressively more widely spaced towards the ventral margin, but never very far apart. Auricles small and poorly developed.

Dimensions (see Text-fig. 2 for explanations):

spec.	Illustr.	h	l	H	L	s	α	δ	valve	h_{\max}
Brü 4		33	28	-	-	(18)	-	55	LV	35
BrüA1		26.5	23.5	-	-	(15.5)	85	-	RV	36
BrüA2	not	26.7	24.5	-	-	19.5	93	40	LV	28.5
BrüA3		28.5	23.5	-	-	17.5	78	50	LV	33.5

DISCUSSION: *Mytiloides peruanus* in the general shape and the double-ridged rugae is morphologically very close, and possibly conspecific with *Mytiloides kossmati* (HEINZ, 1930) [which includes *M. modeliaensis* (SORNAY, 1980) and *M. columbianus* (HEINZ, 1935) – fide WALASZCZYK & COBBAN in KENNEDY & al. 2000]. However, as the holotype and other topotypic material of the species is poorly preserved, we don't want to make any




Fig. 2. Basic measurements on inoceramid specimens

formal taxonomic decision. Such conclusion should be preceded by studies in the original locality.

A number of forms, which in general outline and ornamentation closely resemble BRÜGGEN's species were reported from Columbia by VILLAMIL (1998) and VILLAMIL & ARANGO (1988). This concerns particularly their *Mytiloides subhercynicus* (see VILLAMIL & ARANGO 1988, figs 4, 6E, 7B, 8B and 24 and VILLAMIL 1998, figs 8, 20E), *M. columbianus* (VILLAMIL & ARANGO 1988, figs 6F, G), and partly *Mytiloides kossmati* (VILLAMIL & ARANGO 1988, figs 2-4, 6B, 24 and VILLAMIL 1998, figs 11, 17J, 19).

OCCURRENCE: Lower Turonian of South America (Peru, Ecuador, Colombia).

ADDENDUM

From the Girardot-Nariño section, 4 km NW of Girardot (Cundimarca, Colombia, Lat. 4°21'31", Long. 74°12'), in the middle valley of the Magdalena one of us (N.T.) in the Hondita Fm. (defined by J. DE PORTA 1966) collected specimens of mytiloid inoceramids – together with the following Foraminifera: *Whitenella gigantea*, *W. archaeoretacea*, *W. inornata*, *Concavatotruncana primitiva* and *C. concavata*.

These inoceramids, on preliminary identification, are *Mytiloides scupini* (HEINZ, 1930b) (see TRÖGER 1999) and *M. cf. ratonensis* (WALASZCZYK & COBBAN 2000) taxa that co-occur in the Upper Turonian of the Western Interior. They are illustrated on Text-fig. 1.1 (*M. scupini*) and 1.2 (*M. cf. ratonensis*).

Acknowledgments

We are very grateful to Martin SANDER (Director Goldfuss Museum, Paläontologisches Institut, Rheinische-Friedrich-Wilhelms-Universität, Bonn (Germany) who gave us access to the material studied by BRÜGGEN (1910). A grant from the ABC programme is gratefully acknowledged by I. WALASZCZYK. It made a visit to the RINScB in Brussels possible. We thank Wilfried MISEUR (Brussels) for making the photographs. Peter J. HARRIES and an anonymous reviewer are acknowledged for careful and constructive reviewing of the manuscript.

REFERENCES

- BADILLET, G. & SORNAY, J. 1980. Sur quelques formes du groupe d'*Inoceramus labiatus* décrites par O Seitz. Impossibilité d'utiliser ce groupe pour une datation stratigraphique du Turonien inférieur du Saumurois (France). *Comptes Rendus de l'Académie des Sciences de Paris, Série D* **290**, 323-325.

- BENAVIDES-CACERES, V.E. 1956. Cretaceous system in Northern Peru. *Bulletin of the American Museum of Natural History*, **108**, 355-493.
- BOESE, E. 1923. Algunas faunas cretácicas de Zacatecas, Durango y Guerrero. *Boletín Instituto Geológico de México*, **42**, 219 pp. + IV pp.
- BRONGNIART, A. in CUVIER, G. 1822. Description géologiques des couches des environs de Paris. In: "Les Ossements Fossiles". Vol. II, 2: 229-648, Paris.
- BRÜGGEN, H. 1910. Die Fauna des unteren Senons von Nord-Peru. *Beilage Band Neues Jahrbuch für Mineralogie, Geologie und Paläontologie*, **30**, 717-788.
- BÜRGL, H. 1957. Biostratigrafía de la Sabana de Bogota y sus alrededores. *Boletín Geológico*, **5**, 2, 113-185.
- BÜRGL, H. & DUMIT TOBON, Y. 1954. El Cretáceo superior en la región de Girardot. *Boletín Geológico*, **2**, 1, 23-48.
- HEINZ, R. 1928. Über die Oberkreide-Inoceramen Süd-Amerikas und ihre Beziehungen zu denen Europas und anderer Gebiete. Beiträge zur Kenntnis der oberkretazischen Inoceramen V. *Mitteilungen aus dem Mineralogisch-Geologischen Staatsinstitut, Hamburg*, **10**, 42-97.
- 1930a. In: H. BESAIKIE, Recherches géologiques à Madagascar. Contribution à l'étude des ressources minérales. Thèse présentée à la Faculté des Sciences de l'Université de Paris, 272 pp.
- 1930b. Zur stratigraphischen Stellung der Sonnenberg-schichten bei Waltersdorf in Sachsen (westsüdwestlich von Zittau). Beiträge zur Kenntnis der Inoceramen IX. *Jahresberichte des Niedersächsischen Geologischen Vereins zu Hannover*, **23**, 25-29.
- 1933. Inoceramen von Madagaskar und ihre Bedeutung für die Kreide-Stratigraphie. Beiträge zur Kenntnis der Inoceramen XII. *Zeitschrift der Deutschen Geologischen Gesellschaft*, **85**, 242-259.
- 1935. Unterkreide-Inoceramen von der Kapverden-Insel Maio. Beiträge zur Kenntnis der Inoceramen XVIII. *Neues Jahrbuch für Mineralogie und Geologie, Beilage-Band*, **73**, B, 302-311.
- JAILLARD, E., LAUBACHER, G., BENGTSON, P., DHONDT, A.V., & BULOT, L.G. 1999. Stratigraphy and evolution of the Cretaceous forearc Celica-Lancones basin of southwestern Ecuador. *Journal of South American Earth Sciences*, **12**, 1, 51-68.
- KENNEDY, W.J., WALASZCZYK, I. & COBBAN, W.A. 2000. Pueblo, Colorado, USA, candidate Global Boundary Stratotype Section and Point for the base of the Turonian Stage of the Cretaceous, and for the base of the Middle Turonian Substage, with a revision of the Inoceramidae. *Acta Geologica Polonica*, **50**, 295-334.
- ORBIGNY, A. d'1842. Voyage dans l'Amérique méridionale (le Brésil, la république orientale de l'Uruguay, la république argentine, la Patagonie, la république du Chili, la république de Bolivie, la république du Pérou) exécuté pendant les années 1826-1833. Tome troisième, 4^e partie: Paléontologie, P. Bertrand, Paris; Veuve Levrault, Strasbourg, 188 pp.
- PETRASCHECK, W. 1903. Ueber Inoceramen aus der Kreide Böhmens und Sachsens. *Jahrbuch der Geologischen Reichsanstalt*, **53**, 153-168.
- PORTA, J. DE 1966. La geología del Extremo Sur del Valle Medio del Magdalena entre Honda y Guataqui (Colombia). *Boletín de Geología de la Universidad Industrial de Santander*, **22-23**, 5-347.
- SCHLOTHEIM, E.T von 1813. Beiträge zur Naturgeschichte der Versteinerungen in geognostischer Hinsicht. *Leonhard's Taschenbuch der Mineralogie* **7**, p. 93, Berlin.
- SEITZ, O. 1934. Die Variabilität von *Inoceramus labiatus* v. Schloth. *Jahrbuch der Preussischen Geologischen Landesanstalt*, **55**, 429-474.
- SORNAY, J. 1981. Inocérames (Bivalvia) du Turonien inférieur de Colombie (Amérique du Sud). *Annales de Paléontologie (Invertébrés)*, **67**, 135-148.
- STEINMANN, G. 1929. Geologie von Perú. 448 pp. *Carl Winters Universitätsbuchhandlung*; Heidelberg.
- TRÖGER, K.-A. 1999. *Inoceramus scupini* HEINZ im ober-Turon der Subherzynen Kreidemulde. *Greifswalder Geowissenschaftliche Beiträge*, **6**, 209-215.
- VILLAMIL, T. 1998. Chronology, relative sea-level history and a new sequence stratigraphic model for basinal Cretaceous facies of Colombia. In: J. PINDELL & C. DRAKE (Eds), Eustacy and Tectonostratigraphic Evolution of northern South America. *SEPM Special Publication* **58**, 161-216.
- VILLAMIL, T. & ARANGO, C. 1998. Integrated stratigraphy of latest Cenomanian-Early Turonian facies of Colombia. In: J. PINDELL & C. DRAKE (Eds), Eustacy and Tectonostratigraphic Evolution of northern South America. *SEPM Special Publication* **58**, 129-159.
- WALASZCZYK, I. 1992. Turonian through Santonian deposits of the Central Polish Uplands; their facies development, inoceramid paleontology and Stratigraphy. *Acta Geologica Polonica*, **42** (1-2), 1-122.
- WALASZCZYK, I. & COBBAN, W. A. 2000. Inoceramid faunas and biostratigraphy of the Upper Turonian-Lower Coniacian of the Western Interior of the United States. *Special Papers in Palaeontology*, **64**, 118 pp.

Manuscript submitted: 10th February 2004,

Revised version: 20th August 2004.