

*Wacław Bałuk & Andrzej Radwański*

A new occurrence of fossil acrothoracican cirripedes: *Trypetesa polonica* sp. n. in hermitted gastropod shells from the Korytnica Basin (Middle Miocene; Holy Cross Mountains, Central Poland), and its bearing on behavioral evolution of the genus *Trypetesa* . . . . . 1

*Jarosław Stolarski*

Miocene Scleractinia from the Holy Cross Mountains, Poland; Part 1 — Caryophylliidae, Flabellidae, Dendrophyllidae, and Micrabaciidae . . . . . 37

*Ewa Roniewicz & Jarosław Stolarski*

Miocene Scleractinia from the Holy Cross Mountains, Poland; Part 2 — Archaeocoeniina, Astracina, and Fungiina . . . . . 69

*Leszek Lindner*

Stratigraphy of main Pleistocene loess horizons and paleosols in mid-eastern Europe . . . . . 85

*Wiesław Barczyk*

Succession of the Tithonian to Berriasian brachiopod faunas at Rogoźnik, Pieniny Klippen Belt . . . . . 101

*Stanisław Speczik & El-Sayed A. A. Youssef*

Fluid inclusion studies of Bahariya barites (Western Desert, Egypt) . . . . . 109

*Tadeusz Merta*

A new, universal method of thin-section - to - sieve transformation of granulometric data . . . . . 117

<i>Andrzej Wierzbowski</i> Biostratigraphical correlations around the Oxfordian/Kimmeridgian boundary . . .	149
<i>Antoni Hoffman, Michał Gruszczyński, Krzysztof Malkowski, Stanisław Halas, Bronisław A. Matyja &amp; Andrzej Wierzbowski</i> Carbon and oxygen isotope curves for the Oxfordian of Central Poland . . . . .	157
<i>Maciej Bąbel</i> Dissolution of halite within the Middle Miocene (Badenian) laminated gypsum of southern Poland . . . . .	165
<i>Teresa Czyżewska &amp; Andrzej Radwański</i> Middle Miocene (Badenian) delphinid and phocoenid remains from the Fore-Carpat- hian Depression in southern Poland . . . . .	183
<i>Andrzej Pisera &amp; Adam Bodzioch</i> Middle Triassic lyssacinosan sponges from Upper Silesia (southern Poland), and the history of hexactinosan and lychniscosan sponges . . . . .	193
<i>Stanisław Kwiatkowski</i> Origin of the chert laminae and silico-calcareous nodules in uppermost Rõth cavernous limestone at Gogolin (Lower Silesia) . . . . .	209
<i>Pieter Berendsen &amp; Stanisław Speczik</i> Sedimentary environment of Middle Ordovician iron oolites in northeastern Kansas, U.S.A. . . . .	215

---