

# The oldest Cambrian trilobites from the Holy Cross Mountains, Poland: taxonomic, stratigraphic and biogeographic reappraisal

ANNA ŻYLIŃSKA

*Faculty of Geology, University of Warsaw, Żwirki i Wigury 93, PL-02-089 Warsaw, Poland.  
E-mail: anna.zylinska@uw.edu.pl*

## ABSTRACT:

Żylińska, A. 2013. The oldest Cambrian trilobites from the Holy Cross Mountains, Poland: taxonomic, stratigraphic and biogeographic reappraisal. *Acta Geologica Polonica*, **63** (1), 57–87. Warszawa.

Authorship issues are clarified, new photographic documentation is provided and emended systematic descriptions are presented for the oldest Cambrian trilobite taxa from the Holy Cross Mountains (Poland). Biostratigraphic analysis of the fauna allows correlation with the traditional *Holmia kjerulfi*-group Zone of Scandinavia, the *Callavia* Zone of Britain and Newfoundland, the lower and middle part of the *Sectigena* Zone of Morocco and the Marianian Stage of Spain. The trilobites display a strong biogeographic signal linked with West Gondwana and Avalonia and a suggestion is made that the TESZ margin of Baltica with the Małopolska Massif was liable to currents from those areas that distributed planktonic trilobite larvae.

**Keywords:** Ellipsocephalidae; Holmiidae; Cambrian Series 2; Stratigraphy; Biogeography; Holy Cross Mountains; Trilobita.

## INTRODUCTION

The oldest Cambrian trilobite faunas of the Holy Cross Mountains, Poland, have been known since 1917, when Jan Samsonowicz discovered fine-grained, fossiliferous sandstones along the Koprzywianka River in the vicinity of Gieraszowice in the far south-east of the area (Samsonowicz 1918, 1920); although some trilobites seem to have been found as early as in 1915 (see discussion under *Strenuella polonica* Czarnocki, 1926 in the systematic part). Subsequent papers provided lists of trilobites, even new species names, and tied associated fauna to particular lithologies (Czarnocki 1919, 1926, 1927a, b, 1932, 1933). Unfortunately, the taxa were never described or illustrated by their authors.


Many of the trilobite names given in these reports must be treated as *nomina nuda* because of the loss of a large part of the original collections housed in the Museum of the Polish Geological Institute in Warsaw during the Warsaw Uprising in 1944. The first illustrations and descriptions, based partly on the original specimens, were presented by Samsonowicz (1959a, b, c, and in the posthumous paper, Samsonowicz 1962, prepared from his manuscripts). In these papers, a number of species mentioned by Czarnocki (*op. cit.*) were retained as valid because the original specimens were found with appropriate museum labels. This original authorship usually remained, however, unrecognized in later reports (Bergström 1973b; Orłowski 1974, 1985a; Geyer 1990b).

This paper clarifies the authorship issues for the oldest known Cambrian trilobite taxa from the Holy Cross Mountains, provides new photographic documentation and supplies emended systematic descriptions for some of them. Moreover, the stratigraphic significance of this fauna is discussed in detail as well as in terms of diversity and biogeographic significance, thus supplementing new stratigraphic and biogeographic conclusions on Cambrian trilobites in the southern part of the Holy Cross Mountains (Kielce Region) (Żylińska and Masiak 2007; Żylińska and Szczepanik 2009). Two ellipsocephalid species from this fauna have been recently revised and assigned to *Berabichia oratrix* (Orłowski, 1985a), based on bi- and

multivariate analysis of morphometric parameters and graphic techniques (Żylińska *et al.* in press).

## GEOLOGICAL SETTING

The Holy Cross Mountains (HCM) represent one of the very few areas in Europe that expose Palaeozoic rocks in the direct vicinity of the Teisseyre–Tornquist Line (Text-fig. 1A), with its basement forming a part of the Trans-European Suture Zone (TESZ; Berthelsen 1992), an important geotectonic domain separating the Precambrian East European Craton (EEC) from the Palaeozoic fold-belts of central and western Europe. This WNW–



Text-fig. 1. (A) Sketch-map of Poland with location of the Holy Cross Mountains (HCM) in relation to the East European Craton (EEC) and Teisseyre–Tornquist Line (TTL). Black dot with "G" marks the Goczałkowice IG-1 borehole. (B) Geological sketch-map of the HCM showing the distribution of Cambrian deposits, modified from Orłowski (1975b, 1992). (C) Geological sketch-map of the Klimontów Anticlinorium with the Cambrian lithostratigraphic formations and location of exposures yielding the oldest known trilobites, compiled and modified from Samsonowicz (1962) and Orłowski and Mizerski (1995). LD – Leśniakowa Dębina, JH – Jazwina Hill

ESE-oriented, 70 km long and 50 km wide hilly area, is located in south-central Poland (Text-fig. 1B). The morphology follows the tectonic structure of the area, with the orientation of the particular hilly ranges corresponding closely to the orientation of the main anticlines and synclines in the Palaeozoic core of the Mountains. The area was subjected to multi-stage evolution, in which the Late Cretaceous–Early Palaeogene tectonic inversion and uplift that resulted in partial removal of Permian and Mesozoic strata and exposure of the Palaeozoic, was one of the last structural events (Kutek and Głazek 1972; Krzywicz *et al.* 2009). The latest marine influence in the HCM was in the Miocene, when the sea of the Carpathian Foreland basin encroached on the southern margins of the area (e.g., Radwański 1969).

Different facies development, stratigraphy and tectonic evolution have resulted in the subdivision of the Palaeozoic of the HCM into two regions: the southern, Kielce Region, located on the northern part of the Małopolska Block, and the northern, Łysogóry Region, located on the Łysogóry Block, separated by the WNW–ESE-oriented Holy Cross Fault (e.g., Czarnocki 1919). These regions were considered to be tectono-stratigraphic units (e.g., Czarnocki 1919). Recent studies point to their palaeogeographic character (e.g., Belka *et al.* 2000, 2002; Nawrocki and Poprawa 2006 and publications referenced therein), with contrasting views on whether they are of proximal or exotic provenance with regard to the palaeocontinent of Baltica. These ambiguous views on the nature of both blocks result from inconsistent information on their composition, provenance of detrital material and the biogeographic signature of faunas in their sedimentary cover. Recent geophysical data indicate that both blocks have a similar crustal composition, comparable to that of the EEC (Malinowski *et al.* 2005); according to Nawrocki *et al.* (2007), they represent proximal terranes relocated along the TESZ margin of the Baltica palaeocontinent.

The Cambrian strata in the HCM are composed of siliciclastic facies with an estimated thickness of 2500–3500 m (e.g. Orłowski 1988). The trilobites studied come from the Kielce Region, where the source strata are exposed within the Klimontów Anticlinorium, a Variscan structural unit that covers almost half of the area (Text-fig. 1B, C). A few specimens come from two exposures within the Dyminy Anticline, in the western part of the Kielce Region (Text-fig. 1B). The fossils were encountered in two formations, i.e. the Ociesęki Sandstone, cropping out in the south-easternmost, southern and western part of the area, and the Kamieniec Shale, occurring in the central and eastern part of the unit (Text-fig. 1B, C). The Ociesęki Formation includes fine-grained sandstones and siltstones, often


strongly bioturbated, with thin claystone intercalations (Orłowski 1975b; Kowalczewski *et al.* 2006), representing shallow-marine settings as indicated by sedimentary structures and ichnofossils (Studencki 1988; Orłowski 1989, 1992; Mizerski *et al.* 1999; Orłowski and Żylińska 2002). The Kamieniec Formation is clay- and siltstone-dominated with thin intercalations of fine-grained, non-bioturbated sandstones (Orłowski 1975b; Orłowski and Mizerski 1995); its sedimentary structures and the few ichnofossils point to an outer shelf setting below the storm wave base (Studencki 1988; Mizerski *et al.* 1991).

#### COMPOSITION OF THE FAUNAS

The fauna under discussion is composed of six taxa representing the Holmiidae [*Holmia marginata* Orłowski, 1974; *H. glabra* Orłowski, 1974; *Schmidtellus panowi* (Samsonowicz, 1959a); *S. nodosus* Orłowski, 1985a; *Kjerulfia orcina* Orłowski, 1974; *Postfallotaspis spinatus* Orłowski, 1985a], four taxa representing the Ellipsocephalidae [*Strenuella polonica* Czarnocki, 1926; *S. zbelutkae* Orłowski, 1985a; *Berabichia oratrix* (Orłowski, 1985a); *Termierella sandomirensis* Samsonowicz, 1962], one taxon of the Chengkouiidae (*Acanthomicmacca klimontowi* Orłowski, 1985a), and one taxon of the Atopidae (*Atops granulatus* Orłowski, 1985c) (see Żylińska *et al.* in press, and systematic part below) (Text-fig. 2). The abundance of particular taxa in the material, reaching over 1600 specimens, varies considerably. The fauna is dominated by the Ellipsocephalidae, composing 78% of the material (Text-fig. 3A). Holmiids, including the zone-indicative genera, constitute only 16% of the material. Representatives of the chengkouiids and atopids comprise the remaining 6%. Of the four species of the Ellipsocephalidae, the dominant one is *Berabichia oratrix* (see Żylińska *et al.* in press, for an emended diagnosis of this species), with the Berger-Parker index of dominance for the whole material being 0.48 (Hammer and Harper 2006). The composition of faunas representing particular formations varies. The composition of the fauna from the Ociesęki Formation (1 in Text-fig. 4) more or less corresponds to that of all the material (Text-fig. 3B). However, the composition of the fauna from the Kamieniec Formation (2 in Text-fig. 4) differs considerably (Text-fig. 3C); represented by 145 specimens, it comprises only six species (Berger-Parker index of 0.55): *Strenuella zbelutkae* (55%), *Acanthomicmacca klimontowi* (32%), *Berabichia oratrix* (3%) and three species of Holmiidae (10%). The presence in the Kamieniec Formation of a different species of *Strenuella*, i.e.

family	taxa	localities																									
		Słowik	Telegraf Hill	Szczecno	Stołowa Hill	Cisów	Igrzyczna Hill	Nowa Huta	Majeje	Jazwina Hill	Leśniakowa Dębina	Sterczyna Hill	Ociesęki	Ryj Hill	Wola Wąkopna	Rembów	Zbelutka	Ujazdek	Malikowska Hill	Buczyna	Pranie	Olbierzowice	Kucebrza	Bukówki-Rybnica	Julianów	Dąb	Gieraszowice
Ellipsocephalidae	<i>Berabichia oratrix</i>	+	+			+	+	+		+	+	+				+	+	+	+	+	+						
	<i>Strenuella polonica</i>						+		+	+	+					+	+				+						
	<i>Strenuella zbelutkae</i>										+	+					+								+		
	<i>Termierella sandomirensis</i>																										+
Holmiidae	<i>Holmia glabra</i>												+														
	<i>Holmia marginata</i>			+	+		+			+	+	+			+				+							+	+
	<i>Holmia</i> sp.					+						+							+	+	+		+		+	+	
	<i>Kjerulfia orcina</i>								+		+	+	+		+				+	+					+		
	<i>Postfallotaspis spinatus</i>						+																				
	<i>Schmidtellus nodosus</i>						+				+	+							+	+							
	<i>Schmidtellus panowi</i>										+						+									+	+
other	<i>Acanthomicmacca klimontowi</i>						+					+		+		+					+	+		+	+		
	<i>Atops granulatus</i>										+	+															

Text-fig. 2. Distribution of trilobite taxa (grouped into families) in the particular localities arranged from west to east. Localities in the Kamieniec formation are marked in grey

Text-fig. 3. Pie-chart showing the abundance of particular taxa in the *Holmia-Schmidtellus* Assemblage Zone. A – total; B – Ociesęki formation, C – Kamieniec formation. Grey shades mark representatives of the Ellipsocephalidae

*S. zbelutkae* instead of *S. polonica*, which is very abundant in the Ociesęki Formation, as well as the extremely low representation of *Berabichia oratrix*, known from almost every exposure of the Ociesęki Formation in the studied interval, may suggest a common, e.g. environmental, factor influencing the distribution of particular trilobite species in both formations.

#### BIOSTRATIGRAPHIC ANALYSIS

Both formations with the oldest trilobite faunas are relatively well-constrained in the local zonal scheme for the HCM (Text-fig. 4). The Ociesęki Formation ranges from the lowermost *Holmia-Schmidtellus* Assemblage Zone through the *Protolenus-Issafeniella* Zone of Cambrian Series 2 up to the *Eccaparadoxides insularis* Zone of Cambrian Series 3; and the Kamieniec Formation ranges from the *Holmia-Schmidtellus* Assemblage Zone to the *Protolenus-Issafeniella* Zone of Cambrian Series 2 (Orłowski 1974, 1985a, b; Mizerski *et al.* 1986; Żylińska and Szczepanik 2009).





The *Holmia-Schmidtellus* Assemblage Zone was established by Orłowski (1987); previously this interval was referred to the *Holmia* Zone (e.g. Samsonowicz 1962; Orłowski 1975b, 1985a). Its *Holmia* representatives, i.e. *H. marginata* and *H. glabra*, show great similarity to *H. kjerulfi* (Linnarsson, 1871), the index taxon of the traditional *Holmia kjerulfi*-group Zone of



Scandinavia (Bergström and Ahlberg 1981; Ahlberg *et al.* 1986) or the upper part of the *Holmia kjerulfi* Assemblage Zone of Moczyłowska (1991), and it is commonly accepted that the *Holmia*-bearing beds of the HCM correspond to that part of Cambrian Series 2 in Scandinavia (Bergström 1973b; Bergström and Ahlberg 1981; Ahlberg *et al.* 1986; Ebbestad *et al.* 2003). Other holmiids that co-occur with *Holmia* include *Schmidtiellus panowi* and *S. nodosus*. In Scandinavia, *Schmidtiellus* Moberg (in Moberg and Segerberg 1906) appears in the oldest trilobite zone, i.e. in the traditional *Schmidtiellus mickwitzi* Zone (Bergström 1981; Ahlberg *et al.* 1986). However, based on sequence stratigraphy, the strata yielding this genus were shown to correlate with the traditional *Holmia kjerulfi*-group Zone (Nielsen and Schovsbo 2011). Species of *Kjerulfia* Kiær, 1917 occur in Scandinavia in the traditional *Holmia kjerulfi*-group Zone (Kiær 1917; Ahlberg *et al.* 1986; Ebbestad *et al.* 2003). In a recently proposed revised biostratigraphic scheme for the lower Cambrian of Scandinavia, based on sequence stratigraphy, the traditional *Holmia kjerulfi*-group Zone is now interpreted as the lower part of the *H. kjerulfi*-‘*O.*’ *linnarssoni* Zone (see discussion in Nielsen and Schovsbo 2011). A strange and still poorly known holmiid of the *Holmia*-*Schmidtiellus* Assemblage Zone is *Postfallotaspis spinatus*. It seems quite similar (see systematic part below) to the poorly known holmiid *Iyouella* Geyer and Palmer, 1995, from the *Sectigena* Zone of Morocco (Geyer and Palmer 1995), which corresponds to the upper part of the

*Callavia* Zone in Avalonia and the traditional *Holmia kjerulfi*-group Zone of Scandinavia. Acritarch assemblages recognized recently in the Dąb exposure (Szczepanik and Żylińska 2012) contain such taxa as e.g. *Skiagia ciliosa*, *Solisphaeridium implicatum* and *Cymatiosphaera postii*, suggesting a position in the *Heliosphaeridium dissimulare* – *Skiagia ciliosa* Assemblage Zone (Moczyłowska 1991), which have also been recovered from the traditional *Holmia kjerulfi*-group Zone of Sweden (Vidal 1981; Vidal and Nystuen 1990; Moczyłowska 1991).

Trilobites accompanying the holmiids belong to the Ellipsocephalidae, a family with wide palaeogeographic distribution (Hupé 1953; Geyer 1990b). They are most common in West Gondwana, particularly in Morocco, where they provide a number of index taxa for the Banian and ‘Tissafinian’ zones (Geyer 1990a; Geyer and Landing 2004), and in Spain, where they are index fossils for the Marianian and Bilbilian (e.g. Gozalo *et al.* 2003). In Avalonia, ellipsocephalid species are used for biostratigraphic subdivisions of the upper Comley Series in Great Britain (e.g. Rushton *et al.* 2011) and in Newfoundland, Canada (e.g. Fletcher 2006). Individual species are used for defining the uppermost Cambrian Series 2 zones in Scandinavia (Baltica) (Kiær 1917; Bergström and Ahlberg 1981; Nielsen and Schovsbo 2011) and in the Holy Cross Mountains (Orłowski 1988; Żylińska and Masiak 2007; Żylińska and Szczepanik 2009).

Ellipsocephalids that accompany the holmiids in the *Holmia*-*Schmidtiellus* Assemblage Zone in the HCM are represented by *Berabichia* Geyer, 1990b, *Strenuella* Matthew, 1887 emend. Hupé, 1953, and *Termierella* Hupé, 1953 (Żylińska *et al.* in press, and systematic part below). The two recognized species of *Berabichia* from the Anti-Atlas and High Atlas of Morocco, *B. vertumnia* Geyer, 1990b and *B. stenometopa* Geyer, 1990b (see Żylińska *et al.* in press, for summary), and a number of morphotypes left in open nomenclature, are most abundant in the *Antatlasia guttapliviae* and *Sectigena* zones (Geyer 1990b). *Berabichia wilkesi* (Palmer and Gatehouse, 1972) and *Berabichia subdita* (Palmer and Gatehouse, 1972), from the Argentina Range in Antarctica, are known from erratic boulders that apparently correspond to the Banian of Morocco (Palmer and Gatehouse 1972; Palmer and Rowell 1995). *Berabichia milleri* Westrop (in Westrop and Landing, 2000) from southern New Brunswick, Canada, occurs in the *Kingaspidoidea* cf. *K. obliquoculatus* Zone, thus indicating a younger level than its occurrences in Morocco (Westrop and Landing 2000). The faunule with *Berabichia erratica* Geyer, Popp, Weidner and Förster, 2004, from Pleistocene erratic boulders in a gravel pit

system	series	zones	W	formations	E
CAMBRIAN	SERIES 2	<i>Protolenus-Issafeniella</i>		Ocieszeński Sandstones (part)	 Kamieniec Shales
		<i>Holmia-Schmidtiellus</i>			
	TERRE-NEUVIAN (part)	barren interzone			?

 trilobite assemblages       acritarch assemblages

Text-fig. 4. Stratigraphic occurrence of trilobite faunas and acritarch assemblages in Cambrian Series 2 strata of the Ocieszeński and Kamieniec formations. Trilobite and acritarch assemblages from the *Protolenus*-*Issafeniella* Zone were defined and described in Żylińska and Szczepanik (2009); the acritarch assemblage from the *Holmia*-*Schmidtiellus* Assemblage Zone of the Kamieniec Formation is from Szczepanik and Żylińska (2012). The trilobite faunas from the Ocieszeński and Kamieniec formations discussed herein are marked as 1 and 2, respectively

in northern Germany is considered to correspond to the upper part of the traditional *Holmia kjerulfi*-group Zone of Sweden (Geyer *et al.* 2004). *Berabichia rotundata* (Kiær, 1917), from Scandinavia, is known from beds that yield also *Holmia kjerulfi* (Kiær 1917; Ahlberg and Bergström 1978). Sdzuy (1995) reported *Berabichia* from the Cantabrian Mountains (northern Spain), from the lower member of the Láncara Formation. Álvaro (2007) described *Berabichia eslanensis* Álvaro, 2007, from the same horizon. The age of these occurrences is early Bilbilian (Gozalo *et al.* 2007), i.e. they correlate with the upper part of the *Sectigena* Zone of Morocco and the *Protolenus–Issafeniella* Zone of the HCM in Poland (Żylińska and Szczepanik 2009). *Strenuella* is another ellipsocephalid from the study interval that has a relatively wide distribution. A number of *Strenuella* species have been recognized in Comley, Shropshire, in strata generally corresponding to the *Strenuella sabulosa* Zone of Newfoundland (Fletcher 2006; Rushton *et al.* 2011); however, *Strenuella? spinosa* (Cobbold, 1910) has been noted from the Ac2 horizon in the Comley Limestones (Morris 1988), thus, in slightly older strata, corresponding to the upper part of the *Callavia* Zone. Another *Strenuella* from the *Callavia* Zone mentioned by Morris (1988), *Strenuella comleyensis* Hupé, 1953, is considered by Geyer (1990b) to be a close relative of *Pruvostinoidea angustilineatus* Hupé, 1953. In Massachusetts, USA, *Strenuella strenua* (Billings, 1874) is common in beds yielding *Serrodiscus bellimarginatus* (Shaler and Forster, 1888), suggesting a younger age of these fossils, although Rushton (1966) noted *S. bellimarginatus* from the upper *Callavia* Zone in the Purley Shales of Nuneaton. The Moroccan *Strenuella* specimens of Hupé (1953), *S. rasetti* Hupé, 1953 and *S. howelli* Hupé, 1953, have been referred by Geyer (1990b) to *Pruvostinoidea angustilineatus* and *Ornamentaspis triangularis* (Hupé, 1953), respectively. In Spain, the Marianian Stage commences with the first appearance of *Strenuella* (Liñán *et al.* 2002; Gozalo *et al.* 2003). Species of *Termierella* have been recognized in Morocco in the *Sectigena* and *Hupeolenus* zones (Hupé 1953; Geyer 1990b), and in Spain, in the Marianian Stage (Sdzuy 1961, 1962; Gozalo *et al.* 2003).

The species of *Acanthomicmacca* Hupé, 1953, are known from several localities in Avalonia and Gondwana. *Acanthomicmacca walcotti* (Matthew, 1899) has been noted from the *Callavia broeggeri* Zone of Newfoundland (Fletcher 2006) and the *Olenellus* Limestone (Ac2) at Comley (Cobbold 1931). *Acanthomicmacca* sp. aff. *A. ellipsocephaloides* (Cobbold, 1910) is known

from the overlying *Dipharus attleborensis* Zone in Newfoundland (Fletcher 2006), and *A. sp. cf. A. ellipsocephaloides* (Cobbold, 1910) from the Branchian Series of New Brunswick (Landing *et al.* 2008). Hupé (1953) and Geyer and Malinky (1997) described *Acanthomicmacca neltneri* Hupé, 1953, from the *Marocconus notabilis* Zone of Morocco. Horizons Ac1 and Ac2 of the Comley Limestone at Comley yielded *Acanthomicmacca comleyensis* (Cobbold, 1931) and *A. ellipsocephaloides* (Cobbold, 1910), respectively (Cobbold 1931; Morris 1988), i.e. in strata corresponding to the *Callavia* Zone (Rushton *et al.* 2011). *Micmacca* (*Micmacca*) *coloi* Hupé, 1953 is also probably a representative of *Acanthomicmacca*; it was described from the *Sectigena* Zone of Morocco (Hupé 1953). *Micmacca* aff. *coloi* Hupé, 1953 has been noted in Spain (Sdzuy 1961).

The genus *Atops* Emmons, 1844 is essentially known from Laurentia. However, the *Atops*-group trilobites are also known from the successions of Avalonia and West Gondwana. In Britain, *Pseudatops reticulatus comleyensis* (Cobbold, 1936) was noted in horizon Ac3 of the Comley Limestone (Morris 1988). Fletcher (2006) illustrated *Pseudatops reticulatus* (Walcott, 1890) from the *Dipharus attleborensis* Zone of Newfoundland. Gozalo *et al.* (2003) reported *Atops* as one of the trilobites typical of the Marianian Stage; indeed, *Atops? calanus* Richter and Richter, 1941 and fragments doubtfully assigned to this species, are known from southern Spain (Richter and Richter 1941; Sdzuy 1962).

In conclusion, all trilobites of the *Homia–Schmidtellus* Assemblage Zone in the HCM indicate correlation with the traditional *Holmia kjerulfi*-group Zone of Scandinavia, the *Callavia* Zone of Britain and Newfoundland, the lower and middle parts of the *Sectigena* Zone of Morocco and the Marianian Stage of Spain.

#### BIOGEOGRAPHIC SIGNIFICANCE OF THE STUDIED FAUNA

The discussed fauna is composed of holmiids which allow direct correlation with the Scandinavian successions and thus show a strong Baltic biogeographic affinity, and of ellipsocephalids associated with a chengkouiid and an atopid, which indicate strong similarity to faunas of West Gondwana and Avalonia<sup>1</sup> (Text-fig. 5). Interestingly, the most abundant species (of *Berabichia* and *Strenuella*) are either common to Baltica, Avalonia and West Gondwana or only to Avalonia and West Gondwana. When compared to the time-

<sup>1</sup> Herein, Avalonia is treated as a microcontinent separate from Gondwana already during the Cambrian, following Landing (2005) and references therein.

genus	biogeographic area			
	Baltica	Avalonia	West Gondwana	HCM
<i>Holmia</i>	+			+
<i>Schmidtellus</i>	+			+
<i>Kjerulfia</i>	+		+	+
<i>Postfallotaspis</i>			?	+
<i>Berabichia</i>	+	+	+	+
<i>Strenuella</i>		+	+	+
<i>Termierella</i>			+	+
<i>Acanthomicmacca</i>		+	+	+
<i>Atops</i>		+	+	+

Text-fig. 5. Distribution of trilobite genera from the *Holmia*–*Schmidtellus* Assemblage Zone in the HCM and in other biogeographic areas. See text for references

equivalent successions of Baltica or Avalonia/West Gondwana, the HCM assemblage completely lacks eodiscoids; these appear later, in the *Protolenus*–*Issafeniella* Zone (see Żylińska and Szczepanik 2009 and references therein).

The close proximity of the Małopolska Massif (basement of the Kielce Region of the HCM) to Baltica was discussed in a number of papers; the most recent conclusions were based on provenance of clastic material, palaeomagnetism and biogeography (Nawrocki and Poprawa 2006; Nawrocki *et al.* 2007). During the Cambrian, the HCM edge of the Małopolska Massif was apparently a passive continental margin and was in a position close to Baltica, as evidenced by its Baltic Apparent Polar Wander Path (Nawrocki 2006; Nawrocki *et al.* 2007). Muscovites from the clastic facies of the HCM have a signature suggesting late Neoproterozoic cooling ages, but contain also an igneous-metamorphic overprint of c. 0.8–0.9 and 1.5 Ga, interpreted by Nawrocki *et al.* (2007) as a Fennoscandian source. On the other hand, dextral relocation of the Małopolska Massif along the TESZ margin of Baltica resulted in its close proximity also to the margin of Gondwana (Belka *et al.* 2002; Nawrocki *et al.* 2007), as suggested for the boundary

interval of Cambrian Series 2 and 3 by detailed analysis of trilobites and acritarchs (Żylińska and Szczepanik 2009). This study brings further support for this conclusion. The dominance of Avalonian and West Gondwanan trilobites in the succession, with the proliferation of single species over a large area suggest that the HCM margin of the Małopolska Massif belonged to a region with a mixed fauna and with an interchange of faunal types characteristic of both West Gondwana and Avalonia. This also suggests a continuous influence of ellipsocephalid larvae-carrying currents between Małopolska, West Gondwana and Avalonia. Therefore the TESZ margin of Baltica, along which the Małopolska Massif [together with other constituents of the Teisseyre Terrane Assemblage of Nawrocki and Poprawa (2006) and the Teisseyre-Tornquist Terrane Assemblage of Nawrocki *et al.* (2007)], must have faced West Gondwana and Avalonia during Cambrian epochs 2 and 3. This is a view different from that expressed by the recent palaeogeographic interpretations of Cocks and Torsvik (2005, 2006), in which the TESZ margin of Baltica faced Laurentia and was subjected to currents of the Iapetus Ocean and not of the Ægir Sea or Panthalassic Ocean as was Gondwana with Avalonia. Álvaro *et al.* (2003) proposed a pattern of currents along the margin of Gondwana. However, apart from their applied palaeogeographic reconstruction requiring emendation, the pattern of currents shown in their reconstruction does not explain the migration of faunas along the margin of Gondwana, a fact that was already brought up by Żylińska and Masiak (2007).

The dispersal potential of the holmiids and the ellipsocephalids must have been very different. The ellipsocephalids, with pandemic distribution (e.g. Geyer 1990b), could have been characterized by a long-lived planktonic early stage (i.e. all protaspid growth stages) (strategy I of Chatterton and Speyer 1989). In contrast, the holmiids were much more endemic, with *Holmia* and *Schmidtellus* confined to Baltica; *Callavia* Matthew, 1897 to Avalonia; *Andalusiana* Sdzuy, 1961 and *Cambropallas* Geyer, 1993 to West Gondwana; and *Esmeraldina* Resser and Howell, 1938, *Holmiella* Fritz, 1972 and *Palmettaspis* Fritz, 1995 to Laurentia. Only *Kjerulfia* had a wider distribution, being known from Baltica, West Gondwana and Avalonia (Palmer and Repina 1993). Holmiids apparently had a much lower dispersal potential than ellipsocephalids, possibly with an early benthic strategy of the protaspids (e.g. Chatterton and Speyer 1989). Owing to the poorly known early growth stages of the Olenellina (see e.g. Chatterton and Speyer in Kaesler 1997), this hypothesis cannot be proved.

## CONCLUSIONS

The oldest Cambrian trilobite fauna from the Holy Cross Mountains, Poland, is composed of six species of Holmiidae, four species of Ellipsocephalidae, one chengkouiid and one atopid. It is dominated by the ellipsocephalids (78%), with the abundance of *Berabichia oratrix* exceeding half of the known specimens. Biostratigraphic analysis of the taxa indicate correlation with the traditional *Holmia kjerulfi*-group Zone of Scandinavia, the *Callavia* Zone of Britain and Newfoundland, the lower and middle part of the *Sectigena* Zone of Morocco, and the Marianian Stage of Spain. The taxa display strong biogeographic affinities with West Gondwana and Avalonia, suggesting that the TESZ margin of Baltica with the Małopolska Massif must have been subjected to planktonic larvae-carrying currents effectively dispersing ellipsocephalid trilobites from and to West Gondwana and Avalonia.

## PALAEOLOGICAL NOTES

## Material and methods

The specimens studied herein are preserved as internal moulds or external moulds and their imprints; composite moulds have also been noted. Fossils from fine siliciclastics are usually flattened and in this case the measurable dimensions may not be reliable. Disarticulated sclerites prevail, dominated by cranidia, but trilobites from both formations are represented by a relatively high number of complete exoskeletons compared to other Cambrian successions of the HCM.

Detailed systematic diagnoses and descriptions are supplied only for taxa whose taxonomic assignment is revised herein or required emendation. In the remaining cases synonymy lists have been supplemented and taxon authorships are discussed. Appendix 1 summarizes the distribution of trilobite specimens in particular exposures with their repository numbers.

The terms applied to the trilobite exoskeleton follow the Trilobite Treatise (Kaesler 1997). Measurements were made with digital callipers (0.1 mm accuracy). Character lengths were measured either sagittally (sag.) or exsagittally (exs.), and widths were measured transversely (tr.). For each specimen, the measurements were taken in one plane. Particular parameters were calculated with Microsoft Excel software. A value range is given for two to four measured specimens, whereas for five specimens or more the mean value together with the standard deviation is given; the number of measured specimens is given in parentheses (e.g.

n=6). Before being photographed, the specimens were coated with ammonium chloride. In some cases several photographs of a single specimen were stacked into one with use of computer software (CombineZM or Helicon Focus 5.3).

## Systematic descriptions

Class Trilobita Walch, 1771  
Order Redlichiida Richter, 1932  
Suborder Olenellina Walcott, 1890  
Superfamily Olenelloidea Walcott, 1890  
Family Holmiidae Hupé, 1953  
Subfamily Holmiinae Hupé, 1953

Genus *Holmia* Matthew, 1890

TYPE SPECIES: *Paradoxides kjerulfi* Linnarsson, 1871 from Cambrian Series 2 in the Ringsaker district, Norway, by original designation.

REMARKS: Emended diagnoses of the genus were presented by Bergström (1973b), Ahlberg *et al.* (1986), Palmer and Repina (1993) and Palmer and Repina in Kaesler (1997). The most recent emendation is by Lieberman (1999) based on phylogenetic analyses; however, the Polish specimens of this genus were not included in the analysis (Lieberman 1999, p. 73).

*Holmia marginata* Orłowski, 1974  
(Text-figs 6A–D, 13E)

- ?1918. *Holmia kjerulfi* Linnarss.; Samsonowicz, pp. 702, 705.  
?1919. *Olenellus (Holmia) kjerulfi* Lns.; Czarnocki, pp. 94–96.  
*partim* 1959a. *Holmia kjerulfi* (Lns.); Samsonowicz, pp. 447, 449, pl. 1, figs 1, 3, 5, 6, *non* pl. 1, fig. 2 (= *Kjerulfia orcina*), *non* pl. 1, fig. 4 and pl. 2, fig. 3 (= *Holmia glabra*), *non* pl. 1, figs 7–10 and pl. 2, figs 4–11.  
1974. *Holmia kjerulfi marginata* subsp. n.; Orłowski, pp. 8–10, pl. 1, figs 1–4, pl. 2, figs 1–6.  
1985a. *Holmia marginata* Orłowski; Orłowski, pp. 236, 237, text-fig. 4, pl. 1, figs 1–4.  
1990. *Holmia kjerulfi marginata* Orłowski; Lenzion and Orłowski in Pajchłowa, p. 51, pl. 12, fig. 2.  
1999. *Holmia kjerulfi marginata* Orłowski 1974; Lieberman, p. 75.  
2003. *Holmia marginata*, Orłowski, 1974; Ebbestad, Ahlberg and Høyberget, pp. 1039, 1043.



**HOLOTYPE:** Cephalon MWG ZI/29/0991 (former number: IGP UW 55), illustrated by Orłowski (1974, pl. 1, fig. 1a, b) and Lenzion and Orłowski (in Pajchłowa 1990, pl. 12, fig. 2a, b) as *Holmia kjerulfi marginata*, and in Text-fig. 6A herein, from the *Holmia–Schmidtellus* Assemblage Zone of the Ociesęki Formation at Igrzyczna Hill, HCM.

**AVAILABLE MATERIAL:** Over 100 cephalata (incomplete in many cases), including the holotype, occasionally with incomplete thoraces, 7 hypostomata, 7 librigenae, 4 pygidia, 3 thoraces and a number of fragmentary specimens.

**REMARKS:** This well-known species from the HCM area is closely related to *Holmia kjerulfi* from the traditional lower Cambrian of Scandinavia, as noted in the diagnosis and detailed description by Orłowski (1974). Based on well-preserved specimens, Orłowski (1985a) treated it as a separate species of *Holmia*, instead of as a subspecies of *H. kjerulfi*. This conclusion was confirmed by Ebbestad *et al.* (2003). The species is diagnostic of the *Holmia–Schmidtellus* Assemblage Zone (Orłowski 1987). New photographic documentation is provided herein.

**OCCURRENCE:** Cambrian Series 2, *Holmia–Schmidtellus* Assemblage Zone; Ociesęki Formation at Igrzyczna Hill, Leśniakowa Dębina, Malkowska Hill, Ociesęki, Sterczyna Hill, Stołowa Hill, Szczecno and Wola Wąkopna; Kamieniec Formation at Dąb, HCM.

*Holmia glabra* Orłowski, 1974  
(Text-fig. 6E)


*partim* 1959a. *Holmia kjerulfi* (Lnr.); Samsonowicz, pp. 447, 449, pl. 1, fig. 4; pl. 2, fig. 3, *non* pl. 1, figs 1, 3, 5, 6 (= *Holmia marginata*), *non* pl. 1, fig. 2 (= *Kjerulfia orcina*), *non* pl. 1, figs 7–10 and pl. 2, figs 4–11.

1974. *Holmia glabra* sp. n.; Orłowski, pp. 10–12, pl. 2, figs 7–9.

1990. *Holmia glabra* Orłowski, 1974; Lenzion and Orłowski in Pajchłowa, p. 51, pl. 11, fig. 5.

2003. *Holmia glabra* Orłowski, 1974; Ebbestad, Ahlberg and Høyberget, pp. 1039, 1043.

**TYPES:** Designated as holotype was specimen IGP UW 5 (a cephalon with incomplete thorax), illustrated in Orłowski (1974, pl. 2, fig. 7a–c) and Lenzion and



Text-fig. 6. Holmiinae from the *Holmia–Schmidtellus* Assemblage Zone in the Ociesęki Formation, Holy Cross Mountains. **A–D.** *Holmia marginata* Orłowski, 1974 from Igrzyczna Hill. A – MWG ZI/29/0991, holotype, original of Orłowski (1974, pl. 1, fig. 1); B – MWG ZI/29/0993, original of Orłowski (1974, pl. 1, fig. 2); C – MWG ZI/29/1009; D – MWG ZI/29/1025, original of Orłowski (1985a, pl. 1, fig. 2). **E** – *Holmia glabra* Orłowski, 1974 from Ociesęki, neotype, latex cast of specimen MWG ZI/42/153, original of Samsonowicz (1959a, pl. 1, fig. 4). **F** – *Holmia* sp. from Gieraszwowice, MWG ZI/42/292; original of Samsonowicz (1959a, pl. 1, fig. 7).

Scale-bar represents 0.5 mm

Orłowski (in Pajchłowa 1990, pl. 11, fig. 5), from the *Holmia–Schmidtiellus* Assemblage Zone in the Ociesęki Formation at Bukówki-Rybnica, HCM. However, the specimen is probably lost. Consequently, the fragmentary cephalon MWG ZI/42/153 (former number Os 84) (illustrated in Samsonowicz 1959a, pl. 1, fig. 4, and in Text-fig. 6E herein), from the *Holmia–Schmidtiellus* Assemblage Zone of the Ociesęki Formation, Ociesęki, HCM, is designated the neotype.

AVAILABLE MATERIAL: One cephalon (which is the neotype), one librigena.

REMARKS: The diagnosis of the species and features differentiating it from *H. marginata* were presented in the original description (Orłowski 1974, pp. 10–12). Ebbestad *et al.* (2003) also included this species in their definition of the genus *Holmia*.

OCCURRENCE: Cambrian Series 2, *Holmia–Schmidtiellus* Assemblage Zone; Ociesęki Formation at Ociesęki, HCM.

Genus *Schmidtiellus* Moberg in Moberg and Segerberg, 1906


TYPE SPECIES: *Olenellus mickwitzi* Schmidt, 1888, from the *Schmidtiellus mickwitzi* Zone of Cambrian Series 2 of Estonia; by original designation.

REMARKS: Emended diagnoses of the genus were given in Bergström (1973b), Ahlberg *et al.* (1986), Palmer and Repina (1993) and Palmer and Repina in Kaesler (1997). Based on phylogenetic analysis, Lieberman (1999) supplied the most recent emended diagnosis; the Polish specimens of this genus were not included in his analysis.

*Schmidtiellus panowi* (Samsonowicz, 1959a)  
(Text-fig. 7B, C)

1959a. *Holmia panowi* nova species; Samsonowicz, pp. 449, 450, pl. 2, fig. 12a, b.

1973b. *Holmia panowi* Samsonowicz; Bergström, p. 295.



Text-fig. 7. Holmiinae from the *Holmia–Schmidtiellus* Assemblage Zone in the Ociesęki Formation, Holy Cross Mountains. **A, D, E** – *Schmidtiellus nodosus* Orłowski, 1985a. **A** – MWG ZI/29/1322, original of Orłowski (1985a, pl. 2, fig. 2a), Igrzyczna Hill; **D** – MWG ZI/29/1330, original of Orłowski (1985a, pl. 3, fig. 2), Leśniakowa Dębina; **E** – MWG ZI/29/1327, Sterczyna Hill. **B, C** – *Schmidtiellus panowi* (Samsonowicz, 1959a). **B** – MWG ZI/29/1338, original of Orłowski (1985a, pl. 3, fig. 4), Leśniakowa Dębina; **C** – MUZPIG 1.II.94, holotype, original of Samsonowicz (1959a, pl. 2, fig. 12a) and Orłowski (1974, pl. 1, fig. 5a), Ryj Hill.

Scale-bar represents 0.5 mm

- 1973b. *Schmidtellus panowi* (Samsonowicz); Bergström, pp. 301, 303.
1973. *Holmia grandis* Kiaer; Osmólska after Kotas, p. 37.
1974. *Schmidtellus panowi* (Samsonowicz, 1959); Orłowski, pp. 7, 8, pl. 1, figs 5a, b, 6.
- 1975a. *Schmidtellus panowi* (Samsonowicz, 1959); Orłowski, pp. 380, 381, pl. 1, fig. 1a, b.
- 1985a. *Schmidtellus panowi* (Samsonowicz, 1959); Orłowski, p. 237, text-fig. 5, pl. 3, fig. 4.
1990. *Schmidtellus panowi* (Samsonowicz, 1959); Lendzion and Orłowski in Pajchłowa, p. 52, pl. 13, figs 1, 2.
1999. *Schmidtellus panowi* (Samsonowicz); Lieberman, p. 82.
2003. *Schmidtellus panowi* (Samsonowicz, 1959); Ebbestad, Ahlberg and Høyberget, p. 1039.
2004. *Schmidtellus panowi* (Samsonowicz); Żylińska in Nawrocki *et al.*, text-fig. 6.

**HOLOTYPE:** The incomplete specimen, MUZPIG 1.II.94, illustrated in Samsonowicz (1959a, pl. 2, fig. 12a, b) as *Holmia panowi* nova species, Orłowski (1974, pl. 1, fig. 5a, b), Lendzion and Orłowski (in Pajchłowa 1990, pl. 13, fig. 1a, b) and Text-fig. 7C herein, from the *Holmia–Schmidtellus* Assemblage Zone in the Ociesęki Formation at Ryj Hill near Bardo, HCM. This specimen was erroneously considered the lectotype by Orłowski (1974) and Lendzion and Orłowski (in Pajchłowa 1990).

**AVAILABLE MATERIAL:** Two almost complete exoskeletons (including the holotype), two cranidia, one pleura.

**REMARKS:** Diagnosis and detailed descriptions of the species are provided by Samsonowicz (1959a) and Orłowski (1974, 1985a). This rare species provides direct correlation between the HCM and Upper Silesia. New photographic documentation of the holotype and the other exoskeleton is provided herein.

**OCCURRENCE:** Cambrian Series 2, *Holmia* Zone; Goczałkowice Formation in the Goczałkowice IG-1 Borehole (depth 2793.0 m), Cracow–Silesia region; Cambrian Series 2, *Holmia–Schmidtellus* Assemblage Zone; Ociesęki Formation at Gierasowice, Leśniakowa Dębina and Ryj Hill; Kamieniec Formation at Dąb, HCM.

*Schmidtellus nodosus* Orłowski, 1985a  
(Text-fig. 7A, D, E)

- 1985a. *Schmidtellus nodosus* sp. n.; Orłowski, pp. 237, 238, text-fig. 5, pl. 2, figs 1, 2, pl. 3, figs 2, 3.

1999. *Schmidtellus nodosus* Orłowski; Lieberman, p. 82.
2003. *S. nodosus* Orłowski, 1985; Ebbestad, Ahlberg and Høyberget, pp. 1039, 1049.

**HOLOTYPE:** Cephalon MWG ZI/29/1318 (former number: IGP UW 1.349), illustrated in Orłowski (1985a, pl. 2, fig. 1), from the *Holmia–Schmidtellus* Assemblage Zone, Ociesęki Formation at Malkowska Hill, HCM.

**AVAILABLE MATERIAL:** Thirteen cephalata, including the holotype, two cephalata with thoraces, one librigena, one hypostome, three pleurae, one pygidium tentatively assigned to the species.

**REMARKS:** The species differs from most of the representatives of *Schmidtellus* in possessing only a short occipital spine instead of a prominent one (e.g. Orłowski 1985a; Ebbestad *et al.* 2003). The spine, visible only on imprints of external moulds, is round-based in smaller specimens and changes into a more prominent ridge on the occipital ring in larger specimens.

**OCCURRENCE:** Cambrian Series 2, *Holmia–Schmidtellus* Assemblage Zone; Ociesęki Formation at Igrzyczna Hill, Leśniakowa Dębina, Malkowska Hill, Pranie and Sterczyna Hill, HCM.

Genus *Postfallotaspis* Orłowski, 1985a

**TYPE SPECIES:** *Postfallotaspis spinatus* Orłowski, 1985a, by monotypy.



Text-fig. 8. *Postfallotaspis spinatus* Orłowski, 1985a from the *Holmia–Schmidtellus* Assemblage Zone in the Ociesęki Formation at Igrzyczna Hill, Holy Cross Mountains, MWG ZI/29/2130, holotype, original of Orłowski (1985a, pl. 3, fig. 5a–d). Scale-bar represents 0.5 mm

*Postfallotaspis spinatus* Orłowski, 1985a  
(Text-fig. 8)

1985a. *Postfallotaspis spinatus* sp. n.; Orłowski, p. 235, text-fig. 3, pl. 3, fig. 5.

HOLOTYPE: Cephalon MWG ZI/29/2130 (former number: IGP UW 1.496), illustrated in Orłowski (1985a, pl. 3, fig. 5) and in Text-fig. 8 herein, from the *Holmia-Schmidtellus* Assemblage Zone in the Ociesęki Formation at Igrzyczna Hill, HCM.

AVAILABLE MATERIAL: One incomplete cephalon.

REMARKS: According to Palmer and Repina (1993, p. 32, and in Kaesler 1997, p. 428), it is impossible to ascertain a higher-level taxonomy of this genus and species; these authors placed it in the Superfamily Uncertain in the Suborder Olenellina. According to Jell and Adrain (2003, p. 429), the genus is a representative of the Holmiidae and this view is followed herein. The specimen at hand is distinguished from most of the other Holmiidae by a forwardly tapering glabella. In this respect it resembles specimens assigned to *Iyouella* Geyer and Palmer, 1995, represented so far by only one, incompletely preserved cephalon of *Iyouella contracta* Geyer and Palmer, 1995, from the *Sectigena* Zone, Issafen Formation in Morocco (Geyer and Palmer 1995). Due to the lack of other specimens, detailed comparison with other taxa is not possible.

OCCURRENCE: Cambrian Series 2, *Holmia-Schmidtellus* Assemblage Zone; Ociesęki Formation at Igrzyczna Hill, HCM.

Subfamily Callaviinae Poulsen in Moore, 1959

Genus *Kjerulfia* Kiær, 1917

TYPE SPECIES: *Kjerulfia lata* Kiær, 1917 from Cambrian Series 2 of Tømten, Norway, by original designation.

REMARKS: Emended diagnoses presented by Ahlberg *et al.* (1986), Palmer and Repina (1993) and Lieberman (1999); the Polish species is excluded from the latter analysis.

*Kjerulfia orcina* Orłowski, 1974  
(Text-fig. 9)

*partim* 1959a. *Holmia kjerulfi* (Lnrs.); Samsonowicz, pl. 1, fig. 2, *non* pl. 1, figs 1, 3, 5, 6 (= *Holmia marginata*), *non* pl. 1, fig. 4 and pl. 2, fig. 3 (= *Holmia glabra*), *non* pl. 1, figs 7–10 and pl. 2, figs 4–11.

*partim* 1959a. *Kjerulfia* sp.; Samsonowicz, p. 450, pl. 3, figs 1, 4, 5, *non* pl. 3, figs 2, 3, 6–9.

*partim* 1959a. *Kjerulfia? lagowiensis* Czarnocki; Samsonowicz, p. 450, pl. 3, fig. 11 (only).

1974. *Kjerulfia orcina* sp. n.; Orłowski, pp. 13–15, pl. 3, figs 4, 5, pl. 4, figs 1–8, pl. 5, figs 1–5.

1985a. *Kjerulfia orcina* Orłowski; Orłowski, p. 234, text-fig. 2, pl. 1, figs 5, 6.


1990. *Kjerulfia orcina* Orłowski; Lenzion and Orłowski in Pajchłowa, p. 50, pl. 11, figs 1, 2.

HOLOTYPE: Cephalon MWG ZI/29/1083 (former number: IGP UW 180), illustrated in Orłowski (1974, pl. 4, fig. 1a, b), Lenzion and Orłowski (in Pajchłowa 1990, pl. 11, fig. 1) and Text-fig. 9A herein, from the *Holmia-Schmidtellus* Assemblage Zone, Ociesęki Formation at Ociesęki, HCM.

AVAILABLE MATERIAL: One complete exoskeleton, over sixty cephalae (including the holotype), two cephalae with thoraces, eight thoraces, 16 pleurae, 7 hypostomata, 25 librigenae and a number of fragmentary specimens.

EMENDED DIAGNOSIS: A species of *Kjerulfia* with a clavate glabella, large palpebral lobe and transversely narrow extraocular cheek with broad cephalic border; librigenae with short, broad-based spine; thorax with transversely wide pleurae and long, falcate pleural spines; axial spines present on all thoracic segments, on which they are posteriorly located; total number of segments reaches 17; pygidium small, posteriorly rounded (after Orłowski 1974, supplemented).

REMARKS: The record of a complete, posteriorly enrolled specimen (Text-fig. 9D–F) shows that *Kjerulfia orcina* was a micropygous trilobite with 17 segments in adults and a small, posteriorly rounded pygidium without marginal spines. *Kjerulfia? lagowiensis* Czarnocki (see Samsonowicz 1959a, pl. 3, figs 10–12) is represented only by single pleurae. According to Orłowski (1974, footnote on p. 14) the taxon should be treated as a *nomen nudum*. However, Czarnocki's collection at the Museum of the Polish Geological Institute in Warsaw (MUZPIG) yields several pleurae with labels indicating this name. Some of the pleurae resemble those typical of *Kjerulfia* (e.g. Samsonowicz 1959a, pl. 3, fig. 11).



Text-fig. 9. *Kjerulfia orcina* Orłowski, 1974 from the *Holmia-Schmidtellus* Assemblage Zone in the Ociesęki Formation, Holy Cross Mountains. A – MWG ZI/29/1083, holotype, original of Orłowski (1974, pl. 4, fig. 1), Ociesęki; B – MWG ZI/29/1101, Sterczyna Hill; C – MWG ZI/29/1118, Leśniakowa Dębina; D–F – MUZPIG 1689.II.41, Sterczyna Hill: D – frontal view, E – lateral view, F – posterior view. Scale-bar represents 0.5 mm

**OCCURRENCE:** Cambrian Series 2, *Holmia-Schmidtellus* Assemblage Zone; Ociesęki Formation at Buczyzna, Igrzyczna Hill, Leśniakowa Dębina, Madeje, Malkowska Hill, Ociesęki, Sterczyna Hill and Wola Wąkopna; Kamieniec Formation at Dąb, HCM.

Suborder Redlichiina Richter, 1932  
 Superfamily Ellipsocephaloidea Matthew, 1887  
 Family Ellipsocephalidae Matthew, 1887  
 Subfamily Strenuellinae Hupé, 1953 emend. Geyer, 1990b

REMARKS: According to Geyer (1990b, p. 96), the family encompasses only six genera, *Strenuella* Matthew, 1887, *Myopostrenua* Rushton, 1966, *Comluella* Hupé, 1953 (restricted to the type species *Anomocare platycephalus* Cobbold, 1910), *Pruvostinoidea* Hupé, 1953, *Rinconia* Hupé, 1953 and *Limuolenus* Jago in Courtessole and Jago, 1980. Characteristic features of its representatives, considered relatively primitive in the ellipsocephalid stock, include a non-specialized, usually straight-sided and anteriorly rounded glabella with three lateral glabellar furrows, a narrow preglabellar field in the form of a depression and palpebral lobes passing into eye ridges without any discontinuity but only a slight change in angle.

Genus *Strenuella*

Matthew, 1887 emend. Hupé, 1953

TYPE SPECIES: *Agraulos strenuus* Billings, 1874, from Cambrian Series 2 of south-east Newfoundland; subsequent designation by Kiær (1917, p. 30).

*Strenuella polonica* Czarnocki, 1926  
(Text-fig. 10)

- ?1919. *Protolenus* sp.; Czarnocki, pp. 96, 97.  
1926. *Strenuella polonica* sp. n.; Czarnocki, p. 7.  
1927a. *Strenuella polonica*; Czarnocki, p. 736.  
1927b. *Strenuella polonica* sp. n.; Czarnocki, p. 190.  
1932. *Strenuella polonica*; Czarnocki, p. 77.  
1933. *Strenuella polonica*; Czarnocki, p. 81.  
1959c. *Strenuella polonica* Czarnocki (nomen nudum); Samsonowicz, pp. 525, 526, pl. 1, figs 1–4, ?5, ?6, 7–12.  
partim 1959c. *Strenuella* cf. *lakei* Hupé; Samsonowicz, pl. 2, fig. 3, non pl. 2 figs 1, 2 (= *Strenuella zbelutkae*).  
non 1962. *Strenuella polonica* Czarnocki (nomen nudum) 1927, Samsonowicz 1959; Samsonowicz, p. 12, pl. 3, fig. 9 (= *Termierella sandomirensis*).  
1985a. *Strenuella polonica* Samsonowicz; Orłowski, pp. 241, 242, text-fig. 11, pl. 4, figs 1–6, pl. 5, fig. 6, pl. 7, fig. 9.  
1990. *Strenuella polonica* Samsonowicz; Lendzion and Orłowski in Pajchłowa, p. 55, pl. 15, fig. 4.  
2010. *Strenuella polonica* Czarnocki; Żylińska and Kin, pp. 91–93, fig. 1.


TYPES: According to Samsonowicz (1959c, p. 525), the holotype cannot be found and is most probably lost. The neotype is a partly enrolled imprint of a composite mould (MWG ZI/42/073; former number: IGP UW Os 60) and was selected by Samsonowicz (1959c, p. 525, pl. 1, fig. 1a–e); this specimen was also illustrated by Lendzion and Orłowski in Pajchłowa (1990, pl. 15, fig. 4a, b). Bergström (1973a, p. 27) and Geyer (1990b, p. 56) referred to this specimen as the holotype. The specimen is from the *Holmia–Schmidtellus* Assemblage Zone of the Ociesęki Sandstone Formation at Ociesęki, HCM.

EMENDED DIAGNOSIS: A species of *Strenuella* with prominent glabella and anterior border; three distinct lateral glabellar furrows; anterior border furrow deflected rearwards at midline; occipital spine narrow-based, relatively short, deflected obliquely backwards; librigena with relatively long, slender, broad-based spine; thorax with median knobs, developed into backwardly-directed, curved spines on the 4th to 7th segments; pygidium small, with two pairs of small, inwardly directed marginal spines.

AVAILABLE MATERIAL: Twelve complete specimens, 5 axial shields, 32 cranidia with thoraces (including the neotype), 18 cranidia with librigenae and thoraces, 3 thoracopygons, 31 thoraces, 1 cranidium with librigenae, more than 275 cranidia, 8 librigenae, 11 pygidia, 2 hypostomata.

DESCRIPTION: Medium-sized micropygous trilobites, capable of spiral enrolment. Cranidium sub-square, overall convexity fairly low, glabella and frontal border relatively strongly elevated above the fixigenae, length including occipital spine  $95\pm 11\%$  of maximum width across centre of palpebral lobes when the occipital spine is preserved ( $n=18$ ). Glabella strongly convex, distinctly elevated above the fixigenae,  $80\pm 3\%$  of cranial length including occipital spine ( $n=21$ ) and  $36\pm 3\%$  of cranial width across occipital ring ( $n=18$ ), slightly narrowing forwards. Frontal lobe evenly rounded anteriorly. Three pairs of lateral glabellar furrows, more distinct on internal moulds, shallow and narrow, decreasing in length towards the anterior; posteriormost furrow slightly bent backwards and vaguely bifurcating, the remaining two only slightly oblique with regard to the transverse line. Oc-

Text-fig. 10. *Strenuella polonica* Czarnocki, 1926 from the *Holmia–Schmidtellus* Assemblage Zone in the Ociesęki Formation, Holy Cross Mountains. **A** – AK 300 CD 31–8, two slightly incomplete, outstretched exoskeletons, Sterczyna Hill; **B** – MWG ZI/29/1540, cephalon with incomplete, partly enrolled thorax, Sterczyna Hill; **C** – MUZPIG 1.II.56, incomplete cranidium (note imprint of internal mould of spine), Leśniakowa Dębina; **D** – MUZPIG 1.II.66, pygidium, Leśniakowa Dębina; **E** – reconstruction of cranidium; **F** – UJ 214P-T1, imprint of external mould of thorax (note holes after medial thoracic spines), Sterczyna Hill; **G** – MUZPIG 1.II.45, Leśniakowa Dębina; **H** – MUZPIG 1.II.79, Leśniakowa Dębina; **I** – MWG ZI/29/1488, axial shield with pygidium flex, Leśniakowa Dębina; **J** – MUZPIG 1.II.91, Rembów; **K** – original museum label for specimen MUZPIG 1.II.91 with the annotation: “the first specimen of Lower Cambrian fauna found in the Holy Cross Mountains”. Scale-bar represents 0.5 mm



occipital furrow shallow, wide, shallower at sagittal midline, slightly curved backward. Occipital ring  $17\pm 4\%$  of cranial length ( $n=48$ ),  $25\pm 3\%$  including occipital spine ( $n=21$ ), evenly convex and undifferentiated. Occipital spine with relatively narrow base, rapidly narrowing backwards to reach approximately posterior margin of second thoracic segment, terminating in a fine, sharp tip; broken off or incomplete in most of the studied crania. Axial furrows shallow and wide. Fixigenae almost flat, at a distinctly lower level than glabella, their posterolateral corners slightly sloping downwards towards the margins,  $76\pm 10\%$  of transverse occipital ring width ( $n=61$ ). Palpebral lobes  $27\pm 3\%$  of cranial length including occipital spine ( $n=16$ ), crescent-shaped and slightly elevated above the fixigenae, separated from them by wide and shallow palpebral furrow. Palpebral lobes separated by a very faint, almost exsagittal depression from slightly narrower eye ridges that run adaxially at a slightly changed angle. Eye ridges straight to slightly oblique, disconnected from glabella and curved adaxially close to anterior lobe of glabella. Anterior branch of facial suture almost parallel from palpebral lobes to frontal margin, then curving sharply adaxially. Posterior branch only slightly divergent, shorter than anterior branch. Frontal area  $20\pm 3\%$  of total cranial length including occipital spine ( $n=21$ ), distinctly separated into strongly inflated anterior border and flat preglabellar field. Anterior border  $63\pm 9\%$  of frontal area ( $n=61$ ); anterior border furrow slightly deflected rearwards. Posterior border furrow wide and shallow, wider abaxially. Posterior border with fulcrum located at two-thirds of its length from axial furrow.

Librigena almost flat, transversely very narrow, with relatively wide border extending into prominent, broad-based spine that is almost as long as the exsagittal length of the librigenal field. Genal spine extending posteriorly to the level of the 4th thoracic segment. Lateral border furrow well defined. Librigenal doublure visible on internal or composite moulds, as wide as librigenal border. Librigenal spine following the curvature of the librigenal margin, extended slightly outwards.

Thorax with 13 segments. Thoracic pleurae wider than axial ring, bent strongly downwards (in unflattened specimens) from fulcral process that is located slightly closer to axial ring than half pleural length. Pleural furrows long, shallower close to the axial ring and becoming gradually narrower and more pronounced abaxially, extending from and connected with axial furrow almost up to the abaxial tip of the pleura. Pleural terminations extending into pleural spines that are slightly curved backwards except for the posteriormost pleurae in which the spines are either not curved backwards or not developed. Axial rings in the anterior half of the tho-

rax with prominent median spine that is curved backwards and ends in a sharp tip. Subsequent spines overlap one another when thorax is outstretched. The most prominent spines present in the middle part of thorax, on the 4th to 7th segments, where they are broad-based, thick and long.

Pygidium small, nearly flat, almost twice as wide as long, with axis transversely wider than pleural region. Axis with two obsolescent axial rings, strongly tapering backward and with rounded terminal axial piece that does not reach the posterior margin of the pygidium. Articulating half-ring with lateral margins slightly convergent towards axis. Pleural regions subtriangular, almost undifferentiated except for two pleural furrows, the anterior one being more pronounced. Border flat, very poorly defined from pleural regions. Pygidial margin with two pairs of small, inwardly directed pygidial spines, of which the anterior one is relatively longer and more broad-based. Distance between posterior spines slightly wider transversely than anterior width of axis.

MEASUREMENTS: Cranial length with occipital spine 5.2–18 mm; cranial width across centre of palpebral lobes 6.7–18.8 mm ( $n=73$ ). The largest, almost complete exoskeleton is longer than 30 mm (MWG ZI/29/1519).

REMARKS: *Strenuella polonica* differs from *S. zbelutkae* Orłowski, 1985a, another species of *Strenuella* from the HCM, in a more elevated glabella and a more prominent anterior margin, more distinct lateral glabellar furrows, slightly shorter palpebral lobes in relation to cranial length, a less rounded anterior margin, an anterior border furrow deflected rearwards, and a shorter occipital spine that is less broad-based. From *Strenuella strenua* (Billings, 1874), the type species, it differs in a higher width to length ratio, a sagittally longer frontal area, an anterior border distinctly separated from preglabellar field, more distinct lateral glabellar furrows and a slightly longer occipital spine, as well as a more slender librigenal spine (see e.g., Shaw 1950, pl. 79, figs 2, 5, 14). The medially deflected anterior border furrow also seems to be a distinguishing feature in this case, although Shaw (1950, p. 580) noted a thickening of the border in *S. strenua* from Attleboro, Massachusetts, USA, as a feature of larger crania and ascribed it to maturity or geronticism. Thoraces of *S. polonica*, known from over 80 specimens, are very similar to the thorax assigned to *S. strenua*, illustrated originally by Shaler and Foerste (1888, pl. 2, fig. 21) and re-illustrated by Shaw (1950, pl. 79, figs 11–13), particularly in the arrangement of the thoracic spines (see description in Shaw 1950, p. 580). A distorted but complete specimen of *Strenuella* sp.



from Cierro del Hierro (southern Spain) (Mayoral *et al.* 2008, fig. 6g) shows the same arrangement of thoracic medial spines. Likewise, distorted thoraces of *Strenuella* (*Myopostrenua*) *sabulosa* Rushton, 1966 (now considered a separate genus, see Geyer 1990b, p. 96), illustrated in Bullock *et al.* (2011), show the same, conservative thoracic morphology, seemingly characteristic of this group of trilobites.

Despite the lack of a formal description of the species by Czarnocki, the identity of the taxon was clear to Samsonowicz (1959c), so that he was able to identify the species in old museum collections (many specimens still have original museum labels from Czarnocki's times) and in exposures. Samsonowicz (1959c) referred to Czarnocki (1927a) as the original publication in which the species was mentioned for the first time. However, the first published report with the taxon name was by Czarnocki (1926), and this year should be used with the species name. This is supported also by the fact that the paper by Czarnocki (1927a) is an abstract from the International Geological Congress held in Madrid in 1926.

According to the original museum label for specimen MUZPIG 1.II.91, *Strenuella polonica* was the first Early Cambrian trilobite found in the HCM area already in 1915 (Text-fig. 10K). This very characteristic Cambrian trilobite from the HCM was capable of spiral enrolment, a feature that was noted already by Samsonowicz (1959c), Bergström (1973a), Orłowski (1985a) and Geyer (1990b). The exoskeleton was relatively spinose, with backwardly directed spines located on the occipital ring, librigenae, almost all thoracic axial segments and thoracic pleurae, and also on the pygidial margin. When enrolled, the animal could have posed a serious challenge for potential predators of the Cambrian sea (Żylińska and Kin 2010). The mechanism of enrolment in this species along with the possible causes of such behaviour will be discussed in detail in a separate paper.

OCCURRENCE: Cambrian Series 2, *Holmia–Schmidtellus* Assemblage Zone; Ociesęki Formation at Igrzyczyna Hill, Jaźwina Hill, Leśniakowa Dębina, Madeje, Ociesęki, Pranie, Rembów, Sterczyna Hill, HCM.

*Strenuella zbelutkae* Orłowski, 1985a  
(Text-fig. 11)

*partim* 1959c. *Strenuella* cf. *lakei* Hupé; Samsonowicz, pp. 526, 527, pl. 2, figs 1, 2, *non* pl. 2, fig. 3 (= *Strenuella polonica*).

1985a. *Strenuella zbelutkae* sp. n.; Orłowski, pp. 243, 244, text-fig. 13, pl. 4, figs 7, 8.

HOLOTYPE: Cranidium MWG ZI/42/035 (former numbers: IGP UW Zb2, 1.436), illustrated in Samsonowicz (1959c, pl. 2, figs 1a–c) as *Strenuella* cf. *lakei*, Orłowski (1985a, pl. 4, fig. 7) and Text-fig. 11B herein, from the *Holmia–Schmidtellus* Assemblage Zone, Kamieniec Formation at Zbelutka, HCM.


EMENDED DIAGNOSIS: A species of *Strenuella* with glabella and anterior border only slightly elevated above the fixigenae; two pairs of almost effaced, short lateral glabellar furrows; anterior border furrow slightly less pronounced medially; occipital furrow less pronounced medially; occipital spine broad-based and long, extending into sharp tip.

AVAILABLE MATERIAL: Over 90 cranidia, including the holotype, in variable states of preservation.

DESCRIPTION: Cranidium subquadrate, overall convexity low, glabella and anterior border only slightly elevated above the fixigenae, length including occipital spine  $115 \pm 7\%$  of maximum width across centre of palpebral lobes ( $n=10$ ). Glabella slightly elevated above the fixigenae,  $81 \pm 3\%$  of cranial length including occipital spine and  $40 \pm 3\%$  of cranial width across occipital ring ( $n=10$ ); only slightly narrowing forwards. Frontal lobe slightly rounded to acute anteriorly. Two pairs of short and shallow, obsolescent lateral glabellar furrows. Occipital furrow shallow and wide, less pronounced medially, and slightly curved rearwards. Occipital ring  $20 \pm 3\%$  of cranial length ( $n=5$ ),  $29 \pm 4\%$  of cranial length including occipital spine ( $n=9$ ), of similar transverse convexity as glabella, undifferentiated. Occipital spine with stout base, narrowing evenly backwards, relatively long and terminating in a sharp, broad-based tip. Axial furrows shallow and wide. Fixigenae nearly flat, at a slightly lower level than glabella, with posterior limbs slightly deflected abaxially,  $70 \pm 8\%$  of transverse occipital ring width ( $n=10$ ). Palpebral lobes  $25 \pm 2\%$  of cranial length including occipital spine ( $n=10$ ), crescent-shaped and at a slightly higher level than the fixigenae but not upturned abaxially, separated from fixigenae by an obsolescent palpebral furrow. Palpebral lobes passing without any discontinuity but only slight change of angle into slightly elevated, very narrow eye ridges. Eye ridges slightly oblique, disconnected from glabella, passing into a vaguely marked parafrontal band. Anterior branch of facial suture almost parallel from palpebral lobes to frontal margin, then curving sharply adaxially. Posterior branch slightly divergent, short. Frontal area  $19 \pm 3\%$  of cranial length including occipital spine ( $n=10$ ), poorly separated into only slightly elevated anterior border and flat preglabellar

lar field. Anterior border about half of frontal area. Posterior furrow of medium width and shallow, becoming slightly wider abaxially. Posterior margin with fulcrum located at half width of posterior limb. Librigena, thorax and pygidium unknown.

MEASUREMENTS: Cranial length including occipital spine 11.5–21 mm; cranial width across centre of palpebral lobes 11.8–17.9 mm (n=10). Holotype cranium with occipital spine is 16.5 mm long and 14 mm wide across centre of palpebral lobes.



Text-fig. 11. *Strenuella zbelutkae* Orłowski, 1985a from the *Holmia–Schmidtellus* Assemblage Zone in the Kamieniec (K) or Ociesęki (O) formations, Holy Cross Mountains. A – MUZPIG 110.II.233, locality unknown (O); B – MWG ZI/42/035, holotype, original of Samsonowicz (1959c, pl. 2, fig. 1a–c) and Orłowski (1985a, pl. 4, fig. 7), Zbelutka (K); C – MUZPIG 113.II.2a, Zbelutka (K); D – MWG ZI/42/040, original of Orłowski (1985a, pl. 4, fig. 8); E – MUZPIG 113.II.2C, Zbelutka (K); F – reconstruction of the cranium; G – MWG ZI/42/029, original of Samsonowicz (1959c, pl. 2, fig. 2), Zbelutka (K); H – MUZPIG 113.II.4A, Zbelutka (K); I – MUZPIG 110.II.168B, laterally deformed cranium, locality unknown; J – MWG ZI/29/1286b, Zbelutka (K); K – MUZPIG 113.II.2B, Zbelutka (K); L – MUZPIG 110.II.168C, locality unknown; M – MUZPIG 110.II.160B, Ociesęki (O); N – MWG ZI/29/1289, Zbelutka (K). Scale-bar represents 0.5 mm

REMARKS: Comparison of *S. zbelutkae* with *S. polonica* is presented in the description of the latter species. From *S. strenua*, *S. zbelutkae* differs in a sagittally longer frontal area, longer palpebral lobes and a stout-based, long occipital spine. This species is one of the most abundant taxa in the Kamieniec Shales and seems to represent a species more adapted to the deeper environment of this formation.

OCCURRENCE: Cambrian Series 2, *Holmia-Schmidtellus* Assemblage Zone; Ociesęki Formation at Leśniakowa Dębina and Ociesęki; Kamieniec Formation at Dąb and Zbelutka, HCM.

Subfamily Protoleninae Richter et Richter, 1948  
emend. Geyer, 1990b

*Termierella* Hupé, 1953

TYPE SPECIES: *Termierella latifrons* Hupé, 1953 from the *Sectigena* and *Hupeolenus* zones of the Issafen Syncline, Anti-Atlas, Morocco, by original designation.

*Termierella sandomirensis* Samsonowicz, 1962  
(Text-fig. 12)

*partim* 1962. *Termierella sandomirensis* n. sp.; Samsonowicz, pp. 18, 19, pl. 1, figs 1–6, non fig. 7 (= *Berabichia oratrix*).

1962. *Strenuella polonica* Czarnocki (nomen nudum), 1927, Samsonowicz, 1959; Samsonowicz, p. 20, pl. 3, fig. 9.

1985a. *Strenuella sandomirensis* (Samsonowicz, 1962); Orłowski, pp. 242, 243, text-fig. 12, pl. 4, figs 9–11.

1990. *Termierella sandomirensis* Samsonowicz, 1962; Lendzion and Orłowski in Pajchłowa, p. 57, pl. 16, fig. 1.

HOLOTYPE: Cranidium MWG ZI/42/043 (former number: IGP UW Gr4) from the *Holmia-Schmidtellus* Assemblage Zone at Gieraszwice, HCM, illustrated in Samsonowicz (1962, pl. 1, fig. 3), Orłowski (1985a, pl. 4, fig. 11), Lendzion and Orłowski (in Pajchłowa 1990, pl. 16, fig. 1) and Text-fig. 12A herein.

EMENDED DIAGNOSIS: A small species of *Termierella* with the anteriormost pair of lateral glabellar furrows obsolescent, eye ridges widening close to glabella and occipital ring with short, broad-based spine.

AVAILABLE MATERIAL: 12 cranidia, including the holotype.

DESCRIPTION: Cranidium subtrapezoidal, overall convexity low, length 93–102% of maximum width across centre of palpebral lobes. Glabella distinctly elevated above the fixigenae, 81–88% of cranial length and 46–47% of cranial width across occipital ring; distinctly narrowing forwards. Frontal lobe bluntly rounded anteriorly. Three pairs of shallow, obliquely backward directed lateral glabellar furrows; the posteriormost furrow being the longest and relatively deepest, whereas the anteriormost is obsolescent, probably transverse and short. Occipital furrow transverse, relatively deep and narrow, slightly shallower and deflected forwards at mid-line. Occipital ring 26–29% of cranial length, of similar transverse convexity as glabella, with triangular posterior margin extending into narrow and relatively short occipital spine. Axial furrows of moderate depth, narrow and distinct. Fixigenae flat, distinctly elevated above the surrounding furrows, located at slightly lower level than glabella, about 65% of transverse occipital ring width. Palpebral lobes 27–30% of cranial length, crescent-shaped and slightly elevated above the fixigenae, separated from them by a palpebral furrow of almost the same depth as the axial furrows. Palpebral lobes passing evenly into elevated, almost transverse, wide eye ridges, generally of the same width as the palpebral lobes, then slightly broadening close to glabella and terminating at axial furrows at level of third lateral glabellar furrow. Anterior branch of facial suture only slightly divergent from palpebral lobes to frontal margin, then curving sharply adaxially, shortly anterior to the border furrow to meet the anterior margin anterior to the most elevated part of the preocular fields. Posterior branch divergent and distinctly ventrally deflected. Frontal area 12–19% of cranial length, separated into slightly elevated anterior border and flat preglabellar field. Length of anterior border about one-third of frontal area. Posterior furrow deep and narrow, only slightly wider abaxially. Posterior margin with fulcrum located at approximately half width of posterior limb. Librigena, thorax and pygidium unknown.

MEASUREMENTS: Cranial length with occipital spine 5.3–10.2 mm; cranial width across centre of palpebral lobes 8–10 mm (n=2). The lower values are of the holotype cranidium.


REMARKS: This species is known from 12 cranidia, the original specimens of Jan Samsonowicz from Gieraszwice. They also display some degree of tectonic deformation (wide and narrow specimens) and the

original width to length ratio is not clear. The reconstruction (Text-fig. 12C) is a compromise between sagittally and transversely distorted cranidia, taking into account that the illustrated specimens of other species of *Termierella* are generally wider than long (e.g. Hupé 1953, pl. 9, fig. 24, pl. 10, figs 1–4; Geyer 1990b, pl. 55, figs 4, 9, 11).

From the type species, *Termierella latifrons* Hupé, 1953, it differs in having a lower width to length ratio, a sagittally slightly longer and wider glabella, very short anteriormost lateral glabellar furrows and a longer occipital ring, an occipital spine, and in lacking grooves on the palpebral lobes. According to the footnote in Samsonowicz's posthumous paper (Samsonowicz 1962, p. 10), his manuscripts did not state to which of the subgenera of *Termierella* distinguished at that time this species belonged; an apparent comparison with Hupé's subgenera allowed assignment to *Brevitermierella* Hupé, 1953. At present, *Brevitermierella* is considered a separate genus (Geyer 1990b, p. 209), distinguished from *Termierella* by a sagittally narrow preglabellar

field and a much longer posteriormost lateral glabellar furrow. The specimens at hand have a relatively long preglabellar field sagittally, distinct furrows separating the large palpebral lobes from the fixigenae and in general proportions resemble *Termierella* rather than *Brevitermierella*. In general proportions, the specimens at hand are close to *Termierella* sp. B from Guadalcanal in southern Spain (Sdzuy 1962), but differ in a slightly wider (tr.) glabella, lack of grooves on the palpebral lobes and the presence of an occipital spine. The discussed specimens differ from another species from Spain, *Termierella seviliana* Sdzuy, 1961, in a relatively wider (tr.) glabella, less pronounced lateral glabellar furrows, lack of grooves on the palpebral lobes and anterior facial sutures that are almost parallel and do not diverge abaxially as in *T. seviliana* (Sdzuy 1961).

Orłowski (1985a) attributed these specimens to *Strenuella*, based on the lack of grooves on the palpebral lobes and the presence of an occipital spine. The grooves on the palpebral lobes are generally not present in specimens of *T. latifrons* from Morocco illustrated by Geyer



Text-fig. 12. *Termierella sandomirensis* Samsonowicz, 1962 from the *Holmia-Schmidtellus* Assemblage Zone in the Ocieski Formation at Gierasowice, Holy Cross Mountains. A – MWG ZI/42/043, holotype, slightly deformed cranidium, original of Samsonowicz (1962, pl. 1, fig. 3) and Orłowski (1985a, pl. 4, fig. 11); B – MWG ZI/42/034, original of Samsonowicz (1962, pl. 1, fig. 4) and Orłowski (1985a, pl. 4, fig. 10); C – reconstruction of the cranidium; D – MWG ZI/42/056, original of Samsonowicz (1962, pl. 1, fig. 1) and Orłowski (1985a, pl. 4, fig. 9); E – MWG ZI/42/032, original of Samsonowicz (1962, pl. 1, fig. 6); F – MWG ZI/42/053, original of Samsonowicz (1962, pl. 1, fig. 2). Scale-bar represents 0.5 mm

(1990b, pl. 55, figs 1, 8, 9; pl. 56, fig. 4); moreover, the specimens at hand are poorly preserved and do not allow the recognition of all details. Specimens from the Montagne Noire, assigned by Courtessole and Jago (1980) to “cf. *Termierella* sp. no. 1” and “cf. *Termierella* sp. no. 2”, are rather poorly preserved, but display the cranial proportions, pattern of lateral glabellar furrows and shape of palpebral lobes as in *Termierella*. These specimens do not have grooves on the palpebral lobes and “cf. *Termierella* sp. no. 2” even may have had an occipital spine (Courtessole and Jago 1980, p. 18). A specimen of *T. latifrons* illustrated in Geyer (1990b, pl. 54, fig. 7a) has a small knob on the occipital ring.

OCCURRENCE: Cambrian Series 2, *Holmia-Schmidtellus* Assemblage Zone; Ocieseński Formation at Gieraszowice, HCM.

Order Corynexochida Kobayashi, 1935  
Suborder Corynexochina Kobayashi, 1935  
Family Chengkouiidæ Zhu in Zhang *et al.*, 1980

REMARKS: Geyer and Malinky (1997, pp. 627, 628) discussed the problems with higher-level classification of the family due to mixed ellipsocephalacean and corynexochoid features of the exoskeleton in its representatives. Landing *et al.* (2008) assigned the Chengkouiidæ with doubt to Corynexochida. Allocation to the Corynexochida is also in Adrain (2011).

*Acanthomicmacca* Hupé, 1953

TYPE SPECIES: *Micmacca walcotti* Mathew, 1899, from the upper lower Cambrian basal Brigus Formation, south-east Newfoundland, Canada, by original designation.

REMARKS: The concept of the genus was discussed by Geyer and Malinky (1997). It encompasses also *Myopsomicmacca* Hupé, 1953, *Jaskovitchella* Repina in Repina *et al.*, 1975 and *Chengkouia* Qian and Yao in Lu *et al.*, 1974 (Geyer 1990b; Geyer and Malinky 1997; Jell and Adrain 2003).

*Acanthomicmacca klimontowi* Orłowski, 1985a  
(Text-fig. 13)

*partim* 1962. *Strenuella primaeva* (Brögg.); Samsonowicz, p. 11, pl. 2, fig. 5, *non* pl. 2, figs 1–4, 6, 7 (= *Behrabichia oratrix*).

1962. *Strenuella cf. lakei* Hupé; Samsonowicz, p. 12, pl. 3, figs 6–8.

1985a. *Ellipsocephalus simplex* sp. n.; Orłowski, p. 239; text-fig. 8, pl. 7, fig. 3.


1985a. *Micmacca (Acanthomicmacca) klimontowi* sp. n.; Orłowski, pp. 246, 247, text-fig. 16, pl. 7, figs 5–8.

HOLOTYPE: Cranidium MWG ZI/29/1297 (former number: IGP UW D6), illustrated in Samsonowicz (1962, pl. 3, fig. 7) as *Strenuella cf. lakei*, Orłowski (1985a, pl. 7, fig. 5) and Text-fig. 13A herein, from the *Holmia-Schmidtellus* Assemblage Zone in the Kamieniec Formation at Dąb, HCM.

EMENDED DIAGNOSIS: A species of *Acanthomicmacca* with a parallel-sided glabella both in juveniles and adults; occipital ring with terminal occipital spine; relatively long, distinct eye ridges, relatively wide preocular fields, anterior border almost flat to slightly convex in sagittal section, elevated above the preglabellar field.

AVAILABLE MATERIAL: Over ninety cranidia, including the holotype, in various preservational state.

DESCRIPTION: Cranidium subquadrate, overall convexity modest, glabella distinctly elevated above the fixigenae, length excluding occipital spine  $83 \pm 8\%$  of maximum width across centre of palpebral lobes ( $n=8$ ) and  $100 \pm 9\%$  including occipital spine ( $n=9$ ). Glabella  $77 \pm 6\%$  of cranial length including occipital spine ( $n=9$ ) and  $39 \pm 3\%$  of cranial width across occipital ring ( $n=17$ ), with parallel sides both in juveniles and adults. Frontal lobe bluntly rounded anteriorly. Lateral glabellar furrows almost effaced; in some specimens two(?) backwardly directed furrows are faintly visible. Occipital furrow obsolescent, transverse. Occipital ring  $15 \pm 2\%$  of cranial length ( $n=7$ ),  $26 \pm 7\%$  including occipital spine ( $n=7$ ), of similar transverse convexity as glabella, undifferentiated, slightly wider at sagittal midline, so that the occipital ring clearly tapers towards the axial furrows. Short, terminal occipital spine present, its length approximately the sagittal width of the occipital ring, narrowing backwards and terminating in a sharp tip. In smaller specimens the base of the spine is relatively broader than in adult individuals. Axial furrows shallow, wide, slightly shallower in front of anterior lobe. Fixigenae almost flat, their posterior limbs slightly sloping downwards abaxially,  $78 \pm 11\%$  of transverse occipital ring width ( $n=16$ ). Palpebral lobes  $26 \pm 4\%$  of cranial length including occipital spine ( $n=9$ ), crescent-shaped and nearly flat, slightly elevated above the fixigenae, separated from them by a shallow and narrow palpebral fur-



Text-fig. 13. *Acanthomicmacca klimontowi* Orłowski, 1985a from the *Holmia-Schmidtellus* Assemblage Zone in the Kamieniec (K) or Ociesęki (O) formations, Holy Cross Mountains. **A** – MWG ZI/29/1297, holotype, original of Samsonowicz (1962, pl. 3, fig. 7) and Orłowski (1985a, pl. 7, fig. 5), Dąb (K); **B** – MWG ZI/42/037, original of Orłowski (1985a, pl. 7, fig. 8), Dąb (K); **C** – MWG ZI/42/042, original of Samsonowicz (1962, pl. 4, fig. 2), Dąb (K); **D** – MUZPIG 110.II.214, Igrzyczna Hill (O); **E** – MWG ZI/29/1309, with *Holmia marginata* Orłowski, 1974 (upper specimen), Dąb (K); **F** – reconstruction of the cranidium; **G** – MUZPIG 110.II.186, Ociesęki (O); **H** – MWG ZI/42/050, original of Orłowski (1985a, pl. 7, fig. 7), Dąb (K); **I** – MWG ZI/29/1305, Dąb (K); **J** – MUZPIG 8.II.3, Julianów (K); **K** – MWG ZI/29/1302, Dąb (K); **L** – MWG ZI/29/1299, Dąb (K); **M** – MUZPIG 1.II.8, Zbelutka (K); **N** – MWG ZI/29/1307, Wola Wąkopna (O). Scale-bar equals 0.5 mm

row. Palpebral lobes evenly passing into oblique eyeridges that are slightly elevated, cord-like and disconnected from glabella, terminating approximately slightly posterior to the anterior lobe. Anterior branch of facial suture slightly divergent from palpebral lobes to frontal margin, then curving sharply adaxially. Posterior branch divergent, moderately short. Frontal area  $22\pm 6\%$  of cranial length including occipital spine ( $n=9$ ), poorly separated into only slightly elevated anterior border and flat preglabellar field. Anterior border about the sagittal length of the frontal area. Posterior furrow of medium width and very shallow, becoming slightly wider abaxially. Posterior margin with fulcrum located from axial furrows at about two-thirds width of posterior limb.

MEASUREMENTS: Cranial length with occipital spine 5.8–19 mm; cranial width across centre of palpebral lobes 5.6–20 mm ( $n=9$ ). Holotype cranidium with occipital spine is 15.8 mm long and 16 mm wide across centre of palpebral lobes.

REMARKS: *Acanthomicmacca klimontowi* was introduced by Orłowski (1985a) based on two specimens assigned by Samsonowicz (1962) to *Strenuella* cf. *lakei* Hupé, 1953. At that time, *Acanthomicmacca* Hupé, 1953 was treated as a subgenus of *Micmacca* Matthew, 1895. The morphology and concept of *Micmacca* were discussed by Westrop and Landing (2000), who restricted that genus to its type species, *Micmacca matthevi* Matthew, 1895. *Acanthomicmacca* was treated as a distinct genus by Geyer and Malinky (1997), Westrop and Landing (2000) and Jell and Adrain (2003). The familial assignment of this species is still a matter of debate; the opinion of Geyer (1990b), Geyer and Malinky (1997) and Jell and Adrain (2003) including it in the Chengkouiidae is followed herein.

*Acanthomicmacca klimontowi* differs from the type species as well as other species of the genus (see overview in Geyer and Malinky 1997, p. 631) in a higher length to width ratio (which is equal almost to 1 when the occipital spine is included), a sagittally longer and relatively flat frontal area and relatively wider fixigenae posteriorly (about four-fifths of glabellar width). Similarly as in *A. neltneri* Hupé, 1953 (e.g. Geyer and Malinky 1997, figs 7.4, 7.6), the lateral glabellar furrows are almost effaced. Both the juvenile and adult forms of *A. klimontowi* have a glabella with parallel sides, in contrast to *A. neltneri*, which so far is the only species in *Acanthomicmacca* that possesses a glabella with constricted sides in adults and an anteriorly expanding glabella in juveniles.

A number of specimens representing this species have also been discovered in Samsonowicz's and

Czarnocki's collections in the Museum of the Polish Geological Institute (MUZPIG); they were collected in Zbelutka, Olbierzowice, Julianów and Ociesęki, and assigned previously to *Strenuella* sp. or left undetermined, as is evident from the museum labels. Moreover, the holotype and only specimen of *Ellipsocephalus simplex* (Orłowski 1985a, pl. 7, fig. 3) from Kucebrza is an effaced individual of *A. klimontowi* with the occipital spine not preserved. The analyzed collections yield many effaced or spineless representatives of *A. klimontowi* whose overall appearance is very like this specimen.

OCCURRENCE: Cambrian Series 2, *Holmia-Schmidtellus* Assemblage Zone; Ociesęki Formation at Ociesęki and Wola Wąkopna; Kamieniec Formation at Zbelutka, Dąb, Julianów, Kucebrza and Olbierzowice, HCM.

Order Uncertain

Family Atopidae Hupé, 1953

Genus *Atops* Emmons, 1844

TYPE SPECIES: *Atops trilineatus* Emmons, 1844 from Cambrian Series 2 of the New York State, by monotypy.


*Atops granulatus* Orłowski, 1985c  
(Text-fig. 14)

1985c. *Atops granulatus* n. sp.; S. Orłowski, pp. 976, 977, fig. 2.

HOLOTYPE: Cephalon MWG ZI/29/3732 (former number: IGPUW/T/1), from the *Holmia-Schmidtellus* Assemblage Zone in the Ociesęki Formation at Sterczyna Hill, HCM, illustrated in Orłowski (1985c, fig. 2) and Text-fig. 14A herein.

AVAILABLE MATERIAL: One incomplete cephalon, which is the holotype, one cephalic fragment with ornamentation tentatively assigned to the species, one pleura.

REMARKS: A fragmentary internal mould of the cephalon (Text-fig. 14B) from Czarnocki's collection in the Museum of the Polish Geological Institute (MUZPIG) shows an ornamentation composed of densely distributed larger and smaller pits. The holotype cephalon of *Atops granulatus* (Text-fig. 14A) is provided with large, fairly closely packed pits together with smaller ones on the genal fields, and smaller closely packed pits on the glabella. The small cephalic fragment shows the test with



Text-fig. 14. *Atops granulatus* Orłowski, 1985c from the *Holmia-Schmidtiellus* Assemblage Zone in the Ociesęki Formation, Holy Cross Mountains. **A** – MWG ZI/29/3732, holotype, Sterczyna Hill, original of Orłowski (1985c, fig. 2); **B** – MUZPIG I10.II.154, cephalic fragment with ornamentation, Leśniakowa Dębina. Scale-bar represents 0.5 mm

its characteristic ornamentation and is tentatively assigned to the species because the larger pits seem elongate in transverse section and not round.

**OCCURRENCE:** Cambrian Series 2, *Holmia-Schmidtiellus* Assemblage Zone; Ociesęki Formation at Sterczyna Hill, HCM.

#### Acknowledgements

My warmest gratitude is expressed to Professor Stanisław Orłowski (University of Warsaw) for his everlasting encouragement. The trilobite reconstructions were skilfully drawn by Bogusław Waksmundzki (University of Warsaw). The final version of the manuscript gained significantly from the constructive reviews by Per Ahlberg (Lund University, Sweden), Gerd Geyer (Uppsala University, Sweden) and Rodolfo Gozalo (University of Valencia, Spain), for which I am extremely grateful. Mariusz Niechwedowicz (University of Warsaw), Tatiana Woroncowa-Marcinowska and Katarzyna Skurczyńska-Garwolińska (Polish Geological Institute in Warsaw) assisted with the loan of the specimens and photography. Jose-Antonio Gámez (University of Zaragoza, Spain) and Rodolfo Gozalo (University of Valencia, Spain) kindly helped with the search for old papers. The paper benefited from stimulating discussions on the palaeogeography with Wojciech Kozłowski (University of Warsaw). The research was supported in 2010–2011 by the Institute of Geology, University of Warsaw (BST grants 1536 and 160600/2).

#### REFERENCES

- Adrain, J.M. 2011. Class Trilobita Walch, 1771. In: Z.-Q. Zhang (Ed.), *Animal biodiversity: An outline of higher level classification and survey of taxonomic richness. Zootaxa*, **3148**, 104–109.
- Ahlberg, P. and Bergström, J. 1978. Lower Cambrian ptychopariid trilobites from Scandinavia. *Sveriges Geologiska Undersökning, Ser. Ca*, **49**, 5–41.
- Ahlberg, P., Bergström, J. and Johansson, J. 1986. Lower Cambrian olenellid trilobites from the Baltic Faunal province. *Geologiska Föreningens i Stockholm Förhandlingar*, **108**, 39–56.
- Álvaro, J.J. 2007. New ellipsocephalid trilobites from the lower Cambrian member of the Láncara Formation, Cantabrian Mountains, northern Spain. *Memoirs of the Association of Australasian Paleontologists*, **34**, 343–355.
- Álvaro, J.J., Elicki, O., Geyer, G., Rushton, A.W.A. and Shergold, J.H. 2003. Palaeogeographical controls on the Cambrian trilobite immigration and evolutionary patterns reported in the western Gondwana margin. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **195**, 5–35.
- Belka, Z., Ahrendt, H., Franke, W. and Wemmer, K. 2000. The Baltica–Gondwana suture in central Europe: evidence from K–Ar ages of detrital muscovites and biogeographical data. In: W. Franke, V. Haak, O. Oncken and D. Tanner (Eds), *Orogenic Processes: Quantification and Modelling in the Variscan Belt. Special Publications of the Geological Society of London*, **179**, 87–102.
- Belka, Z., Valverde-Vaquero, P., Dörr, W., Ahrendt, H., Wemmer, K., Franke, W. and Schäfer, J. 2002. Accretion of first Gondwana-derived terranes at the margin of Baltica. In: J.A. Winchester, T.C. Pharaoh and J. Verniers (Eds), *Palaeozoic Amalgamation of Central Europe. Special Publications of the Geological Society of London*, **201**, 19–36.
- Bergström, J. 1973a. Organization, life, and systematics of trilobites. *Fossils and Strata*, **2**, 1–69.



- Bergström, J. 1973b. Classification of olenellid trilobites and some Balto-Scandian species. *Norsk Geologisk Tidsskrift*, **53**, 283–314.
- Bergström, J. 1981. Lower Cambrian shelly faunas and biostratigraphy in Scandinavia. In: M.E. Taylor (Ed.), Short Papers for the Second International Symposium of the Cambrian System 1981. *United States Department of the Interior Geological Survey, Open-File Report*, **81-743**, 22–25.
- Bergström, J. and Ahlberg, P. 1981. Uppermost Lower Cambrian biostratigraphy in Scania, Sweden. *Geologiska Föreningens i Stockholm Förhandlingar*, **103**, 193–214.
- Berthelsen, A. 1992. From Precambrian to Variscan Europe. In: D. Blundell, R. Freeman and S. Muller (Eds), *A Continent Revealed: The European Geotraverse*, pp. 153–164. Cambridge University Press.
- Billings, E. 1874. Palaeozoic Fossils, 2(1). Geological Survey of Canada, 1–144.
- Bullock, R.J., Morris, J.R. and Selby, D. 2011. New findings of body and trace fossils in the St. Bride's area, Cape St. Mary's Peninsula, Newfoundland. Current research, Newfoundland and Labrador Department of Natural Resources, Geological Survey Report, **11-1**, 241–252.
- Chatterton, B.D.E. and Speyer, S.E. 1989. Larval ecology, life history strategies, and patterns of extinction and survivorship among Ordovician trilobites. *Paleobiology*, **15**, 118–132.
- Cobbold, E.S. 1910. On some small trilobites from the Cambrian rocks of Comley (Shropshire). *Quarterly Journal of the Geological Society of London*, **66**, 19–51.
- Cobbold, E.S. 1931. Additional fossils from the Cambrian rocks of Comley, Shropshire. *Quarterly Journal of the Geological Society of London*, **87**, 459–511.
- Cobbold, E.S. 1936. The Conchostraca of the Cambrian area of Comley, Shropshire, with a note on a new variety of *Atops reticulatus* (Walcott). *Quarterly Journal of the Geological Society of London*, **92**, 221–235.
- Cocks, L.R.M. and Torsvik T.H. 2005. Baltica from the late Precambrian to mid-Palaeozoic times: The gain and loss of a terrane's identity. *Earth-Science Reviews*, **72**, 39–66.
- Cocks, L.R.M. and Torsvik T.H. 2006. European geography in a global context from the Vendian to the end of the Palaeozoic. In: D.G. Gee and R.A. Stephenson (Eds), *European Lithosphere Dynamics. Memoirs of the Geological Society of London*, **32**, 82–95.
- Courtessole, R. and Jago, J.B. 1980. Biostratigraphie du Cambrien Inferieur du Cabardes (versant sud de la Montagne Noire, France méridionale). *Mémoire de la Société d'Études Scientifiques de l'Aude*, 3–26.
- Czarnocki, J. 1919. Stratigraphy and tectonics of the Święty Krzyż Mountains. *Prace Towarzystwa Naukowego Warszawskiego*, **28**, 1–172. [In Polish]
- Czarnocki, J. 1926. Cambrian stratigraphy and fauna in the central part of the Holy Cross Mountains. *Posiedzenia Naukowe Państwowego Instytutu Geologicznego*, **14**, 7–9. [In Polish]
- Czarnocki, J. 1927a. Le Cambrien et la faune cambrienne de la partie moyenne du massif de Święty Krzyż (S<sup>te</sup> Croix). Congrès Géologique International, Compte-rendue de la XIV<sup>e</sup> session, en Espagne, 1926, Madrid, 735–750.
- Czarnocki, J. 1927b. Cambrian and its fauna in the central part of the Holy Cross Mountains. *Sprawozdania Polskiego Instytutu Geologicznego*, **4**, 189–207. [In Polish]
- Czarnocki, J. 1932. Cambrian stratigraphy and tectonics in the vicinity of Ociesęki and Orłowiny. *Posiedzenia Naukowe Państwowego Instytutu Geologicznego*, **33**, 76–78. [In Polish]
- Czarnocki, J. 1933. Scientific significance of Cambrian exposures in the vicinity of Ociesęki and Orłowiny. *Zabytki Przyrody Nieożywionej*, **2**, 78–84. [In Polish]
- Ebbestad, J.O.R., Ahlberg, P. and Høyberget, M. 2003. Re-description of *Holmia inusitata* (Trilobita) from the Lower Cambrian of Scandinavia. *Palaeontology*, **46**, 1039–1045.
- Emmons, E. 1844. The Taconic System, based on observations in New York, Massachusetts, Maine, Vermont and Rhode Island. Albany, Caroll and Cook, 1–68.
- Fletcher, T.P. 2006. Bedrock geology of the Cape St. Mary's Peninsula, southwest Avalon Peninsula, Newfoundland. Government of Newfoundland and Labrador, Geological Survey, Department of Natural Resources, St. John's, Report, **06-02**, pp. 1–117.
- Fritz, W.H. 1972. Lower Cambrian trilobites from the Sekwi Formation type section, Mackenzie Mountains, north-western Canada. *Bulletin of the Geological Survey of Canada*, **212**, 1–90.
- Fritz, W.H. 1995. *Esmeraldina rowei* and associated Lower Cambrian trilobites (If fauna) at the base of Walcott's Waucoban Series, southern Great Basin, U.S.A. *Journal of Paleontology*, **69**, 708–723.
- Geyer, G. 1990a. Revised Lower to lower Middle Cambrian biostratigraphy of Morocco. *Newsletters on Stratigraphy*, **22**, 53–70.
- Geyer, G. 1990b. Die marokkanischen Ellipsocephalidae (Trilobita: Redlichiida). *Beringeria*, **3**, 3–363.
- Geyer, G. 1993. The giant Cambrian trilobites of Morocco. *Beringeria*, **8**, 71–107.
- Geyer, G. and Landing, E. 2004. A unified Lower – Middle Cambrian chronostratigraphy for West Gondwana. *Acta Geologica Polonica*, **54**, 179–218.
- Geyer, G. and Malinky, J.M. 1997. Middle Cambrian fossils from Tizi n'Tichka, the High Atlas, Morocco. Part 1. Introduction and trilobites. *Journal of Paleontology*, **71**, 620–637.
- Geyer, G. and Palmer, A.R. 1995. Neltneriidae and Holmiidae (Trilobita) from Morocco and the problem of Early Cam-

- brian intercontinental correlation. *Journal of Paleontology*, **69**, 459–474.
- Geyer, G., Popp, A., Weidner, T. and Förster, L. 2004. New Lower Cambrian trilobites from Pleistocene erratic boulders of northern Germany and Denmark and their bearing on the intercontinental correlation. *Paläontologische Zeitschrift*, **78**, 127–136.
- Gozalo, R., Liñán, E., Dies, M.E., Gámez-Vintaned, J.A. and Mayoral, E. 2007. The Lower–Middle Cambrian boundary in the Mediterranean subprovince. *Special Paper of the Geological Society of America*, **423**, 359–373.
- Gozalo, R., Liñán, E., Palacios, T., Gámez Vintaned, J.A. and Mayoral, E. 2003. The Cambrian of the Iberian Peninsula: An overview. *Geologica Acta*, **1**, 103–112.
- Hammer, Ø. and Harper, D. A. T. 2006. Paleontological data analysis. Blackwell Publishing Ltd., 1–351.
- Hupé, P. 1953. Contribution a l'étude du Cambrien inférieur et du Précambrien III de l'Anti-Atlas marocain. *Notes et Mémoires de la Service géologique du Maroc*, **103**, 1–402. ['1952']
- Jell, P.A. and Adrain, J.M. 2003. Available generic names for trilobites. *Memoirs of the Queensland Museum*, **48**, 331–553.
- Kaessler, R.L. (Ed.) 1997. Treatise on Invertebrate Paleontology, Part O, Arthropoda 1, Trilobita, Revised, pp. i–xxiv, 1–530. Geological Society of America, Boulder, and University of Kansas Press, Lawrence.
- Kiær, J. 1917. The Lower Cambrian *Holmia* Fauna at Tømten in Norway. *Videnskapselskapets Skrifter. I. Matematisk-Naturvidenskapelig Klasse*, **10**, 1–139.
- Kotas, A. 1973. Occurrence of Cambrian formations in the substratum of the Upper Silesia Coal Basin. *Przegląd Geologiczny*, **21**, 37. [In Polish]
- Kowalczewski, Z., Żylińska, A. and Szczepanik, Z. 2006. Cambrian in the Holy Cross Mountains. In: S. Skompski and A. Żylińska (Eds), *Procesy i zdarzenia w historii geologicznej Gór Świętokrzyskich. 77 Zjazd Naukowy Polskiego Towarzystwa Geologicznego*, 14–27. [In Polish]
- Krzywiec, P., Gutowski, J., Walaszczyk, I., Wróbel, G. and Wybraniec, S. 2009. Tectonostratigraphic model of the Late Cretaceous inversion along the Nowe Miasto–Zawichost Fault Zone, SE Mid-Polish Trough. *Geological Quarterly*, **53**, 27–48.
- Kutek, J. and Głazek, J. 1972. The Holy Cross area, Central Poland, in the Alpine cycle. *Acta Geologica Polonica*, **22**, 603–652.
- Landing, E., Johnson, S.C. and Geyer, G. 2008. Faunas and Cambrian volcanism on the Avalonian marginal platform, southern New Brunswick. *Journal of Paleontology*, **82**, 884–905.
- Lieberman, B.S. 1999. Systematic revision of the Olenelloidea (Trilobita, Cambrian). *Bulletin of the Peabody Museum of Natural History, Yale University*, **45**, 1–150.
- Liñán, E., Gozalo, R., Palacios, T., Gámez-Vintaned, J.A., Ugidos, J.M. and Mayoral, E. 2002. Cambrian. In: W. Gibbons and T. Moreno (Eds), *The geology of Spain*, 17–29. The Geological Society, London.
- Linnarsson, G. 1871. Om några försteningar från Sveriges och Norges 'primordialzon'. *Kongliga Svenska Vetenskaps-Akademiens Förhandlingar*, **6**, 789–796.
- Lu, Y., Zhang, W., Qian, Y., Zhu, Z., Lin, H., Zhou, Z., Zhang, S. and Yuan J. 1974. Cambrian trilobites. In: *Handbook of the Stratigraphy and Palaeontology of southwest China*, 82–107. Science Press; Beijing. [In Chinese]
- Malinowski, M., Żelaźniewicz, A., Grad, M., Guterch, A. and Janik, T. 2005. Seismic and geological structure of the crust in the transition from Baltica to Palaeozoic Europe in SE Poland – CELEBRATION 2000 experiment, profile CEL02. *Tectonophysics*, **401**, 55–77.
- Matthew, G.F. 1887. Illustrations of the fauna of the St. John Group, No. 4. On the smaller-eyed trilobites of Division I, with a few remarks on the species of higher divisions of the group. *Canadian Record of Science*, **2**, 357–363.
- Matthew, G.F. 1890. On Cambrian organisms in Acadia. *Transactions of the Royal Society of Canada*, **7**, 135–162.
- Matthew, G.F. 1895. The Protolenus fauna. *Transactions of the New York Academy of Sciences*, **14**, 101–153.
- Matthew, G.F. 1897. What is the Olenellus Fauna? *The American Geologist*, **19**, 396–407. Matthew, G.F. 1899. Studies on Cambrian faunas, no. 4. Fragments of the Cambrian Faunas of Newfoundland. *Transactions of the Royal Society of Canada, Series 2, Section 4*, **5** (for 1899–1900), 67–95.
- Mayoral, E., Liñán, E., Gámez-Vintaned, J.A., Gozalo, R. and Dies, M.E. 2008. El Cámbrico inferior del Cerro del Hierro (Sevilla). Propuesta de itinerario paleontológico. In: I. Bonilla and A. Menor Campillo (coord.), *Investigación científica y conservación en el Parque Natural Sierra Norte de Sevilla*. Consejería de Medio Ambiente – Junta de Andalucía, Sevilla, 27–43.
- Mizerski, W., Orłowski, S., Przybycin, A. and Skurek-Skurczyńska, K. 1999. Large-scale erosional channels in the Lower Cambrian sandstones, Gieraszwice environs (Kielce Block, Holy Cross Mts.). *Geological Quarterly*, **43**, 353–364.
- Mizerski, W., Orłowski, S. and Różycki, A. 1986. Tectonics of the Ocieski and Zamczysko ranges in the Holy Cross Mts. *Geological Quarterly*, **30**, 187–200.
- Mizerski, W., Orłowski, S. and Waksmundzki, B. 1991. New data on geology of the Kamieniec Shale Formation (Lower Cambrian, Holy Cross Mts.). *Geological Quarterly*, **35**, 149–162.
- Moberg, J.C. and Segerberg, C.O. 1906. Bidrag till Kännedomen om Ceratopyge-regionem med särskild hänsyn till dess utveckling i Fogelsångrakten. *Lunds Universitet Årsskrift N.F.*, **2**(2), no. 7, 1–113.
- Moczyłowska, M. 1991. Acritarch biostratigraphy of the

- Lower Cambrian and the Precambrian-Cambrian boundary in southeastern Poland. *Fossils and Strata*, **29**, 1–127.
- Moore, R.C. (Ed.) 1959. Treatise on Invertebrate Paleontology, Part O, Arthropoda 1. pp. i–xix, 1–560. Geological Society of America, Boulder, and University of Kansas Press, Lawrence.
- Morris, S.F. 1988. A review of British trilobites, including a synoptic revision of Salter's monograph. *Palaeontographical Society Monographs*, **574**, 1–316.
- Nawrocki, J. 2006. Palaeogeography of the Holy Cross Mountains in the early Palaeozoic. In: S. Skompski and A. Żylińska (Eds), *Procesy i zdarzenia w historii geologicznej Gór Świętokrzyskich*. 77 Zjazd Naukowy Polskiego Towarzystwa Geologicznego, 9–13. [In Polish]
- Nawrocki, J., Dunlap, J., Pecskay, Z., Krzemiński, L., Żylińska, A., Fanning, M., Kozłowski, W., Salwa, S., Szczepanik, Z. and Trela, W. 2007. Late Neoproterozoic to Early Palaeozoic palaeogeography of the Holy Cross Mountains (Central Europe): an integrated approach. *Journal of the Geological Society, London*, **164**, 405–423.
- Nawrocki, J. and Poprawa, P. 2006. Development of Trans-European Suture Zone in Poland: from Ediacaran rifting to Early Palaeozoic accretion. *Geological Quarterly*, **50**, 59–76.
- Nawrocki, J., Żylińska, A., Buła, Z., Grabowski, J., Krzywiac, P. and Poprawa, P. 2004. Early Cambrian location and affinities of the Brunovistulian terrane (Central Europe) in the light of palaeomagnetic data. *Journal of the Geological Society, London*, **161**, 513–522.
- Nielsen, A.T. and Schovsbo, N.H. 2011. The Lower Cambrian of Scandinavia: Depositional environment, sequence stratigraphy and palaeogeography. *Earth-Science Reviews*, **107**, 207–310.
- Orłowski, S. 1974. Lower Cambrian biostratigraphy in the Holy Cross Mts, based on the trilobite family Olenellidae. *Acta Geologica Polonica*, **24**, 1–16.
- Orłowski, S. 1975a. Lower Cambrian trilobites from Upper Silesia (Goczałkowice borehole). *Acta Geologica Polonica*, **25**, 377–383.
- Orłowski, S. 1975b. Cambrian and Upper Precambrian lithostratigraphic units in the Holy Cross Mts. *Acta Geologica Polonica*, **25**, 431–448. [In Polish with English summary]
- Orłowski, S. 1985a. Lower Cambrian and its trilobites in the Holy Cross Mts. *Acta Geologica Polonica*, **35**, 231–250.
- Orłowski, S. 1985b. New data on the Middle Cambrian trilobites and stratigraphy in the Holy Cross Mts. *Acta Geologica Polonica*, **35**, 251–263.
- Orłowski, S. 1985c. A trilobite with North American affinity in the Lower Cambrian of Poland. *Journal of Paleontology*, **59**, 975–978.
- Orłowski, S. 1987. Stratigraphy of the Lower Cambrian in the Holy Cross Mountains, Central Poland. *Bulletin of the Polish Academy of Sciences, Earth Sciences*, **35**, 91–96.
- Orłowski, S. 1988. Stratigraphy of the Cambrian System in the Holy Cross Mts. *Geological Quarterly*, **32**, 525–532.
- Orłowski, S. 1989. Trace fossils in the Lower Cambrian sequence in Świętokrzyskie Mountains, Central Poland. *Acta Palaeontologica Polonica*, **34**, 211–231.
- Orłowski, S. 1992. Cambrian stratigraphy and stage subdivision in the Holy Cross Mountains, Poland. *Geological Magazine*, **129**, 471–474.
- Orłowski, S. and Mizerski, W. 1995. New data on geology of the Middle Cambrian rocks in the Klimontów Anticlinorium (Holy Cross Mts.). *Geological Quarterly*, **39**, 293–306.
- Orłowski, S. and Żylińska, A. 2002. Lower Cambrian trace fossils from the Holy Cross Mountains, Poland. *Geological Quarterly*, **46**, 135–146.
- Pajchłowa, M. (Ed.) 1990. Atlas of guide and characteristic fossils, Geology of Poland, III, 1a, Older Palaeozoic (with Upper Proterozoic), pp. 1–493. [In Polish, English version published in 1991].
- Palmer, A.R. and Gatehouse, C.G. 1972. Early and Middle Cambrian trilobites from Antarctica. *Geological Survey of America Professional Papers*, **456-D**, 1–37.
- Palmer, A.R. and Repina L.N. 1993. Through a glass darkly: taxonomy, phylogeny, and biostratigraphy of the Olenellina. *The University of Kansas Paleontological Contributions*, **3**, 1–35.
- Palmer, A.R. and Rowell, A.J. 1995. Early Cambrian trilobites from the Shackleton Limestone of the Central Transantarctic Mountains. *The Paleontological Society Memoir*, **45**, 1–28.
- Radwański, A. 1969. Lower Tortonian transgression onto the southern slopes of the Holy Cross Mts. *Acta Geologica Polonica*, **19**, 1–164. [In Polish]
- Repina, L.N., Petrunina, Z.E. and Khajrullina, T.I. 1975. Trilobites. In: Stratigraphy and fauna of the Lower Palaeozoic of the northern submontane belt of Turkestan and Alai Ridges (southern Tyan-shan). Akademia Nauk SSSR, Sibirskoe Otdelenie, Trudy Instituta Geologii i Geofiziki, **278**, 100–248. [In Russian]
- Resser, C.E. and Howell, B.F. 1938. Lower Cambrian *Olenellus* Zone of the Appalachians. *Bulletin of the Geological Society of America*, **49**, 195–248.
- Richter, R. and Richter, E. 1941. Die Fauna des Unter-Kambriums von Cala in Andalusien. *Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft*, **455**, 1–90.
- Richter, R. and Richter, E. 1948. Studien im Paläozoikum der Mittelmeer-Länder, 8. Zur Frage des Unter-Kambriums in Nordost-Spanien. *Senckenbergiana*, **29**, 23–29.
- Rushton, A.W.A. 1966. The Cambrian trilobites from the Purley Shales of Warwickshire. *Palaeontological Society Monographs*, **120**, 1–55.
- Rushton, A.W.A., Brück, P.M., Molyneux, S.G., Williams, M. and Woodcock, N.H. 2011. A revised correlation of the

- Cambrian Rocks in the British Isles. *Geological Society Special Report*, **25**, 1–62.
- Samsonowicz, J. 1918. Materials on the geology of the Holy Cross Mountains. Discovery of the Lower Cambrian in the Holy Cross Mountains. *Sprawozdania z Posiedzeń Towarzystwa Naukowego Warszawskiego*, wydział III, **11** (5), 701–705. [In Polish]
- Samsonowicz, J. 1920. Stratigraphy of the Cambrian and Ordovician in the eastern part of the Holy Cross Mountains. *Sprawozdania Polskiego Instytutu Geologicznego*, **1**, 53–67. [In Polish]
- Samsonowicz, J. 1959a. On the Holmia-Fauna in the Cambrian of the Anticlinorium of Klimontów. *Bulletin de l'Académie Polonaise des Sciences, Série des sciences chimiques, géologiques et géographiques*, **7**, 447–452.
- Samsonowicz, J. 1959b. On *Strenuaeva* from the Lower Cambrian in the Klimontów Anticlinorium. *Bulletin de l'Académie Polonaise des Sciences, Série des sciences chimiques, géologiques et géographiques*, **7**, 521–524.
- Samsonowicz, J. 1959c. On *Strenuella* and *Germanopyge* from the Lower Cambrian in the Klimontów Anticlinorium. *Bulletin de l'Académie Polonaise des Sciences, Série des sciences chimiques, géologiques et géographiques*, **7**, 525–529.
- Samsonowicz, J. 1962. Lower Cambrian fossils from the Klimontów anticlinorium of the Holy Cross Mts., Poland. Prepared for print from the deceased author's MS notes by K. Korejwo and L. Teller. In: E. Passendorfer (Ed.), *Księga Pamiątkowa ku czci Profesora Jana Samsonowicza*, pp. 9–29.
- Schmidt, F. 1888. Ueber eine neuentdeckte untercambrische Fauna in Estland. *Mémoires de l'Académie Imperiale des Sciences de St-Petersbourg (7th series)*, **36**, 1–27.
- Sdzuy, K. 1961. Teil II: Trilobiten. In: F. Lotze and K. Sdzuy (Eds), *Das Kambrium Spaniens. Akademie der Wissenschaften und der Literatur, Abhandlungen der mathematisch-naturwissenschaftlichen Klasse*, **1961** (7, 8), pp. 217–408.
- Sdzuy, K. 1962. Trilobiten aus dem Unter-Kambrium der Sierra Morena (S-Spanien). *Senckenbergiana Lethaea*, **43**, 181–229.
- Sdzuy, K. 1995. Acerca del conocimiento actual del sistema Cámbrico y del Límite Cámbrico Inferior – Cámbrico Medio. In: J.A. Gámez-Vintaned and E. Liñán (Eds), *Memorias de las IV Jornadas Aragonesas de Paleontología: La expansión de la vida en el Cámbrico. Libro homenaje al Prof. Klaus Sdzuy. Institución Fernando el Católico, Zaragoza*, 253–263.
- Shaler, N.S. and Foerste, A.F. 1888. Preliminary description of North Attleboro fossils. *Harvard Museum of Comparative Zoology Bulletin*, **16**, 27–41.
- Shaw, A.B. 1950. A revision of several Early Cambrian trilobites from eastern Massachusetts. *Journal of Paleontology*, **24**, 577–590.
- Studencki, M. 1988. Sedimentary conditions of the Ocieski Sandstone and Kamieniec Shale formations (Lower Cambrian) in the Holy Cross Mountains. *Kwartalnik Geologiczny*, **32**, 533–540. [In Polish]
- Szczepanik, Z. and Żylińska, A. 2012. Dating of Cambrian rocks near Klimontów based on trilobites and acritarchs. In: W. Mizerski and B. Żbikowska (Eds), *Geologia jedna?! II Polski Kongres Geologiczny, Abstrakty*, 86. [In Polish]
- Vidal, G. 1981. Lower Cambrian acritarch stratigraphy in Scandinavia. *Geologiska Föreningens i Stockholm Förhandlingar*, **103**, 183–192.
- Vidal, G. and Nystuen, J.P. 1990. Lower Cambrian acritarchs and the Proterozoic-Cambrian boundary in southern Norway. *Norsk Geologisk Tidsskrift*, **70**, 191–222.
- Walcott, C.D. 1890. The fauna of the Lower Cambrian or Olenellus zone. In: 10<sup>th</sup> Annual Report of the Director, 1888–1889, United States Geological Survey, Part 1 – Geology, 509–774.
- Westrop, S.R. and Landing, E. 2000. Lower Cambrian (Branchian) trilobites and biostratigraphy of the Hanford Brook Formation, southern New Brunswick. *Journal of Paleontology*, **74**, 858–878.
- Zhang, W., Lu, Y., Zhu, Z., Qian, Y., Lin, H., Zhou, Z., Zhang, S. and Yuan, J. 1980. Cambrian trilobite faunas of southwestern China. *Palaeontologica Sinica, new series B*, **159**, 1–497. [In Chinese]
- Żylińska, A. and Kin, A. 2010. Defense strategy in *Strenuella polonica* Czarnocki (Ellipsocephalidae, Trilobita) from the Cambrian of the Holy Cross Mountains. In: M. Zatoń, W. Krawczyński, M. Salamon and A. Bodzioch (Eds), *Kopalne Biocenozy w czasie i przestrzeni, XXI Konferencja Naukowa Sekcji Paleontologicznej Polskiego Towarzystwa Geologicznego, Żarki-Letnisko*, 13–16 września 2010, 91–93. [In Polish]
- Żylińska, A., Kin, A. and Nowicki, J. *in press*. Application of morphometric techniques for taxonomic revision of *Berabichia oratrix* (Orłowski, 1985a) (Trilobita, Cambrian) from the Holy Cross Mountains, Poland. *Geodiversitas*.
- Żylińska, A. and Masiak, M. 2007. Cambrian trilobites from Brzechów, Holy Cross Mountains (Poland) and their significance in stratigraphic correlation and biogeographic reconstructions. *Geological Magazine*, **144**, 661–686.
- Żylińska, A. and Szczepanik, Z. 2009. Trilobite and acritarch assemblages from the Lower–Middle Cambrian boundary interval in the Holy Cross Mountains (Poland). *Acta Geologica Polonica*, **59**, 413–458.

*Manuscript submitted: 20<sup>th</sup> July 2012*

*Revised version accepted: 15<sup>th</sup> November 2012*

## APPENDIX

Distribution of trilobites (in alphabetical order) in particular exposures (those within the Kamieniec Formation are underlined) with their repository numbers [MWG Museum of the Faculty of Geology, University of Warsaw (ZI/29/ collection of Stanisław Orłowski; ZI/42/ collection of Jan Samsonowicz); MUZPIG Museum of the Polish Geological Institute (1.II; 110.II and 113.II collections of Jan Czarnocki; 8.II collection of Jan Samsonowicz); AK collection of Adrian Kin, Museum of the Geoscience Friends Association “Phacops”, Łódź; UJ 214P collection of Michał Stachacz, Institute of Geology, Jagiellonian University, Cracow. Abbreviations: ce – cephalae, cr – cranidia, pl – pleurae, th – thoraces, pyg – pygidia; lib – librigenae; hyp – hypostomata; com – complete specimens; ax – axial shields; fr – various fragments. Type specimens are in **bold**.

*Acanthomicacca klimontowi* Orłowski, 1985a – Dąb: cr [MWG ZI/29/1287, 1288, **1297**, 1298, 1300–1302, 1304–1306, 1309, 1310, 1313–1317, 1573; MWG ZI/42/031, 036–038, 042, 050, 054, 055, 064, 244, 245, 247, 249–251, 256]; Igrzyczna Hill: cr [MUZPIG 110.II.214]; Julianów: cr [MUZPIG 8.II.3]; Kucebrza: cr [MWG ZI/29/1292, 1296, 1311, 1312; MWG ZI/42/027, 028]; Ociesęki: cr [MWG ZI/42/234; MUZPIG 110.II.137, 142, 186]; Olbierzowice: cr [MUZPIG 8.II.7, 8]; Wola Wąkopna: cr [MWG ZI/29/1303, 1307, 1308]; Zbelutka: cr [MWG ZI/42/254; MUZPIG 1.II.8, 8a; MUZPIG 113.II.1A, 14]; unknown locality: cr [MUZPIG 110.II.176A].

*Atops granulatus* Orłowski, 1985c – Leśniakowa Dębina: fr [MUZPIG 110.II.154]; Sterczyna Hill: cr [MWG ZI/29/**3732**], pl [MWG ZI/29/3733].

*Berabichia oratrix* (Orłowski, 1985a) – Buczyna: com [MWG ZI/42/007], cr [MWG ZI/42/237]; Cisów: cr [AK 12/TA 1–9, 12–38], th [AK 12/TA 10, 11], ax [AK 12/TA 5-1]; Igrzyczna Hill: com [MWG ZI/29/**1340**, 1377, 1394, 1418, 1419], cr + th [MWG ZI/29/1395], cr [MWG ZI/29/1344–1347, 1351–1357, 1362–1368, 1398, 1421, 1431, 1435, 1440, 1441, 1447, 1454, 1455, 1458, 1459; MWG ZI/42/204, 209, 213, 233, 288]; Leśniakowa Dębina: com [MWG ZI/29/1434], cr [MUZPIG 1.II.23, 38, 42–44, 47, 57, 59, 63, 64, 68, 73, 80, 85–89; MUZPIG 110.II.29, 36, 37, 38, 56, 62, 68, 70–72, 77, 78, 82, 86, 93, 95–99, 102, 105, 108, 110, 115–123, 126, 130, 133, 135, 141–146, 150, 151, 155–159, 164, 182, 190, 219; MUZPIG 1716.II.2; MWG ZI/29/1103, 1105, 1263, 1341–1343, 1349, 1350, 1358–1360, 1378, 1380–1384, 1386, 1388, 1389, 1391–1393, 1396, 1397, 1399–1409, 1417, 1420, 1422–1425, 1429, 1430, 1442, 1443, 1449, 1452, 1456, 1461, 1463, 1464, 1476, 1478, 1479, 1489, 1493, 1514, 1569; MWG ZI/42/068, 071, 086, 100, 188, 205, 207, 210, 212, 214, 218, 220, 223, 229–232, 236, 239–242, 269, 270, 276, 280–282, 284, 286; UJ 214P/T20, 23, 25], th [MWG ZI/42/205, 235], cr + lib [MWG ZI/29/1477]; Malkowska Hill: com [MWG ZI/29/1410], cr [MUZPIG 8.II.30, 31; MWG

ZI/29/1368, 1372, 1373, 1451, 1457; MWG ZI/42/211, 216, 217, 267, 268, 271, 274, 277, 279], th [MWG ZI/42/287]; Nowa Huta: cr [MWG ZI/42/075]; Ociesęki: com [MWG ZI/29/1453; MUZPIG 1689.II.47], cr + th [MWG ZI/29/1439], cr [MUZPIG 110.II.166, 170–173, 183; MWG ZI/29/1260, 1361, 1371, 1385, 1413, 1415, 1416, 1427, 1432, 1433, 1444–1446, 1450, 1460, 1465, 1526, 1530; MWG ZI/42/021, 023, 070, 072, 077, 078, 081, 082, 084, 173, 178, 179, 182, 184–187, 190, 191, 193, 195–198, 200, 201, 224, 227, 228, 243, 272, 275, 278, 283, 285], th [AK 300/CD 20-1; MWG ZI/42/238]; Pranie: cr [MWG ZI/42/067, 069, 076, 085, 088, 089, 252, 264; MWG ZI/29/1376, 1436, 1437], lib [MWG ZI/42/083]; Rembów: cr [MWG ZI/29/1466, 1470]; Słowik: cr [MWG ZI/42/094; MWG ZI/29/1411, 1412]; Sterczyna Hill: com [MWG ZI/29/1438, 1471, 1472], cr + th [UJ 214P/T5, T6], cr + lib + th [AK 34/TA], cr [MWG ZI/29/1348, 1370, 1374, 1375, 1387, 1390, 1414, 1426, 1448, 1467, 1469, 1473, 1475, 1480–1483; UJ 214P/T11, T12], th [MWG ZI/29/1379, 1428, 1462, 1468, 1474]; Telegraf Hill: cr [MWG ZI/42/087]; Ujazdek: cr [MUZPIG 8.II.35; MWG ZI/42/074, 208]; Zbelutka: cr [MUZPIG 1.II.8B; MUZPIG 113.II.2, 4, 8]; unknown locality: cr [MUZPIG 110.II.174, 176A, 177, 180, 184, 189, 191–203, 205, 213, 229–231, 235], lib [MUZPIG 110.II.231].

*Holmia glabra* Orłowski, 1974 – Ociesęki: ce [MWG ZI/42/**153**], lib [MWG ZI/42/154].

*Holmia marginata* Orłowski, 1974 – Dąb: ce [MWG ZI/29/1052, 1075, 1309; MWG ZI/42/001–003, 008], hyp [MWG ZI/29/1068, 1076], lib [MWG ZI/42/014]; Igrzyczna Hill: ce + th [MWG ZI/29/3723], ce [MWG ZI/29/**0991**–0993, 0995–1002, 1004–1006, 1008–1023, 1025–1032, 1034–1036, 1038–1046, 1048, 1049, 1051, 1053, 1054, 1061, 1063–1065, 1070, 1073, 3722; MUZPIG 110.II.207, 214, 217], lib [MWG ZI/29/1070, 1074], hyp [MWG ZI/29/3725], th [MWG ZI/29/3728], pyg [MWG ZI/29/1062]; Leśniakowa Dębina: ce [MWG ZI/29/1007, 1059, 1066, 1067, 1071, 1077, MUZPIG 110.II.163], hyp [MWG ZI/29/1060, 1082], lib [MWG

ZI/29/1078], pyg [MWG ZI/29/3729, 3734], fr [MWG ZI/29/1079]; Malkowska Hill: ce [MWG ZI/29/1047]; Ociesęki: ce [MWG ZI/29/1050, 1055–1057, 1080; MWG ZI/42/004, 180], ce+lib [MUZPIG 110.II.221], hyp [MUZPIG 110.II.148], lib [MUZPIG 110.II.137, 144], pyg [MWG ZI/29/3730]; Sterczyna Hill: ce [MWG ZI/29/1024, 1058, 1081], hyp [MWG ZI/29/3724]; Stołowa Hill: ce [MWG ZI/29/1037]; Szczecno: ce [MWG ZI/29/1033]; Wola Wąkopna: ce [MWG ZI/29/0994, 1003, 1069], hyp [MWG ZI/29/1069], lib [MWG ZI/29/1072]; unknown locality: ce [MUZPIG 110.II.215, 220, 235], ce+lib [MUZPIG 110.II.160a, 169, 178, 229], lib [MUZPIG 110.II.234].

*Holmia* sp. – Buczyna: lib [MWG ZI/42/289]; Cisów: ce [MUZPIG 1.II.95]; Dąb: hyp [MWG ZI/42/005, 298], lib [MWG ZI/42/296]; Gieraszwice: ce [MWG ZI/42/292], hyp [MWG ZI/42/290], pl [MWG ZI/42/158, 294]; Malkowska Hill: lib [MWG ZI/42/155, 156, 159]; Kucebrza: lib [MWG ZI/42/160]; Ociesęki: ce [MWG ZI/42/197; MUZPIG 110.II.152], hyp [MWG ZI/42/202]; Pranie: lib [MWG ZI/42/006].

*Kjerulfia orcina* Orłowski, 1974 – Buczyna: ce [MWG ZI/42/016], lib [MWG ZI/42/010], pl [MWG ZI/42/015]; Dąb: th [MWG ZI/29/1153]; Igrzyczna Hill: ce [MWG ZI/29/1084, 1094, 1097, 1140, 1141, 1146, 1151, 1157], lib [MWG ZI/29/1061, 1095, 1142], hyp [MWG ZI/29/1180], pl [MWG ZI/29/1145, 1154, 1160, 1164, 1166, 1167]; Leśniakowa Dębina: ce [MWG ZI/29/1089, 1091–1093, 1107–1109, 1116, 1118, 1123, 1126, 1136, 1137, 1143, 1175, 1178; MWG ZI/42/025], lib [MWG ZI/29/1085, 1098, 1100, 1102, 1103, 1104, 1121, 1125, 1129, 1155, 1156, 1159, 1165, 1176], hyp [MWG ZI/29/1105, 1115, 1128, 1150, 1158], th [MWG ZI/29/1086, 1087, 1177], pl [MWG ZI/29/1117, 1148, 1149, 1161; MUZPIG 1.II.19, 20, 22–44]; Madeje: ce [MWG ZI/29/1106, 1119, 1120, 1130, 1134, 1138, 1152; MWG ZI/42/025], hyp [MWG ZI/29/1122], lib [MWG ZI/29/1146], pl [MWG ZI/42/026]; Malkowska Hill: lib [MWG ZI/42/157]; Ociesęki: ce [MWG ZI/29/1083, 1090, 1099, 1113, 1135; MWG ZI/42/021–023; MUZPIG 110.II.211], lib [MWG ZI/29/1112, 1114; MUZPIG 110.II.143], pl [MWG ZI/29/1110, 1111]; Sterczyna Hill: com [MUZPIG 1689.II.41], ce + th [MWG ZI/29/3727, 3735], ce [MWG ZI/29/1088, 1096, 1101, 1127, 1133, 1139, 1162, 1168, 1169, 1171, 1173, 1174, 1179, 1181–1184, 3731], lib [MWG ZI/29/1132, 1185], hyp [MWG ZI/29/3726], th [MWG ZI/29/1124, 1144, 1170], pl [MWG ZI/29/1163, 1172], fr [MWG ZI/29/1186]; Wola Wąkopna: th [MWG ZI/29/1187]; unknown locality: ce [MUZPIG 110.II.187].

*Postfallotaspis spinatus* Orłowski, 1985a – Igrzyczna Hill: ce [MWG ZI/29/2130].

*Schmidtellus nodosus* Orłowski, 1985a – Igrzyczna Hill: ce + th [MWG ZI/29/1322], ce [MWG ZI/29/1321, 1324, 1326, 1329], pl [MWG ZI/29/1323, 1335]; Leśniakowa Dębina: ce [MWG ZI/29/1328, 1330]; Malkowska Hill: ce [MWG ZI/29/1318, 1325], pyg? [MWG ZI/29/1336], pl [MWG ZI/29/1337]; Pranie: lib [MWG ZI/42/066]; Sterczyna Hill: ce + th [MWG ZI/29/1320], ce [MWG ZI/29/1319, 1327, 1331, 1332, 1334], hyp [MWG ZI/29/1333].

*Schmidtellus panowi* (Samsonowicz, 1959a) – Dąb: ce [MWG ZI/42/047]; Gieraszwice – pl [MWG ZI/42/033]; Leśniakowa Dębina: com [MWG ZI/29/1338], ce [MWG ZI/29/1339]; Ryj Hill: ce+ th [MUZPIG 1.II.94].

*Strenuella polonica* Czarnocki, 1926 – Igrzyczna Hill: cr + th [MWG ZI/29/1491, 1551, 1553], cr [MWG ZI/29/1495, 1515, 1568; MUZPIG 110.II.13, 14, 16], lib [MUZPIG 110.II.14], th [MWG ZI/29/1552]; Jaźwina Hill: th [MUZPIG 110.II.42], th + pyg [MUZPIG 110.II.52]; Leśniakowa Dębina: com [MWG ZI/42/073, MWG ZI/29/1494, 1567, 1576], ax [MWG ZI/29/1487, 1488, 1559], cr + lib + th [MWG ZI/29/1486, 1519; MUZPIG 110.II.181], cr + th [MWG ZI/29/1516, 1560–1562, 1596; MWG ZI/42/103], cr [MWG ZI/29/1093, 1341, 1401, 1407, 1408, 1484, 1485, 1489, 1490, 1493, 1496, 1514, 1517, 1569, 1571, 1577, 1587–1589, 1592, 1593, 1595, 1597, 1600, 1601, 1603, 1610–1614, 3739; MWG ZI/42/096, 100, 192, 206, 212, 219, 221–223, 225, 226, 229, 232, 235, 269, 282; MUZPIG 1.II.20, 30, 32, 36–38, 43–48, 52–61, 63, 64, 68–89; MUZPIG 110.II.29, 35, 37, 56, 70–72, 75, 77–79, 82, 84, 86, 93, 95–97, 99, 102, 105, 108, 110, 115, 116, 119–121, 123, 125–127, 133, 137, 139, 145, 150, 151, 164, 181, 182], th [MWG ZI/II/1494, 1518, 1604; MWG ZI/42/174], hyp [MUZPIG 1.II.50; MWG ZI/42/199], lib [MUZPIG 1.II.57, 62, 63, 65, pyg [MUZPIG 1.II.49, 53, 57, 66, 84; MUZPIG 110.II.110]; Madeje: cr [MWG ZI/29/1521]; Ociesęki: com [MWG ZI/42/090], ax [MWG ZI/29/1537], cr + lib [MWG ZI/29/1498], cr + lib + th [MWG ZI/29/1513, 1540], cr + th [MWG ZI/29/1502, 1506, 1510, 1534], cr [MWG ZI/29/1427, 1499, 1500, 1503–1505, 1507–1509, 1511, 1523, 1525, 1526, 1529–1531, 1533, 1535, 1536, 1538, 1541, 1554, 1556, 1570, 1572, 1590, 1591; MWG ZI/42/072, 079, 080, 095, 101, 173, 175–177, 183, 189, 194, 215, 273, 283; MUZPIG 110.II.35, 166], th [MWG ZI/29/1501, 1507, 1522, 1524, 1528], th + pyg [MWG ZI/29/1532], lib [MWG ZI/29/1512], pl [MWG ZI/29/1527], hyp [MWG

ZI/29/1557]; Pranie: cr [MWG ZI/29/1594]; Rembów: cr [MUZPIG 1.II.91]; Sterczyna Hill: com [MWG ZI/29/1546, 1547, 1586, 1599, 1607; AK 300/CD 33-5 (1)], ax [AK 300/CD 33-5 (3)], cr + lib + th [AK 45/TA1, 100/AB 32-3, 100/AB 32-16, 300/CD 31-8 (2); MWG ZI/29/1558, 1566, 1579, 1582, 1584, 1598], cr + th [MWG ZI/29/1497, 1545, 1550, 1606; AK 45/TA2 (3), 100/AB 24-26, 300/BC 31-12, 300/CD 33-5 (2), 300/CD 33-5 (5); UJ 214P/T4, T5, T13, T14, T16, T17], cr [MWG ZI/29/1492, 1520, 1539, 1543, 1549, 1555, 1563-1565, 1574, 1578, 1581, 1583, 1585, 1602, 1609; UJ 214P/T6, T11; MUZPIG 1689.II. 44, 51], th [MWG ZI/29/1544, 1548, 1575, 1580, 1582, 1608; AK 300/CD 33-5 (4); UJ 214P/T9, T10, T15], th + pyg [MWG ZI/29/1574; AK 300/CD 20-1; UJ 214P/T1, T2], lib [AK 100/AB 25-1; MWG ZI/29/1542]; unknown loca-

lity: cr [MUZPIG 110.II.174, 180, 189, 192-194, 196, 198, 202, 203, 205, 235], cr + th [MUZPIG 113.II.5], th [MUZPIG 113.II.5], pyg [MUZPIG 110.II.235].

*Strenuella zbelutkae* Orłowski, 1985a – Dąb: cr [MWG ZI/29/1290, 1291]; Leśniakowa Dębina: cr [MUZPIG 1.II.34; MUZPIG 110.II.179A]; Ociesęki: cr [MUZPIG 110.II.160B]; Zbelutka: cr [MWG ZI/29/1286, 1289; MWG ZI/42/029, 035, 040; MUZPIG 1.II.8b, 9a, 11; MUZPIG 113.II.1b-d, 2a-d, 4, 14, 18, 20, 23, 25, 26]; unknown locality: cr [MUZPIG 110.II.168A, B, C, 176B, C, 215, 233].

*Termierella sandomirensis* Samsonowicz, 1962 – Gieraszowice: cr [MWG ZI/42/030, 032, 034, 039, 041, 043, 045, 051, 053, 056].