

THEODOR NEAGU<sup>1</sup>

CENOMANIAN BENTHONIC FORAMINIFERA  
 IN THE SOUTHERN PART OF THE EASTERN CARPATHIANS  
 (ROMANIA)

(Pl. I—XV and 1 Fig.)

*Otwornice bentoniczne cenomanu południowej części  
 Wschodnich Karpat (Rumunia)*

(Tabl. I—XV i 1 fig.)

**Abstract:** This paper represents the second part of an article published in volume 39/1969 pp. 133—181 dealing with the Cenomanian foraminiferal fauna in the southern part of the Eastern Carpathians. A detailed microbiostratigraphy of these deposits has been established with use of planktonic foraminifera complemented by benthonic forms. A systematic presentation of the benthonic foraminiferal fauna follows a general discussion of the sequence of the planktonic foraminiferal zones which have been recognized.

GENERAL PART

Lithologically, the lowermost Upper Cretaceous (Cenomanian) deposits in the southern area of the Eastern Carpathians are rather uniform and are represented by: marls, marly-limestones, or marly-shales of various colours and of a more or less high degree of sorting of the constituents. These deposits, which as a rule, are transgressive upon the Lower Cretaceous formations begin with a sequence of grey-black or faintly blackish marls with a rich fauna of *Aucellina grypheoides* (Sowberby), accompanied by *Parahibolites tourtiaie* (Wein g). Then follows the Cenomanian proper, represented by a micaceous, reddish, cherry, or brick coloured or grey-black marls or marly-limestones, containing a rich fauna of Cephalopoda such as: *Neohibolites ultimus* (d'Or b.), *Mantelliceras mantelli* (Sowberby), *Acanthoceras rothomagense* (Defrance), *Turrilites* cf. *costatus* (Lamarc k), *Puzosia planulata* (Sowberby) etc., the pelecypod *Inoceramus crippsi* Mantel is also present. This macrofauna confirms with certainty from a stratigraphic viewpoint the presence of the whole Cenomanian stage. The foraminiferal fauna, which is very rich in planktonic, as well as benthonic species, thoroughly confirms in all respects this age determination. From a stratigraphic viewpoint the position of the basal part — also know as the "Aucellina marls" — is also interesting. According to the fauna and the geometrical position, Romanian geologist assign this part to the Vraconian, which they consider as a transitional substage between the Upper Albian and the Cenomanian stages. This transitional

<sup>1</sup> Labor. Paleontologie Univ. București, B-dul N. Bălcescu 1.

position is also reflected by the foraminiferal content, where there may be observed an obvious mixture of elements of two assemblages such as: *Nodosarella (Clarella) bulbosa* ten Dam, *Pleurostomella obtusa* Berthelin, *P. reussi* Berthelin, *Globorotalites brotzeni* Hofker, which are strongly developed in the Albian deposits of an epicontinental type (Moesian Platform), and *Rotalipora apenninica* (Renz), *Planulina lundegreni* Brotzen, *Heterolepa polyraphes* (Reuss), but which are rather rare and are well developed in the Cenomanian. Among the benthonic foraminifera, special mention should be made of *Tritaxia gaultina* (Morozova) which appears here and constitutes one of the characteristic components of the Cenomanian foraminiferal fauna, ranging into the Lower Turonian as well. *Ammobaculites problematicus* Neagu exhibits a similar distribution. These two species disappear at the same time, but, whereas *Tritaxia gaultina* exhibits several morphotypes, some of which are very different from the type, *Ammobaculites problematicus* remains rather constant in character.

Thus, the conclusion may be drawn that micropaleontologically the series of Aucellina beds represents a moment of transition in which old elements which are on the point of disappearing, coexist with the new ones whose evolution is beginning. Microbiostratigraphically, we have separated this moment under the name of the *Hedbergella delrioensis* Zone.

The Lower Cenomanian is represented by a foraminiferal assemblage containing both, benthonic and planktonic species in abundance. Within the planktonic assemblage there is a strong development of the genus *Rotalipora*, and this continues progressively up to the latest part of the Cenomanian, even passing into the basal part of the Lower Turonian. This growth is so strong that within the Middle, but particularly within the Upper Cenomanian, the benthonic fauna of foraminifera is almost entirely concealed. When following the development of the foraminiferal assemblage through the entire Cenomanian stage, the interdependence existing between the degree of development of the benthonic and of the planktonic foraminifera becomes very obvious. The inverse relation of proportionality of these two groups of foraminifera is clearly demonstrated. The remarkable degree of development of the planktonic foraminifera has enabled the writer to subdivide the Cenomanian stage into a sequence of zones, subzones and even zonules. The abundance of species and specimens in the Cenomanian faunas indicate that they do not correspond in any way the assemblages of the flysch type which Carpathian micropaleontologists are used to find. With minor exceptions, these assemblages occur also with the same or greater degree of development within the deposits of an epicontinental type in the Moesian Platform and Dobrogea. It seems that during the Cenomanian the geosynclinal flysch basin at least within this region (the southeastern Carpathians) were under the influence of an epicontinental sea localized more southwards in the area of the Moesian Platform. This opinion is also confirmed by the fact that in some parts of this basin, where the epicontinental influence is not in evidence in the basal part of the series, the foraminiferal fauna is composed exclusively or dominantly of arenaceous form (flysch character). When this influence becomes perceptible — although the planktonic fauna does not make up the basal part as in the other sectors — it nevertheless causes an obvious change which is also discernible in the benthonic fauna. In this sense, the outcrop from Dealul Stîniş (Int. Buzăului) is significant.

Likewise, in the case of the Cenomanian deposits in the southern area

of the Eastern Carpathians, the most distinctive element from an evolutionary viewpoint is the planktonic fauna. The benthonic fauna exhibits a much slower evolution, individual ranges of some species cross one or more stage boundaries. Because of the considerations, a detailed microbiostratigraphy of these formations can be achieved only by carefully determinations the evolution of the planktonic foraminifera. This opinion is supported by the ranges shown in the table (Fig. 1).

Considering the problem from a microbiostratigraphic viewpoint, the Cenomanian deposits in the southern part of the Eastern Carpathians — although structurally belonging to different units, i.e.: the Ceahlău Nappe and Bobu Unit, the Dîmbovicioara Couloir and the Curbicortical Flysch or Teleajen Flysch — may be separated into the following subdivisions:

*Hedbergella* Zone. Stratigraphically, this zone corresponds to the Vraconian and is characterized, as has been shown, from a micropaleontological viewpoint, by the mixture of the two faunal types (Upper Albian and Lower Cenomanian). Within the populations of this Zone, however, the dominant element is the genus *Hedbergella* with one or several species (*H. delrioensis* (Carsey) *H. planispira* (Tappan), *H. trocoidea* (Gandolfi) with which are associated the first occurrence of *Tritaxia gaultina gaultina* (Morozova). This species continues its evolution parallel to the genus *Rotalipora* with a series of morphotypes. The remaining microfauna in this zone, which is rather poor, is composed of: *Reophax minuta* Tappan, *R. deckeri* Tappan, *Kalamopsis grzybowskii* (Dylańska), *Dorothia pupa* (Reuss), *D. oxycona* (Reuss), *Textularia foeda* Reuss, which in some samples — Prahova Valley — become the dominant element, *Globorotalites brotzeni* Hofker, *Nodosarella (Clarella) bulbosa* ten Dam, *Heterolepa polyraphes* (Reuss), *Praeglobotruncana delrioensis* (Plummer) (abundant), *P. stephani stephani* (Gandolfi) (rare), *Rotalipora apenninica* (Renzi) (rare).

Corresponding to this zone in the outcrop of Stîniş Hill-Int. Buzăului, where the *Aucellina* beds are lacking, there develops a series of grey-greenish and cherry-coloured clays in which occur solely or almost alone agglutinated foraminifera such as: *Rhizammina* sp., *Hyperammina gaultina* ten Dam, *Saccammina sphaerica* Sars, *Hormosina ovulum* (Grzybowski), *Reophax minuta* Tappan, *Glomospira charoides* (Jones & Parker), *G. irregularis* (Grzybowski), *Ammodiscus cretaceus* (Reuss), *A. infimus* Franke, *Kalamopsis grzybowskii* (Dylańska), *Plectrorecurvoides alternans* North (as a prevailing element), *Recurvoides imperfectus* Hanzlíková, *Haplophragmoides kirki* Wickenden, *H. gigas minor* Nauss, *Ammobaculites problematicus* Neagu *Trochammina umiatensis* Tappan, *Plectina lenis* (Grzybowski). The fact that these deposits are overlain by a sequence of sediments in which the planktonic forms from below reappear—*Hedbergella planispira* (Tappan), *Praeglobotruncana stephani stephani* (Gandolfi), and then *Rotalipora* and *Schackoina* — as well as the presence of some agglutinated forms which occur also in the *Aucellina* beds, enables us to correlate these assemblages and assign them to the same age i.e., the Vraconian.

*Rotalipora* Zone. This zone corresponds paleontologically to the time when the genus *Rotalipora* constituted a main component of the planktonic foraminiferal fauna. Stratigraphically this zone corresponds almost exactly to the interval assigned to the Cenomanian, and may be subdivided into the following subzones:

— The *Rotalipora apenninica* Subzone;

— The *Rotalipora brotzeni* — *R. globotruncanoides* Subzone;

— The *Rotalipora cushmani* — *R. reicheli* Subzone.

Within the first subzone, the planktonic population is dominated by the *Rotalipora apenninica* group with its various subspecies or morphological types, to which can be added *Rotalipora ticinensis* (Gandolfi) and *R. techamaensis* Marianos & Zingula. Stratigraphically, this subzone begins somewhat below the Vraconian/Lower Cenomanian boundary, so that it also includes the latest part of the Vraconian, and continues into the basal part of the Middle Cenomanian. If one follows the evolution of the *Rotalipora apenninica* group, at the time when the planktonic population is very well developed, one may distinguish two zonules: i.e.; — in the lower part, the *R. apenninica apenninica* Zone, followed in the upper part by the *R. apenninica gandolfi* Zone.

Within the latter zonule the first representatives of the *Rotalipora brotzeni* and *R. globotruncanoides* groups appear, but, of course, they are not common. In the Dealul Stînii outcrop, where, as mentioned before, the epicontinental influences are less pronounced, the *Rotalipora* Zone has a much smaller extent. The lower part of the section, corresponding, as mentioned above, to the *Hedbergella* Zone as well as the entire Lower Cenomanian in addition to the lower part of the Middle Cenomanian, constitutes the *Plectrorecurvoides alternans* Zone. In its lower part, corresponding to the *Hedbergella* Subzone, we have separated the *P. alternans* Subzone in which, as ready mentioned above, the arenaceous foraminifera prevail, while in its upper part, corresponding to the *Rotalipora apenninica* Subzone, we have separated the *Hedbergella planispira* Subzone (this being the first strongly developed planktonic species after the arenaceous phase). Here also *Tritaxia gaultina* (Mrozova) occurs for the first time still in the lower subzone, corresponding to the situation in all other sections. In the composition of the benthonic fauna, the *H. planispira* Subzone is very similar and almost identical with the *R. apenninica apenninica* Subzone, this being also the decisive criterion which permitted the correlation.

The *Rotalipora brotzeni* — *R. globotruncanoides* Subzone follows above, but here the planktonic population is dominated by two species. The boundaries between these two subzones as well as between the other ones, are not of an absolute value. In establishing them we have taken into account the whole population, and not merely the first occurrence of a species or of a group of species.

As a connecting element between these two subzones, and to demonstrate the above statement, the *Rotalipora apenninica evoluta* Zone occurs; it shows an appreciable development in the upper part of the Lower Cenomanian, and in the lower part of the Middle Cenomanian. In the Dealul Stînii outcrop, the genus Schackoina with its various Cenomanian species and subspecies occurs within this subzone.

The *Rotalipora cushmani* — *R. reicheli* Subzone corresponds to the Upper Cenomanian but also embraces the basal part of the Lower Turonian. During this interval, the prevailing elements within the planktonic population are *Rotalipora reicheli* [*R. micheli* (Sacal & Debourle), *R. deecke* (Frank), *R. reicheli* Mornod] group together with a strong development of the „*R. cushmani* group” [*R. cushmani cushmani* (Morrow), *R. cushmani expansa* (Carbonier), *R. cushmani minor* (Mornod), *R. cushmani montsalvensis* (Mornod), *R. cushmani turonica* (Brotzen), *R. cushmani thomei* Hagn & Zeill.]

ALBIA NAPP LIPP	C E N O M A N I A N			LOWER TUO- NIAN	LITHOLOGY	MICROPA- LEONTOLO- GICAL S.A.M - PLES	STRUC- TURAL UNIT	
	VRACONIAN	LOWER	MIDDLE					UPPER
	GREY SILTY MARLS OR DARK GREY SANDY MARLS WITH: AUCELLINA GRYPHEOIDES, PARA- HIB TOURTIAE, LECH- ITES GAUDINI, ETC.	VARIEGATED MARLS; GREY, GREENISH, BLACK AND RED TO VIOLET, WITH NEOHIBOLITES ULTIMUS, MANTELLICERAS SP, PUZOSIA PLANULATA, INOCERAMUS.				THE DÎMBOVITZA VALLEY; THE JALO- MITZA VALLEY; THE LEURDEI VALLEY; THE PRAHOVA VALLEY	THE CEHLIXU BOBU SUBUNIT + THE	
	HEDBERGELLA	ROTALIPORA S ST TRITAXIA GAULTINA		BIG GLOBIGERINAE <sup>7</sup> AND PRAEGLOBOTRUNCANA DIV SP	ASSEMBLAGE ZONES	THE RUCĂR-PODUL DÎMBOVITZA BASIN	DÎMBOVICIOARA ZONE (DÎMBOVI- CIOARA COULOIR)	
	HEDBERGELLA DELRIODENSIS	ROTA LIPORA APENNINICA PRAEGLOBOTR DELRIOENSIS PRAEGL. STEPHANI	R. BROTZENI - GLOBOTRUNCANOIDES PRAEGL STEPHANI - STEPHANI		R. REICHEL - R CUSHMANI PRAEGL STEPHANI GIBBA			SUB- ZO- NES
		ROTA LIPORA APENNINICA APENNINICA GANDOLFII	R. BROTZENI R. GLOBOTRUNCANOIDES R. APENNINICA EVOLUTA		R. MICHELI R. DEECKEI R. REICHEL ROTALIPORA CUSHMANI TURONICA			ZO- NU- LES
	CONGLOMERATES SANDSTO- NES MARLY SANDSTONES WITH: AUCELLINA GRYPHEO- IDES, PARAHIB TOURTIAE, LECHITES GAUDINI, SCHLOENBACHIA, ETC.	GREY THIN BEDDED MARLS, CALCAREOUS MARLS AND MARLY SILTSTONES (RUCĂR MARLS) WITH: MANTELLICERAS MANTELLI, SCHLOENBACHIA VARIANS, SCAPHITES AEOUALIS, PUZOSIA PLA- NULATA, CALLICOCERAS SP., INOCERAMUS.			LITHOLOGY			
		ROTALIPORA S ST TRITAXIA GAULTINA			ASSEMBLAGE ZONES			
		ROTA LIPORA APENNINICA	R. BROTZENI - GLOBO- TRUNCANOIDES	R. REICHEL - CUSHMANI	SUB- ZO- NES			
		R. APENNINICA APENNINICA R. TICINENSIS	ROTA LIPORA APENNINICA GANDOLFII R. BROTZENI R. GLOBOTRUNCANOIDES	R. DEECKEI R. REICHEL ROTALIPORA CUSHMANI CUSHMANI TURONICA	ZO- ZONULES			
	„AUCELLINA BEDS” DARK GREY SANDY MARLS WITH A. GRYPHEO- IDES, PARAHIBOLITES TOURTIAE, ETC.	VARIEGATED MARLS: GREENISH, BLACK AND RED TO VIOLET, WITH: NEOHIBOLITES ULTIMUS, OR BLACK SHALES AND GREY MARLY - LIMESTONES WITH PUZOSIA PLANULATA, MANTELLICE- RAS MANTELLI, TURRILITES COSTATUS, INOCERAMUS			TUFFACEOUS RED AND GREY CLAYS	THE TELIU VALLEY	THE „CURBICORTICAL” <sup>9</sup> FLYSCH NAPPE (THE TELEA- JEN NAPPE)	
	?	ROTALIPORA S. ST TRITAXIA GAULTINA			ASSEMBLAGE ZONES			
	?	R. APENNINICA APENNINICA PRAEGLOBOTRUNCANA	R. GLOBOTRUNCANOIDES STEPHANI - STEPHANI	R. REICHEL R. CUSHMANI PRAEGL STEPHANI GIBBA RADIOLARIA BIOFACIES	P. SEUDOBO- LIVINA VARIABLES			
	RED AND GREEN CLAYS	RED MARLS, GREEN AND RED, MARLY LIMESTONES		BLACK SHALES AND GREY MAR- LY LIMESTONES WITH SCALES OF FISH	LITHOLOGY	INTORSURA BUZĂULUI (STĂNIU HILL) VAMA BU- ZĂULUI (THE BOULUI CREEK PRAEȘCU HILL)		
		PLECTORECURVOIDES ALTERNANS TRITAXIA GAULTINA	ROTALIPORA S ST		ASSEMBLAGE ZONES			
	PLECTORECURVOIDES ALTERNANS	HEDBERGELLA PLANISPIRA	R. BROTZENI SCHACKOINA	R. GLOBOTRUNCANOIDES CENOMANA	R. CUSHMANI TURONICA	SUBZO- NES		

From the terminal part of the Middle Cenomanian up to the lower half of the Upper Cenomanian in the group of the "high-spined *Rotaliporae*" (*R. reicheli*) the following forms develop: *Rotalipora micheli* and *R. deecke* which allows the separation of a lower zonule, while in the upper part of the Upper Cenomanian only *Rotalipora reicheli* remains, and it disappears almost abruptly at the boundary with the Lower Turonian.

Using the group of the "flat-spined *Rotaliporae*" (*R. cushmani*) which appear in the middle part of the Middle Cenomanian, we can recognize two zonules. Of these the *Rotalipora cushmani cushmani* Z o n u l e extends from the middle part of the Middle Cenomanian to the middle part of the Upper Cenomanian. The second one, the *R. cushmani turonica* Z o n u l e follows above and reaches the Upper Cenomanian/Lower Turonian boundary. Within this latter zonule, the following species are well developed: *Rotalipora cushmani thomei*, *R. cushmani minor*, *R. cushmani montsalvensis*, all of which accompany *Rotalipora cushmani turonica* until its disappearance.

In the Dealul Stînişii outcrop this zonule is lacking and probably corresponds to the series of grey-blackish marly-limestones and marls with fish remains but without microfauna. It is found again however, further southwards in the Prădescu Hill (Vama Buzăului area), but only with *R. cushmani turonica* occurring in a series of grey-blackish marls.

In the planktonic fauna, along with the *Rotalipora* group, within the sequence of Cenomanian deposits in the southern area of the Eastern Carpathians *Praeglobotruncana*, *Clavhedbergella* and *Hedbergella* genera are also found. Of particular interest among these genera is the genus *Praeglobotruncana* which does not show such a spectacular evolution permitting to establish subzones or zonules. Nevertheless, it can be seen that a series of Lower Cenomanian deposits is dominated by *Praeglobotruncana stephani stephani* (G a n d o l f i), which later in the Middle Cenomanian is joined by *P. marginaculeata* (L o e b l i c h & T a p p a n), and towards its terminal part also by the first bicarinate species — *P. algeriana* C a r o n. All species of this genus cross the Cenomanian/Turonian boundary; within the Lower Turonian this genus has a strong evolutionary outburst resulting in a great number of new species, some with a trend towards enormity in size and with reduced keels, others showing an ever increasing trend toward the development of two peripheral keels.

Whereas the planktonic foraminifera allow a microbiostratigraphical zonation, which in some cases is quite detailed, such a zonation cannot be established however, on the basis of benthonic foraminifera because of the great uniformity and extremely slow evolution, so that the rapidly evolving species occur but very rarely. The main components of this population include arenaceous foraminifera, among which *Haplophragmoides gigas minor* N a u s s, *Ammobaculites problematicus* N e a g u, *Spiroplectamina roemeri* L a l i c k e r, *S. gandolfii* C a r b o n i e r, *S. complanata* (R e u s s), *Tritaxia gaultina gaultina* (M o r o z o v a), *T. gaultina intermedia* (N e a g u), *T. gaultina carinata* (N e a g u), *T. gaultina discjuncta* (C u s h m a n), *Dorothia oxycona* (R e u s s), *D. pupa* (R e u s s), *D. concinna* (R e u s s), occur from the basal to the upper part of the Cenomanian. Among the calcareous species, the Nodosariidae are generally abundant together with various species of *Nodosaria*, *Dentalina*, *Lenticulina*, *Marginalina* and *Vaginulina*. Among the *Buliminaceae*, a special mention should be made of *Pyramidina minima robusta* N e a g u. s.ssp. which is placed in the Middle and Upper Cenomanian. Another feature of the benthonic

fauna to be emphasized is the particularly great development of the Pleurostomellidae. This development is rarely observed below or above these substages. In addition to the fact that the genera *Pleurostomella*, *Ellipsoidea*, *Nodosarella* (*Clarella*) and *Ellipsoglandulina* are represented by one to three species the specimens are in some cases of gigantic size, and their frequency is appreciable.

Among the *Rotaliidae*, an appreciable and sometimes very great abundance is shown by the Osangulariidae which in most cases are represented by very well developed Albian species such as *Osangularia cretacea* (Carbonier), *Gyroidinoides nitidus* (Reuss), *Globorotalites brotzeni* Hofker. They are accompanied by several cenomanian species such as: *Gyroidinoides mauretanicus* (Carbonier), *Globorotalites multiseptus* (Brotzen).

The same situation is shown by the Anomalinidae where the genera *Heterolepa*, *Gavelinella* and *Lingulogavelinella* are represented by several species which are characteristic for the Cenomanian such as: *Heterolepa polyraphes* (Reuss), *Gavelinella schoembachi* (Reuss), *Lingulogavelinella globosa* (Brotzen), and *Orostella aumalensis* (Sigal).

#### PALEONTOLOGICAL PART

##### *Hypocrepina depressa* Vašiček

Pl. I, fig. 4

*Hypocrepina depressa* Vašiček 1947, p. 243, pl. 1, fig. 1—2; Huss 1957, pl. 1, fig. 2 (3); Geroch 1959, pl. 12, fig. 15; 1960, pl. 6, fig. 1; 1966, p. 435, fig. 6 (8—13).

Occurrence: Vraconian-Cenomanian, D. Stînii Hill, Teliu Valley, Boului Greek Prahova Valley, Leurdei Valley.

Dimensions: length 0,96 mm; breadth 0,24 mm.

Hypotypes: L.P.B. 9206.

##### *Hyperammina gaultina* ten Dam

Pl. II, fig. 2

*Hyperammina gaultina* ten Dam 1950, p. 5, pl. 1, fig. 2; Huss 1957, pl. 1, fig. 2 (2); Geroch 1966, pl. 435, fig. 6 (14—18).

*Hyperammina elongata* Brady; Neagu 1962, p. 55, pl. 1, fig. 5.

Occurrence: Vraconian-Lower/Cenomanian, D. Stînii Hill, Boului Creek, Teliu Valley.

Dimensions: length 0,39 mm.

Hypotypes: L.P.B. 9321

##### *Psammosphaera fusca* Schulze

Pl. II, fig. 1

*Psammosphaera fusca* Schulze; Grzybowski 1896, p. 270, pl. 8, fig. 14; Cushman 1910, p. 35, text-fig. 25—28; 1918, p. 34, pl. 13, fig. 1—6; Franke 1928, p. 8, pl. 1, fig. 3; Neagu 1962, p. 53, pl. 1, fig. 3; Grün & all. 1964, p. 247, pl. 3, fig. 3; Huss 1966, p. 15, pl. 1, fig. 1—3.

Occurrence: Cenomanian, Teliu Valley, D. Stînii Hill, Boului Creek, Prahova Valley, Leurdei Valley.

Dimensions: diameter 0,39 mm.

Hypotypes: L.P.B. 9207.

*Ammodiscus cretaceus* (Reuss)

Pl. II, fig. 12

*Operculina cretacea* Reuss 1845, p. 35, pl. 13, fig. 64—65.

*Cornuspira cretacea* Reuss 1860, p. 177, pl. 1, fig. 1; 1862, p. 43, pl. 1, fig. 10, (not 11—12); Egger 1899, p. 18, pl. 22, fig. 1—2; Cushman 1926, p. 608, pl. 21 fig. 3; Franke 1928, p. 16, pl. 1, fig. 22; White 1928, p. 188, pl. 27, fig. 9.

*Ammodiscus cretaceus* (Reuss), Marie 1941, p. 18, pl. 1, fig. 5—6; Cushman 1946, p. 17, pl. 1, fig. 35; Hagn 1953, p. 4, pl. 1, fig. 3; Frizzell 1954, p. 58, pl. 1, fig. 15; Said & Kenawy 1956, p. 120, pl. 1, fig. 4; Belford 1960, p. 22, pl. 6 fig. 1; Tappan 1962, p. 130, pl. 30, fig. 1—2; Graham & Church 1963, pl. 1, fig. 17; Huss, 1966, p. 16, pl. 2, fig. 13—16.

Occurrence: Vraconian-Cenomanian, Teliu Valley, D. Stîinii, Boului Creek Prahova Valley, Leurdei Valley, Rucăr basin.

Dimensions: diameter 0,36 mm.

Hypotypes: L.P.B. 9208.

*Ammodiscus infimus* Franke

Pl. II, fig. 19

*Ammodiscus infimus* Franke; Grün & all. 1964, p. 258, pl. 4, fig. 11; Geroch 1966, p. 437, fig. 8 (13—14).

Occurrence: Vraconian-Cenomanian, Teliu Valley, D. Stîinii Hill, Boului Creek, Prahova Valley, Leurdei Valley, Ialomitza Valley, Rucăr basin.

Dimensions: diameter 0,50—1,20 mm.

Hypotypes: L.P.B. 9209/1—3.

*Glomospira charoides* (Jones & Parker)

Pl. II, fig. 6—10

*Ammodiscus charoides* Jones & Parker; Grzybowski 1896, pl. 8, fig. 39—43.

*Glomospira charoides* (Jones & Parker); Neagu 1962, p. 57, pl. 4, fig. 49—51; Grün & all. 1964, p. 260, pl. 5, fig. 14.

Occurrence: Vraconian-Cenomanian, Teliu Valley, D. Stîinii Hill, Boului Creek, Prahova Valley, Leurdei Valley, Rucăr basin.

Dimensions: diameter 0,26 mm.

Hypotypes: L.P.B. 9210.

*Glomospira gordialis* (Jones & Parker)

Pl. II, fig. 14—15

*Ammodiscus gordialis* Jones & Parker; Grzybowski, 1896, p. 281, pl. 8, fig. 44, 45.

*Glomospira gordialis* (Jones & Parker); Tollmann 1960, p. 149, pl. 6, fig. 5; Grün & all. 1964, p. 261, pl. 5, fig. 2.

Occurrence: Vraconian-Cenomanian, Teliu Valley, D. Stîinii Hill, Boului Creek, Prahova Valley, Leurdei Valley, Ialomitza Valley, Rucăr basin.

Dimensions: diameter 0,29 mm — 0,43 mm.

Hypotypes: L.P.B. 9211/1—2.

*Glomospira irregularis* (Grzybowski)

Pl. II, fig. 11

*Ammodiscus irregularis* Grzybowski 1898, p. 285, pl. 11, fig. 2—3; 1901, p. 27.



*Glomospira irregularis* (Grzybowski); Glaessner 1937, p. 359, pl. 1, fig. 7;  
Neagu 1962, p. 57, pl. 4, fig. 54, pl. 6, fig. 85—86; Grün & all. 1964, p. 263, pl. 4,  
fig. 4.

Occurrence: Vraconian-Cenomanian, Teliu Valley, D. Stîniş Hill, Boului  
Creek, Prahova Valley, Leurdei Valley, Ialomitza Valley.

Dimension: length 0,67.

Hypotypes: L.P.B. 9212.

*Lituotuba incerta* Franke

Pl. II, fig. 16—18

*Lituotuba incerta* Franke 1928, p. 15, pl. 1, fig. 11; Neagu 1962, p. 58, pl. 4,  
fig. 62—63.

Occurrence: Vraconian-Cenomanian, D. Stîniş Hill, Boului Creek, Pra-  
hova Valley.

Dimensions: length 0,46—0,70 mm.

Hypotypes: L.P.B. 9213/1—2.

*Kalamopsis grzybowskii* (Dylażanka)

Pl. I, fig. 1—3

*Hyperammina grzybowskii* (Dylażanka); Geroch 1960, p. 39, pl. 1, fig. 22,  
23, pl. 10, fig. 7.

*Kalamopsis grzybowskii* (Dylażanka); Grün & all. 1964, p. 254, pl. 3, fig. 14,  
Geroch 1966, p. 438, fig. 6 (27—29).

Occurrence: Vraconian-Cenomanian, Teliu Valley, D. Stîniş Hill,  
Boului Creek, Prahova Valley, Leurdei Valley, Rucăr basin.

Dimensions: length: 1,12—1,72 mm; thickness 0,14—0,17 mm.

Hypotypes: L.P.B. 9214/1—3.

*Hormosina ovulum* (Grzybowski)

Pl. II, fig. 5

*Reophax ovulum* Grzybowski 1896, p. 267, pl. 8, fig. 19—21; 1901, p. 268, pl. 7,  
fig. 3.

*Hormosina ovulum* (Grzybowski); Glaessner 1937, p. 357, pl. 1, fig. 5;  
Neagu 1962, p. 58, pl. 6, fig. 92; Geroch 1959, p. 116, pl. 13; Grün & all.  
1964, p. 254, pl. 5, fig. 9.

Occurrence: Vraconian-Cenomanian, Teliu Valley, D. Stîniş Hill,  
Boului Creek, Prahova Valley, Leurdei Valley, Ialomitza Valley, Rucăr  
basin.

Dimensions: diameter 0,40 mm; length 0,58 mm.

Hypotypes: L.P.B. 9215.

*Reophax deckeri* Tappan

Pl. II, fig. 3—4

*Reophax deckeri* Tappan 1940, p. 94, pl. 14, fig. 3; 1943, p. 479, pl. 77, fig. 3;  
Frizzell 1954, p. 57, pl. 1, fig. 7.

Occurrence: Vraconian-Cenomanian, D. Stîniş Hill, Prahova Valley.

Dimensions: length 0,86—1,00 mm; thickness 0,36—0,40 mm.

Hypotypes: L.P.B. 9216/1—2.

*Reophax minuta* Tappan

Pl. II, fig. 13

*Reophax minuta* Tappan 1940, p. 94, pl. 14, fig. 4; 1943, pl. 77, fig. 4; Frizzell 1954, p. 57, pl. 1, fig. 11; Tappan 1962, p. 132, pl. 30, fig. 10; Geroch 1960, p. 41, pl. 6, fig. 3; 1966, p. 439, fig. 7 (7—17).

Occurrence: Vraconian-Cenomanian, Teliu Valley, D. Stînii Hill, Boului Creek Prahova Valley, Rucăr basin.

Dimensions: length 0,60 mm; thickness 0,14 mm.

Hypotypes: L.P.B. 9217.

*Haplophragmoides gigas minor* Nauss

Pl. V, fig. 8—11

*Haplophragmoides gigas minor* Nauss 1947, p. 338, pl. 49, fig. 10; Geroch 1959, p. 117, pl. 12, fig. 19, 1966, p. 441, pl. 10, fig. 1—3.

Occurrence: Vraconian-Cenomanian, Teliu Valley, D. Stînii Hill, Boului Creek, Prahova Valley, Belia Valley, Leurdei Valley, Iaşomîţa Valley, Rucăr basin.

Dimensions: diameter 0,39—0,55 mm.

Hypotypes: L.P.B. 9218/1—2.

*Haplophragmoides bulloides* (Beissel)

Pl. II, fig. 20—21

*Haplophragmium bulloides* Beissel 1891, p. 17, pl. 4, fig. 24—30.

*Haplophragmoides bulloides* (Beissel), Huss 1966, p. 23, pl. 3, fig. 17—24.

Occurrence: Vraconian-Cenomanian, D. Stînii Hill, Prahova Valley, Rucăr basin.

Dimensions: diameter 0,36 mm.

Hypotypes: L.P.B. 9219.

*Haplophragmoides kirki* Wickenden

Pl. IV, fig. 14—19

*Haplophragmoides kirki* Wickenden; Cushman 1946, p. 21, pl. 2, fig. 23; Neagu 1959, pl. 4, fig. 11—16; Huss 1966, p. 27, pl. 4, fig. 17—24; Sliter 1968, p. 44, pl. 2, fig. 2.

Occurrence: Vraconian-Cenomanian, Teliu Valley, D. Stînii Hill, Boului Creek, Prahova Valley, Leurdei Valley, Rucăr basin.

Dimensions: diameter 0,34—0,48 mm.

Hypotypes: L.P.B. 9220/1—3.

*Haplophragmoides* sp. aff. *H. nonioninoides* Geroch (not Reuss)

Pl. I, fig. 19—21

*Haplophragmoides* cf. *chapmani* Crespin; Geroch 1959, p. 117, pl. 12, fig. 17—18.

*Haplophragmoides* aff. *nonioninoides* Geroch (not Reuss), 1966, p. 440 fig. 9 (1—19), fig. 11 (1 a—e).

Occurrence: Vraconian, Boului Creek (Vama Buzăului area).

Dimensions: diameter 0,46—0,50 mm; thickness 0,21—0,24 mm.

Remarks: Our specimens correspond with those described and figured by S. Geroch (1966), who shows (p. 464) "On the other hand the Carpathian specimens differ from *H. nonioninoides* (Reuss 1863) in being thicker, in having wider umbilici and fewer chambers in the last whorl" In our opinion this is a new species which S. Geroch has the priority to describe.

Hypotypes: L.P.B. 9221/1—5.

*Recurvoides imperfectus* (Hanzliková)

Pl. I, fig. 5—6

*Haplophragmoides imperfectus* (Hanzliková), 1965, p. 38, fig. 7.

*Recurvoides imperfectus* Hanzliková 1953, pl. 9, fig. 1 (nomen nudum); Geröch 1966, p. 443, fig. 10 (4—5).

Occurrence: Vraconian-Cenomanian, Teliu Valley, D. Stînii Hill, Boului Creek, Prahova Valley, Leurdei Valley.

Dimensions: diameter 0,50 mm; thickness 0,43 mm.

Hypotypes: L.P.B. 9222.

*Trochamminoides irregularis* (White)

Pl. III, fig. 17

*Trochammina irregularis* White 1928, p. 307, pl. 42, fig. 1.

*Trochamminoides irregularis* (White); Neagu 1962, p. 59, pl. 4, fig. 65; Grün & all. 1964, p. 265, pl. 4, fig. 3.

Occurrence: Cenomanian, D. Stînii Hill, Boului Creek, Prahova Valley, Leurdei Valley, Rucăr basin.

Dimensions: diameter 0,77 mm.

Hypotypes: L.P.B. 9223.

*Ammobaculites problematicus* Neagu

Pl. III, fig. 1—4

*Ammobaculites agglutinans problematicus* Neagu 1962, p. 61, pl. 2, fig. 22—24.

*Haplophragmium aequicameratum* Huss 1966, p. 32, pl. 9, fig. 10—25.

Occurrence: Vraconian-Cenomanian, Teliu Valley, D. Stînii Hill, Boului Creek, Prahova Valley, Leurdei Valley, Rucăr basin.

Dimensions: length 0,46—0,65 mm; thickness 0,12—0,14 mm.

Hypotypes: L.P.B. 9224/1—4.

*Haplophragmium aequale* (Roemer)

Pl. XII, fig. 23

*Haplophragmium aequale* (Roemer); Reuss 1862, p. 29, pl. 1, fig. 1—7; Groiss 1964, p. 7, pl. 3, fig. 3—4, pl. 4, fig. 10.

Occurrence: Cenomanian, Teliu Valley, D. Stînii Hill, Prahova Valley;

Dimensions: length 1,17—1,20 mm; thickness 0,53—0,84.

Hypotypes: L.P.B. 9316/1—10.

*Spiroplectammina gandolfii* Carbonier

Pl. III, fig. 5—8

*Spiroplectammina gandolfii* Carbonier 1952, p. 112, pl. 5, fig. 2.

Occurrence: Cenomanian, Teliu Valley, D. Stînii Hill, Prahova Valley, Leurdei Valley, Rucăr basin.

Dimensions: length 0,46—0,67 mm; breadth 0,29—0,34 mm; thickness 0,09—0,12 mm.

Hypotypes: L.P.B. 9226/1—5.

*Spiroplectammina complanata* (Reuss)

Pl. III, fig. 9—10

*Proroporus complanatus* Reuss 1860, p. 231, pl. 12, fig. 5.

*Spiroplectammina complanata* (Reuss); Neagu 1959, pl. 4, fig. 20—21.

Occurrence: Cenomanian, Teliu Valley, D. Stînii Hill, Prahova Valley.

Dimensions: length 0,58 mm; breadth 0,34 mm; thickness 0,09 mm.  
Hypotypes: L.P.B. 9227.

*Spirolectammia roemeri* Lalicker

Pl. III, fig. 11—12

*Spirolectammia roemeri* Lalicker 1935, p. 9, pl. 2, fig. 5.

*Spirolectammia laevis cretosa* Cushman; Neagu 1959, pl. 4, fig. 18—19.

Occurrence: Cenomanian, Teliu Valley, D. Stîinii Hill, Prahova Valley, Belia Valley, Leurdei Valley, Rucăr basin.

Dimensions: length 0,65 mm; breadth 0,58 mm; thickness 0,40 mm.  
Hypotypes: L.P.B. 9228.

*Textularia foeda* Reuss

Pl. I, fig. 7—8

*Textularia foeda* Reuss 1846, p. 109, pl. 43, fig. 12—13.

Occurrence: Vraconian, Prahova Valley.

Dimensions: length 0,39 mm; breadth 0,26—0,29 mm.

Hypotypes: L.P.B. 9229/1—5.

*Pseudobolivina parvissima* Neagu

Pl. I, fig. 11

*Pseudobolivina parvissima* Neagu 1970 (in press).

Occurrence: Cenomanian, Teliu Valley, D. Stîinii Hill.

Dimensions: length 0,48 mm; thickness 0,14 mm.

Hypotypes: L.P.B. 9230.

*Plectorecurvoides alternans* Noth

Pl. II, fig. 22—23

*Plectorecurvoides alternans* Noth 1952, p. 117, text-fig. 1—2; Geroch 1960, p. 54, pl. 7, fig. 4; 1962, p. 296, pl. 3, fig. 11, 15.

Occurrence: Vraconian-Cenomanian, D. Stîinii Hill, Boului Creek, Prahova Valley, Leurdei Valley.

Dimensions: diameter 0,34—0,44 mm.

Hypotypes: L.P.B. 9231/1—5.

*Trochammina umiatensis* Tappan

Pl. V, fig. 12—17

*Trochammina umiatensis* Tappan 1957, p. 214, pl. 67, fig. 27—29.

Occurrence: Vraconian-Cenomanian, Teliu Valley, D. Stîinii Hill, Prahova Valley, Boului Creek, Leurdei Valley, Rucăr basin.

Dimensions: diameter 0,65 mm.

Hypotypes: L.P.B. 9232.

*Trochammina quinqueloba* Geroch

Pl. XII, fig. 1—3

*Trochammina quinqueloba* Geroch 1959, p. 118, pl. 12, fig. 1—3; 1966, p. 452, fig. 14 (18); Neagu 1962, p. 63, pl. 2, fig. 34—36.

Occurrence: Vraconian-Cenomanian, D. Stîinii Hill, Prahova Valley.

Dimensions: diameter 0,24 mm.

Hypotypes: L.P.B. 9233.

*Verneulina polystropha* (Reuss)

Pl. III, fig. 13—14, Pl. XI, fig. 1—4

*Bulimina polystropha* Reuss, 1846, p. 109, pl. 24, fig. 53; Berthelin 1880, p. 30, pl. 2, fig. 3.

*Verneulina polystropha* (Reuss) Cushman 1937, p. 11, pl. 1, fig. 14.

Occurrence: Cenomanian, D. Stîinii Hill, Prahova Valley, Rucăr basin.

Dimensions: length 0,36—0,48 mm.

Hypotypes: L.P.B. 9235/1—3; 9236/1—5.

*Gaudryina carinata* Franke

Pl. I, fig. 15—18

*Gaudryina* (*Siphogaudryina*) *carinata* Franke; Cushman 1937, p. 76, pl. 11, fig. 12—14.

Occurrence: Cenomanian, D. Stîinii Hill, Prahova Valley, Belia Valley, Leurdei Valley, Rucăr basin.

Dimensions: length 0,43—0,46 mm; breadth 0,26—0,31 mm.

Hypotypes: L.P.B. 9237/1—5.

*Spiroplectinata annectens* (Parker & Jones)

Pl. I, fig. 10

*Spiroplectinata annectens* (Jones & Parker); Cushman 1937, p. 101, pl. 14, fig. 10—12; Neagu 1965, p. 6, pl. 2, fig. 19.

Occurrence: Vraconian-Lower Cenomanian.

Prahova Valley, Belia Valley, Leurdei Valley, Rucăr basin.

Dimensions: length 0,48 mm; breadth 0,21 mm.

Hypotypes: L.P.B. 9238.

*Tritaxia amorpha* (Cushman)

Pl. IV, fig. 6—8

*Clavulina amorpha* Cushman 1926, p. 583, pl. 17, fig. 3.

*Tritaxia amorpha* (Cushman); Huss 1966, p. 47, pl. 7, fig. 7—13.

Occurrence: Cenomanian, Teliu Valley, D. Stîinii Hill, Prahova Valley, Rucăr basin, Leurdei Valley.

Dimensions: length 0,53—0,98 mm; thickness 0,21—0,31 mm.

Hypotypes: L.P.B. 9239, 9240.

*Tritaxia gaultina carinata* (Neagu)

Pl. IV, fig. 1—5

*Clavulinoides gaultinus carinatus* Neagu 1962, p. 420, pl. 40, fig. 14—20.

Occurrence: Cenomanian, Teliu Valley, D. Stîinii Hill, Prahova Valley, Belia Valley, Leurdei Valley, Rucăr basin.

Dimensions: length 0,29—1,05 mm; thickness 0,21—0,39 mm.

Hypotypes: L.P.B. 9241/1—5.

*Tritaxia gaultina disjuncta* (Cushman)

Pl. I, fig. 13—14

*Clavulinoides disjuncta* (Cushman), 1937, p. 125, pl. 18, fig. 1—3.

Occurrence: Upper Cenomanian, Belia Valley.

Dimensions: length 0,70—0,96 mm; thickness 0,19—0,24 mm.

Hypotypes: L.P.B. 9242.

*Uvigerinammina moesiana* Neagu

Pl. XII, fig. 4—7

*Uvigerinammina moesiana* Neagu 1965, p. 5, pl. 2, fig. 11—18.

Occurrence: Vraconian-Lower Cenomanian.

Dimensions: length 0,34—0,36 mm; breadth 0,24—0,29 mm.

Hypotypes: L.P.B. 9250/1—5; 9240.

*Dorothia oxycona* (Reuss)

Pl. I, fig. 22

*Gaudryina oxycona* Reuss 1860, p. 220, pl. 12, fig. 3.

*Marsonella oxycona* (Reuss); Cushman 1937, p. 56, pl. 5, fig. 27—29, pl. 6, fig. 1—17; Ebensberger 1962, p. 22, pl. 1, fig. 1, pl. 12, fig. 10.

Occurrence: Vraconian-Cenomanian, Teliu Valley, D. Stînii Hill, Prahova Valley, Belia Valley, Leurdei Valley, Rucăr basin.

Dimensions: length 0,46 mm; thickness 0,29 mm.

Hypotypes: L.P.B. 9257.

*Dorothia pupa* (Reuss)

Pl. I, fig. 9, Pl. IV, fig. 9—13

*Textularia pupa* Reuss 1860, p. 232, pl. 13, fig. 4.

*Dorothia pupa* (Reuss); Cushman 1937, p. 78, pl. 9, fig. 20—24.

Occurrence: Cenomanian, Teliu Valley, D. Stînii Hill, Prahova Valley, Belia Valley, Leurdei Valley, Rucăr basin.

Dimensions: length 0,46—0,74 mm; breadth 0,43—0,65 mm.

Hypotypes: L.P.B. 9243/1—3; 9244/1—3.

*Dorothia conulus* (Reuss)

Pl. III, fig. 18—19, 23—24

*Textularia conulus* Reuss 1845, p. 38, pl. 8, fig. 69, pl. 13, fig. 75.

*Dorothia conula* (Reuss); Cushman 1937, p. 76, pl. 8, fig. 11—17.

*Dorothia conulus* (Reuss), Tollmann 1960, p. 162, pl. 10, fig. 7—8.

Occurrence: Cenomanian, Teliu Valley, D. Stînii Hill, Prahova Valley, Rucăr basin, Leurdei Valley.

Dimensions: length 0,40—0,66 mm.

Hypotypes: L.P.B. 9246/1—5.

*Dorothia concinna* (Reuss)

Pl. III, fig. 20—22

*Textularia concinna* Reuss 1846, p. 109, pl. 24, fig. 54.

*Dorothia concinna* (Reuss); Cushman 1937, p. 75, pl. 8, fig. 8—10.

Occurrence: Cenomanian, Teliu Valley, D. Stînii Hill, Rucăr basin.

Dimensions: length 0,62—0,98 mm.

Hypotypes: L.P.B. 9247/1—5.

*Arenobulimina conoidea* (Perner)

Pl. XI, fig. 5

*Bulimina conoidea* Perner 1892, p. 55, pl. 3, fig. 5.

*Arenobulimina conoidea* (Perner); Cushman 1937, p. 38, pl. 4, fig. 1—2; Gavor-Biedova 1969, p. 80, pl. 5, fig. 6, pl. 7, fig. 4—5.

Occurrence: Cenomanian, Prahova Valley.

Dimensions: length 0,36 mm.

Hypotypes: L.P.B. 9249/1—5.

*Arenobulimina frankei* Cushman

Pl. I, fig. 12

*Arenobulimina frankei* Cushman; Cushman; 1937, p. 37, pl. 4, fig. 21; Gawor-Biedova 1969, p. 85, pl. 5, fig. 4, 5, pl. 7, fig. 6, 7, 8, text-fig. 5, 6.

Occurrence: Vraconian-Lower Cenomanian, Prahova Valley, Rucăr basin.

Dimensions: length 0,43 mm.

Hypotypes: L.P.B. 9314/1—5.

*Arenobulimina advena* (Cushman)

Pl. V, fig. 1—7

*Hagenowella advena* Cushman; Cushman 1937, p. 174, pl. 21, fig. 3—4.

*Arenobulimina advena* (Cushman); Gawor-Biedova 1969, p. 86, pl. 8, fig. 1—4.

Occurrence: Cenomanian, D. Stînii Hill, Prahova Valley, Leurdei Valley, Rucăr basin.

Dimensions: length 0,60—0,74 mm.

Hypotypes: L.P.B. 9248/1—5.

*Plectina lenis* (Grzybowski)

Pl. III, fig. 15—16

*Spiroplecta lenis* Grzybowski 1896, p. 28, pl. 9, fig. 24—25.

*Plectina lenis* (Grzybowski); Huss 1966, p. 50, pl. 8, fig. 1—5.

Occurrence: Cenomanian, Teliu Valley, D. Stînii Hill, Prahova Valley, Leurdei Valley, Rucăr basin.

Dimensions: length 0,60—0,72 mm; thickness 0,14—0,19 mm.

Hypotypes: L.P.B. 9245/1—5.

*Quinqueloculina kochi* (Reuss)

Pl. XI, fig. 6—7

*Triloculina kochi* Reuss 1855, p. 289, pl. 11, fig. 6—7.

Occurrence: Cenomanian, Teliu Valley, D. Stînii Hill, Prahova Valley, Rucăr basin, Leurdei Valley.

Dimensions: length 0,26 mm; thickness 0,21 mm.

Hypotypes: L.P.B. 9251, 9317.

*Nodosaria lepida* Reuss

Pl. VI, fig. 1—2

*Nodosaria lepida* Reuss 1860, p. 178, pl. 1, fig. 2; Egger 1899, p. 56, pl. 6, fig. 18.

Occurrence: Cenomanian, D. Stînii Hill, Prahova Valley, Rucăr basin.

Dimensions: length 1,80—2,21 mm; thickness 0,34—0,50 mm.

Hypotypes: L.P.B. 9252/1—2.

*Nodosaria oligostegia* Reuss

Pl. VI, fig. 16

*Nodosaria oligostegia* Reuss 1845, p. 27, pl. 13, fig. 19—20.

*Dentalina oligostegia* (Reuss); Reuss 1851, p. 25, pl. 1, fig. 10; Tollmann 1960, p. 169, pl. 13, fig. 11—12.

Occurrence: Cenomanian, D. Stînii Hill, Prahova Valley,

Dimensions: length 0,50 mm; thickness 0,19 mm.

Hypotypes: L.P.B. 9253.

*Nodosaria limbata* d'Orbigny

Pl. VI, fig. 17

*Nodosaria limbata* d'Orbigny 1840, p. 12, pl. 1, fig. 1.

Occurrence: Cenomanian, D. Stîinii Hill, Rucăr basin.

Dimensions: length 1,08 mm; thickness 0,48 mm.

Hypotypes: L.P.B. 9256.

*Nodosaria rugosa* ten Dam

Pl. VII, fig. 20

*Nodosaria rugosa* ten Dam 1946, p. 575, pl. 88, fig. 7; Neagu 1965, p. 22, pl. 5, fig. 27.

Occurrence: Cenomanian, D. Stîinii Hill.

Dimensions: length 0,36 mm; thickness 0,24 mm.

Hypotypes: L.P.B. 9254.

*Nodosaria obscura* Reuss

Pl. VII, fig. 17—19

*Nodosaria obscura* Reuss 1845, p. 26, pl. 13, fig. 7—9; 1875, p. 81, pl. 20, fig. 1—4.

Occurrence: Cenomanian, Teliu Valley, D. Stîinii Hill, Prahova Valley, Leurdei Valley, Rucăr basin.

Dimensions: length 0,55—1,15 mm; thickness 0,21—0,39 mm.

Hypotypes: L.P.B. 9255/1—3.

*Dentalina pseudochrysalis* Reuss

Pl. VI, fig. 3

*Dentalina pseudochrysalis* Reuss 1862, p. 40, pl. 2, fig. 12.

Occurrence: Cenomanian, Teliu Valley, Stîinii Hill, Prahova Valley.

Dimensions: length 1,44 mm; thickness 0,26 mm.

Hypotypes: L.P.B. 9260.

*Dentalina cylindroides* Reuss

Pl. VI, fig. 9—11

*Dentalina cylindroides* Reuss 1860, p. 185, pl. 1, fig. 8.

Occurrence: Cenomanian, D. Stîinii Hill, Teliu Valley.

Dimensions: length 1,12—2,18 mm; thickness 0,26—0,39 mm.

Hypotypes: L.P.B. 9258/1—3.

*Dentalina nana* Reuss

Pl. VI, fig. 4—7

*Dentalina nana* Reuss 1862, p. 39, pl. 2, fig. 10, 18; Neagu 1959, pl. 6, fig. 19.

Occurrence: Cenomanian, Teliu Valley, D. Stîinii Hill, Prahova Valley, Rucăr basin, Leurdei Valley.

Dimensions: length 1,15—8,86 mm; thickness 0,24—0,36 mm.

Hypotypes: L.P.B. 9259/1—5.

*Dentalina legumen* Reuss

Pl. VI, fig. 8, 12, 13, 15

*Dentalina legumen* Reuss 1851, p. 10, pl. 1, fig. 14; 1860, p. 187, pl. 3, fig. 5.

Occurrence: Cenomanian, D. Stîinii Hill, Rucăr basin.

Dimensions: length 0,60—0,96 mm; thickness 0,12—0,19 mm.

Hypotypes: L.P.B. 9261/1—4.



*Pseudonodosaria mutabilis* (Reuss)

Pl. VI, fig. 19—20

*Glandulina mutabilis* Reuss 1862, p. 58, pl. 5, fig. 7—11.

Occurrence: Cenomanian, D. Stînii Hill, Rucăr basin, Prahova Valley.

Dimensions: length 0,67—1,10 mm; thickness 0,34—0,62 mm.

Hypotypes: L.P.B.: 9263/1—2.

*Marginulina linearis* Reuss

Pl. VI, fig. 21—23

*Marginulina linearis* Reuss 1862, p. 60, pl. 5, fig. 15; Neagu 1959, pl. 12, fig. 14.

Occurrence: Cenomanian, Teliu Valley, D. Stînii Hill, Prahova Valley.

Dimensions: length 0,65—1,12 mm; thickness 0,14—0,17 mm.

Hypotypes: L.P.B. 9264

*Marginulina bullata* Reuss

Pl. VIII, fig. 7

*Marginulina bullata* Reuss 1845, p. 29, pl. 13, fig. 34—38; 1860, p. 61, pl. 6, fig. 4—6; Pożaryska 1957, p. 106, pl. 12, fig. 6.

Occurrence: Cenomanian, D. Stînii Hill, Teliu Valley.

Dimensions: length 0,43 mm; thickness 0,26 mm.

Hypotypes: L.P.B. 9265.

*Marginulina cephalotes* (Reuss)

Pl. VIII, fig. 5—6

*Cristellaria cephalotes* Reuss 1862, p. 67, pl. 7, fig. 4—6

*Vaginulinopsis cephalotes* (Reuss); Neagu 1959, pl. 7, fig. 17.

Occurrence: Cenomanian, Teliu Valley, D. Stînii Hill, Prahova Valley, Leurdei Valley, Rucăr basin.

Dimensions: length 1,20—1,51 mm; thickness 0,39—0,48 mm.

Hypotypes: L.P.B. 9266/1—2.

*Lenticulina comptoni* (Sowerby)

Pl. VII, fig. 13—14

*Lenticulina comptoni* (Sowerby); Brotzen 1936, p. 48, pl. 2, fig. 5; Pożaryska 1957, p. 121, pl. 17, fig. 5, text-fig. 6—8, 12, 27.

Occurrence: Cenomanian, Teliu Valley, D. Stînii Hill, Prahova Valley, Rucăr basin, Leurdei Valley.

Dimensions: diameter 0,46—0,62 mm; thickness 0,13—0,19 mm.

Hypotypes: L.P.B. 9267/1—2.

*Lenticulina navicula* (d'Orbigny)

Pl. VII, fig. 11, 12

*Cristellaria navicula* d'Orbigny 1840, p. 27, pl. 2, fig. 19—20.

*Lenticulina navicula* (d'Orbigny); Said & Kenawy 1956, p. 131, pl. 2, fig. 9.

*Saracenaria navicula* (d'Orbigny); Tollmann 1960, p. 172, pl. 13, fig. 18.

Occurrence: Cenomanian, Teliu Valley, D. Stînii Hill, Prahova Valley.

Dimensions: length 0,43—0,48 mm; breadth 0,24—0,34 mm; thickness 0,24—0,26 mm.

Hypotypes: L.P.B. 9262/1—2.

*Lenticulina nuda* (Reuss)

Pl. VIII, fig. 20—21

*Cristellaria nuda* Reuss 1861, p. 328, pl. 6, fig. 1—3; 1862, p. 72, pl. 8, fig. 2.

Occurrence: Cenomanian, D. Stînii Hill, Prahova Valley, Rucăr basin.

Dimensions: length 0,46 mm; breadth 0,31 mm.

Hypotypes: L.P.B. 9320.

*Lenticulina subaperta* (Reuss)

Pl. VII, fig. 15—16, Pl. VIII, fig. 8—9

*Cristellaria subaperta* Reuss 1862, p. 73, pl. 8, fig. 5.

Occurrence: Cenomanian, D. Stînii Hill.

Dimensions: length 0,48—0,82 mm; breadth 0,18—0,36 mm.

Hypotypes: L.P.B. 9268.

*Lenticulina pachynota* (Reuss)

Pl. VIII, fig. 10—13

*Cristellaria pachynota* Reuss 1862, p. 69, pl. 7, fig. 11.

*Darbyella* cf. *subcretacea* Tappan; Neagu 1959, pl. 8, fig. 8—11.

Occurrence: Cenomanian, Teliu Valley, D. Stînii Hill.

Dimensions: diameter 0,72—0,74 mm; breadth 0,46—0,48 mm; thickness 0,19—0,21 mm.

Hypotypes: L.P.B. 9269/1—2.

*Astacolus crepidulus* (Fichtel & Moll)

Pl. VII, fig. 1—10

*Cristellaria (Astacolus) crepidula* (Fichtel & Moll); Gandolfi 1942, p. 57, pl. 11, fig. 2, text-fig. 14.

*Astacolus crepidula* (Fichtel & Moll); Tollmann 1960, p. 169, pl. 13, fig. 7.

Occurrence: Cenomanian, Teliu Valley, D. Stînii Hill, Prahova Valley, Leurdei Valley, Rucăr basin.

Dimensions: length 0,48—0,82 mm; breadth 0,19—0,36 mm.

Hypotypes: L.P.B. 9270/1—9.

*Vaginulina recta* Reuss

Pl. VIII, fig. 3—4

*Vaginulina recta* Reuss 1862, p. 48, pl. 3, fig. 14—15; Neagu 1965, p. 24, pl. 5, fig. 31, pl. 6, fig. 1—2.

Occurrence: Cenomanian, D. Stînii Hill, Prahova Valley, Rucăr basin.

Dimensions: Length 0,84—1,08 mm; breadth 0,14—0,17 mm.

Hypotypes: L.P.B. 9271/1—2.

*Vaginulina strombecki* Reuss

Pl. VIII, fig. 1—2

*Vaginulina strombecki* Reuss 1862, p. 46, pl. 3, fig. 8.

Occurrence: Cenomanian, Teliu Valley, D. Stînii Hill, Prahova Valley.

Dimensions: length 0,60—0,67 mm; breadth 0,24 mm.

Hypotypes: L.P.B. 9272/1—2.

*Lagena apiculata* Reuss

Pl. VIII, fig. 22—24

*Oolina apiculata* Reuss 1851, p. 22, pl. 1, fig. 1.

*Lagena apiculata* (Reuss); Reuss 1863, p. 318, pl. 1, fig. 4—8, 10—11.

Occurrence: Cenomanian, Teliu Valley, D. Stînii Hill, Prahova Valley, Rucăr basin.

Dimensions: length 0,34—0,39 mm; thickness 0,21—0,26 mm.

Hypotypes: L.P.B. 9287/1—2.

*Globulina prisca* Reuss

Pl. VIII, fig. 25—26, Pl. VII, fig. 21—22

*Globulina prisce* Reuss 1862, p. 79, pl. 9, fig. 8; Neagu 1965, p. 28, pl. 7, fig. 15—17.

Occurrence: Cenomanian, Teliu Valley, D. Stînii Hill, Prahova Valley.

Dimensions: length 0,31—0,50 mm; thickness 0,12—0,31 mm.

Hypotypes: L.P.B. 9273/1—4.

*Ramulina novaculeata* Bullard

Pl. IX, fig. 3—4

*Ramulina novaculeata* Bullard 1953, p. 346 (non pl. 46, fig. 26); Neagu 1965, p. 28, pl. 7, fig. 17—18.

Occurrence: Cenomanian, Teliu Valley, D. Stînii Hill, Prahova Valley, Leurdei Valley, Rucăr basin.

Dimensions: length 0,48—0,58 mm.

Hypotypes: L.P.B. 9274/1—2.

*Ramulina arkadelphiana* Cushman

Pl. IX, fig. 5—6

*Ramulina arkadelphiana* Cushman; Cushman 1946, p. 99, pl. 43, fig. 3—8; Neagu 1959, pl. 8, fig. 19.

Occurrence: Cenomanian, D. Stînii Hill.

Dimensions: length 0,72—1,00 mm.

Hypotypes: L.P.B. 9318/1—2.

*Ramulina globotubulosa* Cushman

Pl. IX, fig. 1—2

*Ramulina globotubulosa* Cushman; Cushman 1946, p. 100, pl. 43, fig. 10; Neagu 1959, pl. 8, fig. 17—18.

Occurrence: Cenomanian, Teliu Valley, D. Stînii Hill.

Dimensions: diameter 0,43—0,48 mm.

Hypotypes: L.P.B. 9275/1—2.

*Tristix excavata* (Reuss)

Pl. VIII, fig. 14—15

*Rhabdogonium acutangulum* Reuss 1862, p. 55, pl. 4, fig. 14.

*Tristix acutangulus* (Reuss); Kalantari 1969, p. 163, pl. 14, fig. 16—18.

Occurrence: Cenomanian, Teliu Valley, D. Stînii Hill, Prahova Valley, Rucăr basin.

Dimensions: length 0,53 mm; thickness 0,19 mm.

Hypotypes: L.P.B. 9276.

*Tristix excavata* (Reuss)

Pl. VIII, fig. 16—17

*Rhabdogonium excavatum* Reuss 1862, p. 91, pl. 12, fig. 8.

*Tristix excavata excavata* (Reuss); Michael 1967, p. 73, pl. 11, fig. 4.

Occurrence: Cenomanian, Teliu Valley, D. Stîinii Hill, Prahova Valley, Leurdei Valley, Rucăr basin.

Dimensions: length 0,40 mm; thickness 0,26 mm.

Hypotypes: L.P.B. 9277.

*Tristix maertensi* (Reuss)

Pl. VIII, fig. 18—19

*Rhabdogonium maertensi* Reuss 1862, p. 56, pl. 5, fig. 4.

*Quadratina maertensi* (Reuss); Michael 1967, p. 72, pl. 11, fig. 1.

Occurrence: Cenomanian, D. Stîinii Hill, Prahova Valley, Rucăr basin,

Dimensions: length 0,62 mm; thickness 0,26 mm.

Hypotypes: L.P.B. 9278.

*Pyramidina minima robusta* Neagû n.ssp.

Pl. IX, fig. 7—12

Holotype: L.P.B. 9279, pl. 9, fig. 11—12.

Stratum typicum: Middle-Upper Cenomanian.

Locus typicus: D. Stîinii Hill, Int. Buzăului-Covasna district, Romania.

Diagnosis: Test free, triserial, subtriangular in transversal section, with plate to feeble concave lateral faces; the 2—3 last formed chambers are globular; the sutures are smooth and feeble distinct; aperture, an elongated loop-shaped opening, is situated on the inner basal part of the last formed chamber.

Description: Test free, typical triserial; the edges of the test are right or a little twisted, acute in the early stage and rounded in the last one; the lateral faces of the test are plate or feeble concave; the feeble distinct sutures from the early stage become a little depressed to at the last 2—3 formed chambers which have more or less globular aspect. The aperture, an elongated loops-shaped opening, is situated at the base of the inner part of the last chamber.

Occurrence: Middle-Upper Cenomanian, Teliu Valley, D. Stîinii Hill, Rucăr basin.

Dimensions: holotype; length 0,31 mm; thickness 0,21 mm; paratypes: length 0,31—0,39 mm; thickness 0,16—0,19 mm.

Remarks: This new subspecies differ from the Brotzen's *P. minima* 1936, by the general aspect of the test, by the bigger size and especially by the disposition of the last formed chambers.

Holotype: L.P.B. 9279.

Paratypes: L.P.B. 9280 (Teliu Valley) (1—17; D. Stîinii Hill) 18—21; Rucăr basin.

*Eponides karsteni* (Reuss)

Pl. XIII, fig. 19—24

*Rotalia karsteni* Reuss 1855, p. 273, pl. 9, fig. 6.

*Eponides karsteni* (Reuss); Gandolfi 1942, p. 96.

Occurrence: Cenomanian, Prahova Valley.

Dimensions: diameter 0,43 mm; thickness 0,26 mm.

Hypotypes: L.P.B. 9282.

*Planulina lundegreni* Brotzen

Pl. XII, fig. 10—12; Pl. XIII, fig. 16—18

*Planulina lundegreni* Brotzen 1936, p. 181, pl. 14, fig. 1.

*Planulina* cf. *spissicostata* Cushman; Neagu 1959, p. 169, pl. 1, fig. 22—24.

Occurrence: Cenomanian, Teliu Valley, D. Stîinii Hill, Prahova Valley, Belia Valley, Leurdei Valley, Rucăr basin.

Dimensions: diameter 0,34—0,50 mm; thickness 0,09—0,17 mm.

Hypotypes: L.P.B. 9305/1—10.

*Pleurostomella obtusa* Berthelin

Pl. IX, fig. 17—19, 23—24, Pl. X, fig. 13

*Pleurostomella obtusa* Berthelin 1880, p. 29, pl. 1, fig. 9.

Occurrence: Cenomanian, Teliu Valley, D. Stîinii Hill, Prahova Valley, Belia Valley, Leurdei Valley, Rucăr basin.

Dimensions: length 0,65—0,91 mm; thickness 0,19—0,24 mm.

Hypotypes: L.P.B. 9284/1—5.

*Pleurostomella reussi* Berthelin

*Pleurostomella reussi* Berthelin, 1880, p. 28, pl. 1, fig. 10—12.

Occurrence: Cenomanian, Teliu Valley, D. Stîinii Hill, Prahova Valley, Belia Valley, Leurdei Valley, Rucăr basin.

Dimensions: length 0,91 mm; thickness 0,24 mm.

Hypotypes: L.P.B. 9283.

*Ellipsoidella macrocephala* (Storm)

Pl. IX, fig. 13—16, Pl. X, fig. 14—16, Pl. XII, fig. 8—9

*Ellipsodimorphina macrocephala* Storm 1929, p. 53, text-fig. 14.

*Nodosarella* sp. Neagu 1959, p. 161, pl. 2, fig. 16—17.

*Nodosarella coalingensis* Cushman; Neagu 1959, p. 9, fig. 6—7.

Occurrence: Cenomanian, Teliu Valley, D. Stîinii Hill, Prahova Valley, Belia Valley, Leurdei Valley, Rucăr basin.

Dimensions: length 0,40—0,53 mm; thickness 0,24—0,36 mm (A forms); length 0,29—0,31 mm; thickness 0,14—0,19 mm (B forms).

Hypotypes: 9286/1—5 (A forms), 9288/1—5 (B forms).

*Ellipsoidella gracillima* (Cushman)

Pl. X, fig. 11—12

*Nodosarella gracillims* Cushman; Cushman 1946; p. 134, pl. 55, fig. 19—21; Neagu 1959, pl. 9, fig. 11—12.

Occurrence: Cenomanian, Teliu Valley, D. Stîinii Hill, Prahova Valley, Belia Valley, Rucăr basin.

Dimensions: length 1,20—1,58 mm; thickness 0,31—0,40 mm.

Hypotypes: L.P.B. 9289/1—3.

*Nodosarella (Clarella) bulbosa* (ten Dam)

Pl. IX, fig. 25, Pl. X, fig. 6, 10

*Nodosarella bulbosa* ten Dam 1950, p. 45, pl. 3, fig. 16; Neagu 1959, pl. 9, fig. 8.

Occurrence: Prahova Valley, Cenomanian.

Dimensions: length 0,58—1,08 mm; thickness 0,21—0,29 mm.

Hypotypes: L.P.B. 9290/1—2.

*Nodosarella (Clarella) divergens* (Storm)

Pl. X, fig. 1—3

*Ellipsodimorphina divergens* Storm 1929, p. 52, text-fig. 10.

Occurrence: Middle-Upper Cenomanian, D. Stîinii Hill, Prahova Valley, Leurdei Valley, Rucăr basin.

Hypotypes: L.P.B. 9291/13; 9292/1—3.

*Nodosarella (Clarella) liebusi* (Storm)

Pl. X, fig. 4—5

*Ellipsonodosaria liebusi* Storm, 1929, p. 50, text-fig. 13.

Occurrence: Upper Cenomanian, D. Stîinii Hill, Belia Valley, Rucăr basin.

Dimensions: length 1,10—1,44 mm; thickness 0,19—0,24 mm.

Hypotypes: L.P.B. 9293/1—2 (D. Stîinii Hill), 9294 (Belia Valley).

*Nodosarella (Clarella) frequens* (Storm)

Pl. X, fig. 7—9

*Ellipsodimorphina frequens* Storm 1929, p. 52, text-fig. 12.

Occurrence: Upper Cenomanian, D. Stîinii Hill, Belia Valley.

Dimensions: length 0,43—1,41 mm; thickness 0,17—0,43 mm.

Hypotypes: L.P.B. 9295 (Belia Valley); 9296/1—3 (D. Stîinii Hill).

*Ellipsoglandulina exponens* (Brady)

Pl. X, fig. 17—18, Pl. IX, fig. 20—22

*Ellipsoglandulina exponens* (Brady); Cushman 1946, p. 137, pl. 56, fig. 34—36; Neagu 1959, pl. 9, fig. 14.

Occurrence: Cenomanian, Teliu Valley, D. Stîinii Hill, Prahova Valley.

Dimensions: length 0,40—0,50 mm; thickness 0,24—0,29 mm.

Hypotypes: L.P.B. 9297.

*Osangularia cretacea* (Carbonier)

Pl. XI, fig. 8—10

*Parella cretacea* Carbonier 1952, p. 115, pl. 5, fig. 9; Neagu 1959, pl. 11, fig. 22—24.

*Osangularia cretacea* (Carbonier); Neagu 1968, text-fig. 2 (72).

Occurrence: Vraconian-Cenomanian, Teliu Valley, D. Stîinii Hill, Prahova Valley, Belia Valley, Leurdei Valley, Rucăr basin.

Dimensions: diameter 0,43 mm.

Hypotypes: L.P.B. 9298/1—5.

*Globorotalites multisepta* (Brotzen)

Pl. XIV, fig. 1—7

*Globorotalia multisepta* Brotzen 1936, p. 161, pl. 11, fig. 6—7.

Occurrence: Middle-Upper Cenomanian, Prahova Valley, Leurdei Valley.

Dimensions: diameter 0,40—0,55 mm; thickness 0,26—0,39 mm.

Hypotypes: L.P.B. 9300/1—5.

*Globorotalites brotzeni* Hofker

Pl. XIV, fig. 8—10

*Globorotalites brotzeni* Hofker 1957, text-fig. 455.

Occurrence: Vraconian, Belia Valley.

Dimensions: diameter 0,40—0,43 mm; thickness 0,26—0,29 mm.

Hypotypes: L.P.B. 9299/1—3.

*Gyroidinoides nitidus* (Reuss)

Pl. XII, fig. 13—15, 19—22

*Rotalina nitida* Reuss 1845, p. 35, pl. 8, fig. 52, pl. 12, fig. 8, 20.

*Gyroidinoides nitidus* (Reuss); Hofker 1957, p. 393, text-fig. 437—440; Kalantari 1969, p. 197, pl. 26, fig. 9.

Occurrence: Vraconian-Cenomanian, Teliu Valley, D. Stîinii Hill, Prahova Valley, Belia Valley, Leurdei Valley, Rucăr basin.

Dimensions: diameter 0,29—0,40 mm; thickness 0,21—0,26 mm.

Hypotypes: L.P.B. 9301/1—3 (D. Stîinii Hill); 9302/1—5 (Prahova Valley).

*Gyroidinoides mauretanicus* (Carbonier)

Pl. XII, fig. 16—18, Pl. XIV, fig. 11—13

*Gyroidina mauretanicus* Carbonier 1952, p. 113, pl. 5, fig. 5; Neagu 1959, pl. 10, fig. 7—9.

*Gyroidinoides mauretanicus* (Carbonier); Neagu 1968, text-fig. 2 (66).

Occurrence: Cenomanian, Teliu Valley, D. Stîinii Hill, Prahova Valley, Belia Valley, Leurdei Valley, Rucăr basin.

Dimensions: diameter 0,31—0,48 mm; thickness 0,21—0,40 mm.

Hypotypes: L.P.B. 9303/1—2 (D. Stîinii Hill); 9304/1—3 (Belia Valley).

*Gavelinella schloembachi* (Reuss)

Pl. XV, fig. 7—12

*Rotalia schloembachi* Reuss 1862, p. 84, pl. 10, fig. 5.

*Discorbis? schloembachi* (Reuss); Neagu 1959, pl. 10, fig. 25—27

Occurrence: Cenomanian, Teliu Valley.

Dimensions: diameter 0,39 mm; thickness 0,12 mm.

Hypotypes: L.P.B. 9311/1—5.

*Heterolepa polyraphes* (Reuss)

Pl. XIII, fig. 7—15

*Rotalia polyraphes* Reuss 1845, p. 35, pl. 12, fig. 18; 1851, p. 135, pl. 3, fig. 1.

*Planorobulina polyraphes* (Reuss); Reuss 1874, p. 114, pl. 23, fig. 10.

*Cibicides polyraphes polyraphes* (Reuss); Neagu 1959, pl. 11, fig. 15—17.

*Cibicides polyraphes praeeriksdalensis* (Vasilenko); Neagu, 1959, pl. 11, fig. 18—20, 1968, text-fig. 2 (68—69).

Occurrence: Cenomanian, Teliu Valley, D. Stîinii Hill, Prahova Valley, Belia Valley, Leurdei Valley, Rucăr basin.

Dimensions: diameter 0,34—0,55 mm; thickness 0,17—0,26 mm.

Hypotypes: L.P.B. 9306/1—5.

*Orostella aumalensis* (Sigal)

Pl. XI, fig. 11—19; Pl. XIII, fig. 1—6; Pl. XIV, fig. 14—16;

Pl. XV, fig. 13—17

*Anomalina aumalensis* Sigal 1952, p. 27, text-fig. 26; Kalantari 1969, p. 211, pl. 22, fig. 4—6.

*Gavelinella aumalensis* Sigal var. 1962, pl. 13, fig. 6.

*Orostella aumalensis* (Sigal); Butt 1966, p. 178.

*Valvulineria allomorphinoides* (Reuss); Neagu 1959, pl. 9, fig. 19—24; pl. 10, fig. 1—6, pl. 11, fig. 12—14.

Occurrence: Middle-Upper Cenomanian, Teliu Valley, D. Stîinii Hill, Prahova Valley, Belia Valley, Leurdei Valley, Rucăr basin.

Dimensions: diameter 0,34—0,60 mm; thickness 0,17—0,29 mm.

Hypotypes: L.P.B. 9309/1—9 (D. Stîinii Hill), 9310/1—10 (Teliu Valley).

*Lingulogavelinella globosa* (Brotzen)

Pl. XV, fig. 1—6

*Anomalinoides globosa* Brotzen 1945, p. 55, pl. 2, fig. 6.

Occurrence: Middle-Upper Cenomanian, Teliu Valley, Leurdei Valley.

Dimensions: diameter 0,31—0,46 mm; thickness 0,21—0,24 mm.

Hypotypes: L.P.B. 9312/1—8 (Leurdei Valley), 9313 (Teliu Valley).

WYKAZ LITERATURY

REFERENCES

- Beissel J. (1891), Die Foraminiferen der Aachener Kreide. *Abh. Preuss. Geol. Landesanst.*, n. ser. 3.
- Belford D.J. (1960), Upper Cretaceous foraminifera from the Toolong calcilutite and Gingin Chalk, Western Australia. *Bur. Min. Resources, Geol.-Geophys. Bull.* 57.
- Berthelin M. (1880), Mémoire sur les foraminifères fossiles de l'étage Albien de Montcley (Doubs). *Mém. Soc. Geol. France*, ser. 3, 1.
- Brotzen Fr. (1936), Foraminiferen aus dem Schwedischen Unter-Senon von Eriksdal in Schonen. *Sver. Geol. Unders.*, ser. C, 396.
- Brotzen Fr. (1942), Die Foraminiferengattung Gavelinella n. gen. und die Systematik der Rotaliiformes. *Sver. Geol. Unders.* ser. C. 452.
- Brotzen Fr. (1945), De geologiska resultaten från boringarna vid Höllviken; Preliminär rapport Del I: Kritan. *Sver. Geol. Unders.* ser. C. 465.
- Bullard F.J. (1953), Polymorphinidae of the Cretaceous (Cenomanian) Del Rio Shale. *J. Paleont.* 27, 3.
- Butt A.A. (1966), Foraminifera of the type Turonian. *Micropaleontology*, 12, 2.
- Carbonier A. (1952), Sur un gissement de foraminifères d'âge Cenomanien supérieur, provenant de la région de Taza (Maroc). *Bull. Soc. Geol. France*, ser. 16, vol. 2, fasc. 1—3.
- Cushman J.A. (1910), A monograph of the foraminifera of the North Pacific Ocean, pt. 1. *U.S. Nat. Mus., Bull.* 71.
- Cushman J.A. (1918), Foraminifera of the Atlantic Ocean, *U.S. Mus. Bull.* 104.
- Cushman J.A. (1926), The foraminifera of the Velasco Shale of the Tampico Embayment. *Bull. Amer. Assoc. Petrol. Geol.*, 10, no. 6.
- Cushman J.A. (1937), A monograph of the foraminiferal family Verneuilinidae. *Cush. Labor. Foram. Research. Spec. Publ.* no. 7.
- Cushman J.A. (1937) b) A monograph of the foraminiferal family Valvulinidae, *Cush. Labor. Foram. Research. Spec. Publ.* no. 8.
- Cushman J.A. (1946), Upper Cretaceous foraminifera of the Gulf Coastal region of United States and adjacent areas. *U.S. Geol. Survey, Prof. Paper* 206.
- Dam A. ten (1946), Arenaceous foraminifera and Lagenidae from the Neocomian (Lower Cretaceous) of the Netherlands. *J. Paleont.* vol. 20, no. 6.
- Dam A. ten (1950), Les foraminifères de l'Albien des Pays Bas. *Mém. Soc. Geol. France*, n. ser., vol. 29, pt. 4, no. 63.
- Ebensberger H. (1962), Stratigraphische und Mikropaläontologische Untersuchungen in der Aachener Oberkreide, besonders der Maastricht-Stufe. *Palaeontographica*, 120 A.
- Egger J.G. (1899), Foraminiferen und Ostracoden aus den Kreidemergeln der Oberbayerischen Alpen. *K. Bayer. Akad. Wiss. München, Math.-Physik, Cl., Abh.*, 21, no. 1.
- Franke A. (1928), Die Foraminiferen der Oberen Kreide Nord- und Mitteldeutschlands. *Abh. Preuss. Geol. Landesanst.*, n. ser., vol. 111.



- Frizzell L.D. (1954), Handbook of Cretaceous foraminifera of Texas. *Texas Univ. Bur. Econ. Geol. Rept. Investig.*, no 22.
- Geroch S. (1959), Stratigraphic significance of the arenaceous foraminifera in the Carpathian Flysch. *Paläont. Z.*, vol. 33, no. 1—2.
- Geroch S. (1960), Microfaunal assemblages from the Cretaceous and Paleogene Silesian Unit in the Beskid Śląski Mts. (Western Carpathians). *Bull. Inst. Geol.*, 153.
- Geroch S., (1966), Lower Cretaceous small foraminifera of the Silesian series Polish Carpathians. *Ann. Soc. Geol. Pol.*, vol. 36, fasc. 4.
- Gandolfi R. (1942), Recerche micropaleontologiche e stratigrafiche sulla Scaglia et sul Flysch Cretacici dei dintorni di Balerna (Canton Ticino). *Riv. Ital. Paleont.*, 48, mem. 4.
- Gawor-Biedowa E. (1969), The genus *Arenobulimina* Cushman from the Upper Albian and Cenomanian of the Polish Lowlands. *Ann. Soc. Geol. Pol.*, vol. 39, fasc. 1—3.
- Groiss J.T. (1964), Eine Mikrofauna aus des albüberdeckenden Kreide der südlichen Frankenalb. *Abh. Erlanger Geol.*, H. 53.
- Graham J. and Cusch C. (1963), Campanian foraminifera from the Stanford University campus California. *Stanford Univ. Publ. Geol. Sci.*, 8, no. 1.
- Glaessner M. (1937), Studien über Foraminiferen aus der Kreide und dem Tertiär des Kaukasus. *Moskow Univ. Labor. Palaeont., Probl. Palaeont.* 2—3.
- Grün W., Lauer G., Niedermayr G., Schnabel W. (1964), Die Kreide-Tertiär-Grenze im Wienerwaldflysch bei Hochstrass. *Verh. Geol. Bundesanst. H. 2.*
- Grzybowski J. (1896), Otwornice czerwonych ilów z Wadowic. *Rozpr. Wydz. math.-przyr. Akad. Umiej.*, 30.
- Grzybowski J. (1896), Otwornice pokładów naftonośnych okolicy Krosna. *Rozpr. Wydz. math.-przyr. Akad. Umiej.* 33.
- Hanzliková E. (1966), Die Foraminiferen der Lhoty-Schichten. *Acta Musei Moraviae*, 5, sect. St. Natur.
- Hofker J. (1957), Foraminiferen der Oberkreide von Nordwestdeutschland und Holland. *Jb. Geol.* 27.
- Huss F. (1957), Stratigraphy of the Węglówka unit in the light of its microfauna. *Acta geol. pol.*, 7.
- Huss F. (1966), Les foraminifères agglutinats de la serie sous-silesienne de l'unité petrolifère de Węglówka. *Pol. Akad. Nauk. Komis. Nauk. Geol., Pr. geol.* 34.
- Kalantari A. (1969), Foraminifera from the Middle Jurassic-Cretaceous successions of Koppet-Dagh region. (N. E. Iran). *National Iran. Oil Comp. Geol. Labor.*, Publ. no. 3.
- Lalicker C. G. (1935), New Cretaceous Textulariidae. *Cush. Labor. Foram. Research. Contrib.*, 11, pt. 1.
- Loeblich A.R.Jr., Tappan H. (1964), Sarcodina, in R. C. Moore, Treatise on invertebrate Paleontology, part. C, Protista 2.
- Malapris M. (1965), Les Gavelinellidae et formes affines du gisement Albien de Courcelles (Aube). *Rev. Micropal.* 8, no. 3.
- Malapris M. (1967), Les Lingulogavelinelles de l'Albien inférieur et moyen de l'Aube. *Rev. Micropal.* 10, no. 2.
- Marie P. (1941), Les foraminifères de la Craie a *Belemnitella mucronata* du Bassin de Paris. *Mus. Nat. Hist., Mém. n. ser.*, 12, fasc. 1.
- Marks P. (1967), Foraminifera from the Craie de Theligny (Cenomanian, dept. Sarthe, France). *Koninkl. Nederl. Akad. Wetensch., Amsterdam. Proc. ser. B*, 70, no. 4.
- Marks P. (1968), Smaller foraminifera from the „Couches à *Orbitolina complanata*” (Cenomanian) at Bailon (Sarthe, France). *Koninkl. Nederl. Akad. Wetensch., Amsterdam, Proc. ser. B*, vol. 71, no. 1.

- Michael E. (1967), Die Mikrofauna des NW-deutschen Barreme. Teil. 1. Die Foraminiferen des NW-deutschen Barreme. *Palaeontographica*, Suppl. 12.
- Neagu T. (1959), Studiul micropaleontologic al Cretacicului superior din Valea Teliu. *Ann. Univ. Buc., ser. St. Natur.* no. 12.
- Neagu T. (1962), Studiul foraminiferelor aglutinante din argilele cretacic superioare de pe Valea Sadova și basinul superior al Văii Buzăului. *Acad. Rom. Stud. Cercet. Geol.*, 7, no. 1.
- Neagu T. (1962 b), *Clavulinoides gaultinus* (Morozowa) 1948 (Foraminifera) dans le Flysch crétacé en Roumanie. *Ann. Soc. Geol. Pol.* 32, fasc. 3.
- Neagu T. (1965), Albian foraminifera of the Rumanian Plain. *Micropaleontology*, 11, no 1.
- Neagu T. (1968), Biostratigraphy of Upper Cretaceous deposits in the southern Eastern Carpathians near Brasov, *Micropaleontology*, 14, no. 2.
- Neagu T. (1970), Micropaleontological and stratigraphical study of the Upper Cretaceous deposits (between the Upper Valleys of the Buzău and Rîul Negru Rivers). *Mem. Inst. Geol.* 12.
- Noth R. (1951), Foraminiferen aus Unter- und Oberkreide des Oesterreichischen Anteils an Flysch, Helvetikum und Vorlandvorkommen. *Jb. Austria Geol. Bundesanst.* Sonderband 3.
- Noth R. (1952), *Plectorecurvoides* eine neue Foraminiferengattung. *Verh. Geol. Bundesanst.* no. 3.
- Orbigny A. D. (1840), Mémoire sur les foraminifères de la Craie Blanche des Paris. *Mém. Soc. Geol. France*, vol. 4, no. 1.
- Perner J. (1892), Über die Foraminiferen des boehmischen Cenomans. *Palaeontographica Bohemiae*, 4.
- Požaryska K. (1957), Lagenidae du Crétacé supérieur de Pologne. *Palaeont.*, no. 3.
- Reuss A. E. (1845—46), Die Versteinerungen der böhmischen Kreideformation, Stuttgart.
- Reuss A. E. (1851), Die Foraminiferen und Entomostraceen des Kreidemergels von Lemberg. *Abh. Haidinger's Naturwis.* 4.
- Reuss A.E. (1855), Ein Beitrag zur genaueren Kenntniss des Kreidegebilde Mecklenburgs. *Z. Deutsch. Geol. Ges.* 7, fasc. 1.
- Reuss A. E. (1860), Die Foraminiferen der Westphalischen Kreideformation. *Sitzber. k. k. Akad. Wiss. Wien, Math-Naturw. Cl.* 40.
- Reuss A. E. (1862), Die Foraminiferen des norddeutschen Hils und Gault. *Sitzber. k. k. Akad. Wiss. Wien, Math.-Naturw. Cl.*, 46.
- Reuss A.E. (1874), Die Foraminiferen, Bryozoen und Ostracoden des Pläners. *Palaeontographica*, vol. 20, pt. 2.
- Said R., Kenawy A. (1956), Upper Cretaceous and Lower Tertiary foraminifera from Northern Sinai, Egypt. *Micropaleontology*, vol. 2, no. 2.
- Sigal J. (1952), Aperçu stratigraphique sur la micropaleontologie du Crétacé. *XIX Cong. Geol. Intern. Monogr. region, ser. 1, Algerie*, 26.
- Sigal J., Dardenne M. (1962), Correlations dans la craie du Bassin de Paris. Perimetre de Dammartin-en-Geole. *Ann. Soc. Geol. Nord* 80.
- Sliter W. V. (1968), Upper Cretaceous foraminifera from southern California and northwestern Baja California, Mexico. *Univ. Kansas Paleont. Contrib. Protozoa*, art. 7.
- Storm H. (1929), Zur Kenntnis der Foraminiferenfauna in Oberturon und Emscher der Boehmischen Kreideformation. *Lotos*, 77, fas., 3.
- Tappan H. (1940), Foraminifera from the Grayson Formation of northern Texas. *J. Paleont.* 14, no. 5.
- Tappan H. (1943), Foraminifera from the Duck Creek Formation of Oklahoma and Texas. *J. Paleont.* 17, no. 5.

- Tappan H. (1957), New Cretaceous index foraminifera from northern Alaska. *Bull. U.S. Nat. Mus.*, 215.
- Tappan H. (1962), Foraminifera from the Arctic Slope of Alaska, pt. 3 Cretaceous foraminifera. *U.S. Geol. Survey, Prof. Paper*, 236-C.
- Tollmann A. (1960), Die Foraminiferen des Oberconiac aus der Gosau des Ausseer Weissenbachtals in Steiermark. *Jb. Geol. Bundesanst.* 103.
- Vašiček M. (1947). Remarks on the microbiostratigraphy of Magura Flysch in Moravia. *Vest. Stat. Geol. Ustav.*, 22.
- White M.P., (1928), Some index foraminifera of the Tampico Embayment Area of Mexico. *J. Paleont.* 2, no. 3, no. 4.

## STRESZCZENIE

Artykuł niniejszy jest drugą częścią pracy zamieszczonej w Roczniku Pol. Tow. Geol. tom 30 zeszyt 1—3 (1969) dotyczącej fauny otwornic cenomanu południowej części Wschodnich Karpat rumuńskich. Szczegółowa biostatygrafia tych osadów opiera się na otwornicach planktonicznych i częściowo na otwornicach bentonicznych. Wykaz systematyczny otwornic bentonicznych jest poprzedzony dyskusją odnośnie do następstwa wyróżnionych poziomów otwornic planktonicznych.

## EXPLANATION OF PLATES OBJAŚNIENIA TABLIC

### PLATE — TABLICA I

- Fig. 1—3. *Kalamopsis grzybowskii* (Dylażanka), Boului Creek, L.P.B. 9214/1—3; ×70
- Fig. 4. *Hippocrepina depressa* Vašiček, Boului Creek, L.P.B. 9206; ×70
- Fig. 5—6. *Recurvoides imperfectus* (Hanzliková), D. Stînii Hill, L.P.B. 9222; ×70
- Fig. 7—8. *Textularia foeda* Reuss, Prahova Valley, L.P.B. 9229/1—2; ×70
- Fig. 9. *Dorothia pupa* (Reuss), Belia Valley, L.P.B. 9243; ×65
- Fig. 10. *Spiroplectinata annectens* (Jones & Parker), Belia Valley, L.P.B. 9238; ×85
- Fig. 11. *Pseudobolivina parvissima* Neagu, D. Stînii Hill, L.P.B. 9230; ×85
- Fig. 12. *Arenobulimina frankei* Cushman, Podișor Hill (Rucăr basin), L.P.B. 9314/1; ×85
- Fig. 13—14. *Tritaxia gaultina disjuncta* (Cushman), Belia Valley, L.P.B. 9242/1—2; ×70
- Fig. 15—18. *Gaudryina carinata* Franke, Belia Valley, L.P.B. 9237/1—2; ×70
- Fig. 19—21. *Haplophragmoides* aff. *nonioninoides* (non Reuss), Geroch, Boului Creek, L.P. 9221; ×80
- Fig. 22. *Dorothia oxycona* (Reuss) L.P.B. 9257; ×75

### Plate — Tablica II

- Fig. 1. *Psammosphaera fusca* Schultze, D. Stînii Hill, L.P.B. 9207; ×75
- Fig. 2. *Hyperammina gaultina*, ten Dam, D. Stînii Hill, L.P.B. 9321; ×85
- Fig. 3—4. *Reophax deckeri* Tappan, D. Stînii Hill, L.P.B. 9216/1—2; ×70
- Fig. 5. *Hormosina ovulum* (Grzybowski), D. Stînii Hill, L.P.B. 9215; ×100
- Fig. 6—10. *Glomospira charoides* (Jones & Parker), D. Stînii Hill, L.P.B. 9210; ×50
- Fig. 11. *Glomospira irregularis* (Grzybowski), D. Stînii Hill, L.P.B. 9212; ×65

- Fig. 12. *Ammodiscus cretaceus* (Reuss), D. Stinii Hill, L.P.B. 9208;  $\times 50$   
Fig. 13. *Reophax minuta* Tappan, D. Stinii Hill, L.P.B. 9217;  $\times 70$   
Fig. 14—15. *Glomospira gordialis* (Jones & Parker), D. Stinii Hill, L.P.B. 9211/  
1—2;  $\times 50$   
Fig. 16—18. *Lituotuba incerta* Franke, D. Stinii Hill, L.P.B. 9213/1;  $\times 50$   
Fig. 19. *Ammodiscus infimus* Franke, D. Stinii Hill, L.P.B. 9209/2;  $\times 75$   
Fig. 20—21. *Haplophragmoides bulloides* (Beissel), D. Stini Hill, L.P.B. 9219;  $\times 70$   
Fig. 22—23. *Plectorecurvoides alternans* Noth, D. Stinii Hill, L.P.B. 9231/1;  $\times 60$

Plate — Tablica III

- Fig. 1—4. *Ammobaculites problematicus* Neagu, D. Stinii Hill, L.P.B. 9224/1—4;  
 $\times 75$   
Fig. 5—8. *Spiroplectammina gandolfii* Carbonier, D. Stinii Hill, L.P.B. 9226/1;  
 $\times 100$   
Fig. 9—10. *Spiroplectammina complanata* (Reuss), D. Stinii Hill, L.P.B. 9227;  $\times 75$   
Fig. 11—12. *Spiroplectammina roemeri* Lalicker, D. Stinii Hill, L.P.B. 9228;  $\times 65$   
Fig. 13—14. *Verneuilina polystropha* (Reuss), D. Stinii Hill, L.P.B. 9235/1—2;  $\times 75$   
Fig. 15—16. *Plectina lenis* (Grzybowski), D. Stinii Hill, L.P.B. 9245/1—2;  $\times 70$   
Fig. 17. *Trochamminoides irregularis* (White), D. Stinii Hill, L.P.B. 9225;  $\times 70$   
Fig. 18—19, 23—24. *Dorothia conulus* (Reuss), D. Stinii Hill, L.P.B. 9246/1—2;  $\times 75$   
Fig. 20—22. *Dorothia concina* (Reuss), D. Stinii Hill, L.P.B. 9247/1—2;  $\times 75$

Plate — Tablica IV

- Fig. 1—5. *Tritaxia gaultina carinata* (Neagu), D. Stinii Hill, L.P.B. 9241/1—4;  $\times 75$   
Fig. 6—8. *Tritaxia amorpha* (Cushman), D. Stinii Hill, L.P.B. 9239 (fig. 6—7),  
9240 (fig. 8);  $\times 75$   
Fig. 9—13. *Dorothia pupa* (Reuss), D. Stinii Hill, L.P.B. 9244/1—5;  $\times 75$   
Fig. 14—19. *Haplophragmoides kirki* Wickenden, D. Stinii Hill, L.P.B. 9220/1—5;  
 $\times 75$

Plate — Tablica V

- Fig. 1—7. *Arenobulimina advena* (Cushman), D. Stinii Hill, L.P.B. 9248/1—5;  $\times 75$   
Fig. 8—11. *Haplophragmoides gigas minor* Nauss, D. Stinii Hill, L.P.B. 9218/1—2;  
 $\times 50$   
Fig. 12—17. *Trochammina umiatensis* Tappan, D. Stinii Hill, L.P.B. 9232 (fig. 15—17),  
9234 (fig. 12—14);  $\times 75$

Plate — Tablica VI

- Fig. 1—2. *Nodosaria lepida* Reuss, D. Stinii Hill L.P.B. 9252/1—2;  $\times 35$   
Fig. 3. *Dentalina pseudochrysalis* Reuss, D. Stinii Hill, L.P.B. 9260;  $\times 50$   
Fig. 4—7. *Dentalina nana* Reuss, D. Stinii Hill, L.P.B. 9259/1—4;  $\times 40$   
Fig. 8, 12, 13, 15. *Dentalina legumen* Reuss, D. Stinii Hill, L.P.B. 9261/1—4;  $\times 40$   
Fig. 9, 10, 11. *Dentalina cylindroides* Reuss, D. Stinii Hill, L.P.B. 9258/1—3; (fig. 9  
 $\times 45$ , fig. 10, 11  $\times 35$ )  
Fig. 14. *Dentalina* sp., Dealul Stinii Hill.  
Fig. 16. *Nodosaria oligostegia* Reuss, D. Stinii Hill, L.P.B. 9253;  $\times 70$   
Fig. 17. *Nodosaria limbata* d'Orbigny, D. Stinii Hill, L.P.B. 9256;  $\times 35$   
Fig. 18. *Dentalina?* sp. D. Stinii Hill.  
Fig. 19—20. *Pseudonodosaria mutabilis* (Reuss), D. Stinii Hill, L.P.B. 9263/1—2;  $\times 45$   
Fig. 21—23. *Marginulina linearis* (Reuss), D. Stinii Hill, L.P.B. 9264/1—3;  $\times 75$   
Figs. 6, 7, 9, 17, the tests with many borings made by the young gastropods

Plate — Tabela VII

- Fig. 1—10. *Astacolus crepidulus* (Fichtell & Moll), D. Stînii Hill, L.P.B. 9270/1—9;  $\times 70$   
Fig. 11—12. *Lenticulina navicula* (d'Orbigny), D. Stînii Hill, L.P.B. 9262/1—2;  $\times 75$   
Fig. 13—14. *Lenticulina comptoni* (Sowerby), D. Stînii Hill, L.P.B. 9267;  $\times 70$   
Fig. 15—16. *Lenticulina subaperta* (Reuss), D. Stînii Hill, L.P.B. 9268/2;  $\times 75$   
Fig. 17—19. *Nodosaria obscura* Reuss, D. Stînii Hill, L.P.B. 9254;  $\times 70$   
Fig. 20. *Nodosaria rugosa* ten Dam, D. Stînii Hill, L.P.B. 9254;  $\times 70$   
Fig. 21—22. *Globulina prisca* Reuss, D. Stînii Hill, L.P.B. 9273/1—2;  $\times 60$

Plate — Tablica VIII

- Fig. 1—2. *Vaginulina strombecki* Reuss, D. Stînii Hill, L.P.B. 9272/1—2;  $\times 70$   
Fig. 3—4. *Vaginulina recta* Reuss, D. Stînii Hill, L.P.B. 9271/1—2;  $\times 70$   
Fig. 5—6. *Marginulina cephalotes* (Reuss), D. Stînii Hill, L.P.B. 9266/1—2; (fig. 5  $\times 35$ , fig. 6  $\times 54$ )  
Fig. 7. *Marginulina bullata* Reuss, D. Stînii Hill, L.P.B. 9265;  $\times 70$   
Fig. 8—9. *Lenticulina subaperta* (Reuss), D. Stînii Hill, L.P.B. 9268/1;  $\times 45$   
Fig. 10—13. *Lenticulina pachynota* (Reuss). D. Stînii Hill, L.P.B. 9269/1—2;  $\times 55$   
Fig. 14—15. *Tristix acutangula* (Reuss), D. Stînii Hill, L.P.B. 9276;  $\times 70$   
Fig. 16—17. *Tristix excavata* (Reuss), D. Stînii Hill, L.P.B. 9277;  $\times 70$   
Fig. 18—19. *Tristix maertensi* (Reuss), D. Stînii Hill, L.P.B. 9278;  $\times 70$   
Fig. 20—21. *Lenticulina nuda* (Reuss), D. Stînii Hill, L.P.B. 9320;  $\times 80$   
Fig. 22—24. *Lagena apiculata* Reuss, D. Stînii Hill, L.P.B. 9287/1—3;  $\times 75$   
Fig. 25—26. *Globulina prisca* Reuss, D. Stînii Hill, L.P.B. 9273/3—4;  $\times 85$

Plate — Tablica IX

- Fig. 1—2. *Ramulina globotubulosa* Cushman, D. Stînii Hill, L.P.B. 9275/1—2;  $\times 90$   
Fig. 3—4. *Ramulina novaculeata* Bullard, D. Stînii Hill, L.P.B. 9274/1—2;  $\times 90$   
Fig. 5—6. *Ramulina arkadelphiana* Cushman, D. Stînii Hill, L.P.B. 9318/1—2;  $\times 90$   
Fig. 7—12. *Pyramidina minima robusta* Neagu n. sp. fig. 11—12, holotype, L.P.B. D. Stînii Hill, L.P.B. 9279, fig. 7 paratype, Teliu Valley, L.P.B. 9280, fig. 8, 9—10, D. Stînii Hill, L.P.B. 9280/2—3;  $\times 100$   
Fig. 13—16. *Ellipsoidella macrocephala* (Storm), D. Stînii Hill, L.P.B. 9288/2—3;  $\times 90$   
Fig. 17—19, 23—24. *Pleurostomella obtusa* Berthelin, D. Stînii Hill, L.P.B. 9284/1—4;  $\times 75$   
Fig. 20—22. *Ellipsoglandulina exponens* (Brady), D. Stînii Hill, L.P.B. 9297/3—5;  $\times 80$   
Fig. 25. *Nodosarella (Clarella) bulbosa* (ten Dam), Prahova Valley, L.P.B. 9290/2;  $\times 75$

Plate — Tablica X

- Fig. 1—3. *Nodosarella (Clarella) divergens* (Storm), D. Stînii Hill, L.P.B. 9292/1—3;  $\times 45$   
Fig. 4—5. *Nodosarella (Clarella) liebusi* (Storm), D. Stînii Hill, L.P.B. 9293/1—2;  $\times 70$   
Fig. 6, 10. *Ellipsoidella bulbosa* (ten Dam) Prahova Valley, L.P.B. 9290/1;  $\times 70$   
Fig. 7—9. *Nodosarella (Clarella) frequens* (Storm), D. Stînii Hill, L.P.B. 9296/1—3;  $\times 45$   
Fig. 11—12. *Ellipsoidella gracillima* (Cushman), D. Stînii Hill, L.P.B. 9289/1—2;  $\times 50$   
Fig. 13. *Pleurostomella obtusa* Berthelin, D. Stînii Hill, L.P.B. 9284/5;  $\times 70$

- Fig. 14—16. *Ellipsoidella macrocephalla* (Storm) microsphaerical generation? D. Stînii Hill, L.P.B. 9286/1—2; ×75  
Fig. 17—18. *Ellipsoglandulina exponens* (Brady), D. Stînii Hill, L.P.B. 9297/1—2; ×70

Plate — Tablica XI

- Fig. 1—4. *Verneuilina polystropha* (Reuss), Prahova Valley, L.P.B. 9236/1—2; ×85  
Fig. 5. *Arenobulimina conoidea* (Perner), Prahova Valley, L.P.B. 9249; ×85  
Fig. 6—7. *Quinqueloculina kochi* (Reuss), D. Stînii Hill, L.P.B. 9251; ×120  
Fig. 8—10. *Osangularia cretacea* (Carbonier), D. Stînii Hill, L.P.B. 9298; ×90  
Fig. 11—19. *Orostella aumalensis* (Sigal), D. Stînii Hill, L.P.B. 9309/5—8; ×95

Plate — Tablica XII

- Fig. 1—3. *Trochammina quinqueloba* Geroch, D. Stînii Hill, L.P.B. 9233; ×115  
Fig. 4—7. *Uvigerinammina moesiana* Neagu, Prahova Valley, L.P.B. 9250/1—2; ×90  
Fig. 8—9. *Ellipsoidella macrocephalla* (Storm) D. Stînii Hill, L.P.B. 9288/1; ×90  
Fig. 10—12. *Planulina lundegreni* Brotzen, Belia Valley, L.P.B. 9305/1; ×85  
Fig. 13—15. 19—22. *Gyroidinoides nitidus* (Reuss), fig. 13—21, Prahova Valley; L.P.B. 9302/1—2; ×85; fig. 22. D. Stînii Hill, L.P.B. 9301/3; ×85  
Fig. 16—18. *Gyroidinoides mauretanicus* (Carbonier), D. Stînii Hill, L.P.B. 9303/1; ×85  
Fig. 23. *Haplophragmium aequale* (Roemer), D. Podișor (Rucăr basin), L.P.B. 9316/1; ×70

Plate — Tablica XIII


- Fig. 1—6. *Orostella aumallensis* (Sigal), D. Stînii Hill, L.P.B. 9309/1—3; ×85  
Fig. 7—15. *Heterolepa polyraphes* (Reuss), D. Stînii Hill, L.P.B. 9306/1, 3; ×85  
Fig. 16—18. *Planulina lundegreni* Brotzen, Belia Valley, L.P.B. 9305/2; ×95  
Fig. 19—21. *Eponides karsteni* (Reuss), Prahova Valley, L.P.B. 9282; ×85


Plate — Tablica XIV

- Fig. 1—7. *Globorotalites multisepta* (Brotzen), Prahova Valley, L.P.B. 9300/1—3; ×85  
Fig. 8—10. *Globorotalites brotzeni* Hofker, Belia Valley, L.P.B. 9299; ×85  
Fig. 11—13. *Gyroidinoides mauretanicus* (Carbonier), D. Stînii Hill; L.P.B. 9303/2; ×85  
Fig. 14—16. *Orostella aumallensis* (Sigal), D. Stînii Hill;

Plate — Tablica XV


- Fig. 1—6. *Lingulogavelinella globosa* (Brotzen), Leurdei Valley, L.P.B. 9312/1—2; ×85  
Fig. 7—12. *Gavelinella schloembachi* (Reuss), Teliu Valley, L.P.B. 9311/1—2; ×85  
Fig. 13—17. *Orostella aumalensis* (Sigal), D. Stînii Hill, L.P.B. 9309/6; ×85







T. Neagu









*T. Neagu*








T. Neagu



T. Neagu










T. Neagu









T. Neagu





T. Neagu



T. Neagu