

PROLIFIC DEVELOPMENT OF PACHYTHECALIINES IN LATE BARREMIAN, BULGARIA: CORAL TAXONOMY AND SEDIMENTARY ENVIRONMENT

Bogusław KOŁODZIEJ¹, Marin IVANOV² & Vyara IDAKIEVA²

¹ *Institute of Geological Sciences, Jagiellonian University, ul. Oleandry 2a, 30-063 Kraków, Poland;
e-mail: boguslaw.kolodziej@uj.edu.pl*

² *Department of Geology, Paleontology and Fossil Fuels, Sofia University 'St. Kliment Ohridski', 15 Tzar Osvoboditel
Bd., 1000 Sofia, Bulgaria; e-mail: mivanov@gea.uni-sofia.bg; idakieva@gea.uni-sofia.bg*

Kołodziej, B., Ivanov, M. & Idakieva, V., 2012. Prolific development of pachythecaliines in Late Barremian, Bulgaria: coral taxonomy and sedimentary environment. *Annales Societatis Geologorum Poloniae*, 82: 291–330.

Abstract: Diversified and abundant corals of the suborder Pachythecaliina (order Hexanthiniaria) are described from Upper Barremian, biostromal reefs of the Emen Formation, Lovech Urganian Group, north central Bulgaria. The corals are mostly of the phaceloid growth form and represent 14 species (six new), 12 genera (three new), belonging to five families. Pachythecaliines occur with the small, monopleurid cylindrical rudist *Mathesia darderi*. The rudists frequently are densely clustered, occur between coral branches or are in contact with them. Other corals, with the exception of the phaceloid *Calamophylliopsis*, and other rudists, are rare. Non-laminated microbialite crusts provided additional, structural support for bioconstruction development. Microbialites (auto-micrites) can be interpreted as a product of microbial activity, or alternatively, as a result of carbonate precipitation, brought about by non-living organic substrates (organomineralization s.s.). In addition to microbialites, metazoans are encrusted by heterotrophic skeletal microorganisms, while photophilic and oligotrophic micro-encrusters, usually common in other coral-bearing limestones of the Emen Formation, are very rare. The section at the Rusalya Quarry (NW of Veliko Tarnovo), about 42 m thick, provides the sedimentary and environmental context for the reefal biostromes. The vertical biotic and sedimentary succession displays a general shallowing trend: from the outer carbonate platform with bioclastic limestones containing small boundstone patches (corals, but not pachythecaliines, *Lithocodium aggregatum*), to the inner platform with rudist biostromes. The pachythecaliine-rich biostromes, 2.5 m thick, were developed in a low-energy environment, referred to the distal part of the rudist-dominated area of the platform. The development of microbialites was facilitated by a low sedimentation rate, and possibly by increased nutrient level. Only poorly diversified and non-phaceloid pachythecaliines occur in other coral-rich limestones and marls of the Urganian complex in Bulgaria. The assemblage described is the most remarkable, Early Cretaceous coral community worldwide, with regard to pachythecaliines. Phaceloid pachythecaliines are only more common in the Upper Jurassic rocks, being particularly diversified in the Tithonian–Lower Berriasian Štramberk Limestone (Czech Republic) and its equivalent in the Polish Outer Carpathians. However, their sedimentary context differs from that described for the corals of the Emen Formation.

Key words: corals, taxonomy, carbonate platform, palaeoecology, Cretaceous, Fore-Balkan, Bulgaria.

Manuscript received 12 November 2012, accepted 20 December 2012

INTRODUCTION

Pachythecaliina Eliášová, 1976 (= *Amphiastraeina* Alloiteau, 1952) is an extinct suborder (Late Triassic–Late Cretaceous) that has focussed the attention of coral researchers in the context of the origin of corals from the order Scleractinia Bourne, 1900 in the Middle Triassic, and their possible relationship to the order Rugosa Milne-Edwards et Haime, 1851. Many pachythecaliines display “archaic”, skeletal features, which are unique among post-Palaeozoic corals. These corals usually were or still are classi-

fied in the Scleractinia. However, because of their skeletal architecture, especially in the Late Triassic Zardinophyllidae Montanaro Gallitelli, 1975 and the Jurassic–Cretaceous *Amphiastraeidae* Ogilvie, 1897 – similar to the late Palaeozoic plerophyllines – some authors classified them directly into the Rugosa (Koby, 1888; Ogilvie, 1897) or assumed their rugosan ancestry (Cuif, 1975, 1977, 1981, 2010; Melnikova and Roniewicz, 1976; Stolarski, 1996). Alternatively, pachythecaliines (in a narrow or broad meaning) were




Fig. 1. General location of study area on geological map (1: 500 000) (after Cheshitev *et al.*, 1989; modified and corrected) with location of sampling sites for corals, rudists *Mathesia darderi*, as well as ammonites, used for local biostratigraphy

classified as the separate order Hexanthiniaria Montanaro Gallitelli, 1975 (Montanaro Gallitelli, 1975; Eliášová, 1976b, 1978; Roniewicz, 2008; Melnikova and Roniewicz, 2012; Morycowa, 2012).

Phaceloid (branched, built of parallel corallites, pseudocolonial) pachythecaliine corals were moderately common in the Late Jurassic and locally diversified taxonomically (Tithonian–Lower Berriasian Štramberk Limestone), but were rare during the Cretaceous. The coral communities, dominated by diversified, phaceloid pachythecaliines, recently discovered in the Upper Barremian limestones of the Emen Formation in north central Bulgaria (Kołodziej *et al.*, 2009, 2011b), are unique among post-Jurassic coral communities. The aims of this paper are the taxonomic and palaeoecological analysis of these corals, as well as the interpretation of the sedimentary environment of pachythecaliine-rich bioconstructions in the Emen Formation.

GEOLOGICAL SETTING AND MATERIAL

During the Barremian–Early Aptian several carbonate platforms existed on the northern, Tethyan margin, located in the present territory of Bulgaria (Lovech Urganian Group, Vratsa Urganian Group, Russe Formation). In particular, the Lovech Urganian Group in the central Fore-Balkan (north central Bulgaria) contains diverse coral and rudist assemblages. This Urganian complex consists of four terrigenous and four carbonate formations (Figs 1, 2; Khrichev, 1966; Nikolov, 1969; Peybernès *et al.*, 1998; review in Minkovska *et al.*, 2002; Nikolov *et al.*, 2007). Coral communities – from level-bottom assemblages to coral-microbialite reefs – represent both clear- and turbid-water environments. Up to now, more than 100 coral species were described, mainly from soft marls, but were analysed more rarely with respect to their palaeoecology (e.g., Toula, 1889;




Fig. 2. Lithostratigraphic scheme of study area and stratigraphic position of sampling sites

Zlatarski, 1967, 1968a, b, c; Tchechmedjieva, 2001; Idakieva, 2002, 2003, 2007, 2008; Idakieva and Ivanov, 2002; Kołodziej *et al.*, 2007; see also Ilcheva and Motchurova-Dekova, 2011). Diversified coral assemblages occur also in pure or marly limestones (Kołodziej *et al.*, 2011c). However, apart from the present paper, they have not yet been the subject of detailed, taxonomic publications.

The pachythealiine corals studied occur in limestones of the Emen Formation in the Veliko Tarnovo (also transliterated as Veliko Turnovo, Veliko Trnovo, Veliko Tyrnovo) area. They were collected mainly in quarries at Rusalya and Vishovgrad. Sampling was supplemented in the Zarapovo-ecotrack and Hotnitsa-ecotrack, referred to below as Zarapovo and Hotnitsa, respectively (Fig. 1). This study also includes samples from Veliko Tarnovo, or its immediate vi-

cinity, referred to below as Veliko Tarnovo, collected by the late Prof. Ryszard Marcinowski (Warsaw University), but the exact collecting location is unknown.

The Emen Formation, situated in the middle part of the Lovech Urganian Group, was established by Khrishev (1966) in the region of Emen village, about 20 km NW of Veliko Tarnovo (Fig. 1). The best outcrops of this formation occur between the Osum and Jantra Rivers, and the type section is located at the Alexander Stambolijski Dam near to Gorsko Kosovo village, 40 km west of Veliko Tarnovo. Previous studies showed that the limestones of the Emen Formation were deposited on a shallow-water, carbonate platform, largely in an open lagoon. Intercalations of siliciclastic sediments are relatively rare (Khrishev, 1966; Khrishev and Bakalova, 1974). Microfacies analyses, performed by Min-




Fig. 3. General view of Rusalya Quarry. **A** – view of south wall, showing lowermost part of section with units 1, 2 and part of unit 3. Maximum height of wall is about 30 meters. **B** – view of WNW wall with approximate boundaries between units 3, 4 and 5. Straight line does not reflect exact relief of pachythealiine-*Mathesia* biostrome. Depression behind quarry is built of siliciclastics of Gorna Oryahovitsa Formation

kovska (1996) in three sections revealed sediments of (1) the inner platform (the Preobrazhenski Monastir section – close to Veliko Tarnovo; cored wells), (2) external parts of the inner platform (the area around Emen village), and (3) various facies, representing the distal to the inner part of the carbonate platform (Alexander Stambolijski Dam section). In the Emen Formation bioclastic and coral-bearing limestones dominate, while rudist occurrences are uncommon (Minkovska, 1996). The section in Rusalya was studied for the first time by the present authors (see Fenerci-Masse *et al.*, 2011).

In the study area, the Emen Formation grades laterally into the siliciclastic Gorna Oryahovitsa Formation (Figs 2, 3B). The Emen Formation consists here of two tongues bracketed by marly units with ammonites appearing to correspond to the upper part of the Lower Barremian (probably *Kotetishvilia compressissima* Zone) and the middle part of the Upper Barremian (*Gerhardtia sartousiana* Zone) (Khrishev, 1992; Ivanov, 1995; Ivanov and Nikolov, 1995; Stoykova and Ivanov, 2000; Ivanov and Idakieva, 2009; see also Fenerci-Masse *et al.*, 2011).

The stratigraphy of the sampling sites ranges from the early Late Barremian (*Toxancyloceras vandenheckii* Zone; site Hotnitsa) up to the early middle Late Barremian (*Gerhardtia sartousiana* Zone; sites: Rusalya and Vishovgrad Quarries, Zarpovo). The presence of the latest Early Barremian *Moutoniceras moutonianum* Zone in Hotnitsa cannot be ruled out.

More detailed studies of the sedimentary succession were performed in the Rusalya Quarry. The Vishovgrad Quarry was inaccessible for detailed studies. Microfacies studies were performed on thin sections from all sampling sites. The results, presented here, are based on the study of 136 thin sections; 94 thin sections are of standard size (4 × 2.7 mm) and 42 are large (6 × 5 mm). The specimens are deposited at the Institute of Geological Sciences, Jagiellonian University, Kraków (collection UJ 225 P). Twenty thin sections from three samples, collected by the late Prof. R. Marciniowski, are housed at the Institute of Palaeobiology, Polish Academy of Sciences, Warszawa (collection abbreviated as ZPAL Bułg).

SEDIMENTARY SUCCESSION AT RUSALYA QUARRY

As noted above, more detailed studies of the sedimentary succession bearing pachythealiine-rich biostromes were performed only in the Rusalya Quarry. In the following chapter the results of field observations and microscopic analysis of the material from all sites are presented.

Sedimentological studies at Rusalya were carried out on a well-exposed section, about 42 m thick (Fig. 3). The section consists of five main, lithological units (Fig. 4):

(1) ~10 m. Bioclastic packstone, rarely wackestones and grainstones, containing small, decimetre-scale boundstone patches, with irregular rims, built of small corals and calcified sponges, encrusted by *Lithocodium aggregatum* and microbial structures with vesicular, ‘bacinellid’ fabric (Fig. 4A, B). Colonial corals and their fragments are small (usually no more than a few centimetres in size). Associated biota include calcified sponges (chaetetids, stromatoporoids) and rare rudists. Pachythealiine corals (or other phaceloid corals), abundant in the coral-*Mathesia*-microbialite biostromes (unit 4), have not been recognized in unit 1.

(2) ~15 m. Bioclastic limestones interlayered with rudist (mostly monopleurids) limestones (Fig. 4C).

(3) ~8 m. Packstone-dominated, bioclastic limestones with chaetetids, stromatoporoids, subordinately with rudists and small, colonial corals, but without pachythealiines (Fig. 4D).

(4) ~2.5 m. Coral-rudist-microbialite biostromes and possibly also low-relief bioherms. The dominant metazoan components are large, phaceloid corals: pachythealiines (Fig. 4E, see also Figs 11–13, 17–19, 22–24) and *Calamophylliopsis* sp. and small monopleurid rudists *Mathesia darderi* (Astre, 1933), densely clustered locally (e.g., Fig. 4F; see also Figs 5C, 6A). Microbialites are common (e.g., Fig. 7B, C, E; see also Fig. 23B, C). Access to the upper part of the section was difficult, therefore establishing a more detailed pattern of the lateral distribution of corals and rudists, as well as the lateral extension and geometry of bioconstructions (biostromes, low-relief bioherms), requires supple-




Fig. 4. Generalized lithostratigraphic section of Upper Barremian succession at Rusalya Quarry, showing main lithologic units, with examples of biofacies. **A, B** – unit 1: bioclastic limestone with small boundstone patches (arrows in A), built of small corals, calcified sponges, *Lithocodium aggregatum* (B) and ‘bacinellid’, microbial structures. **C** – unit 2: bioclastic limestone with monopleurid rudists. **D** – unit 3: bioclastic limestone with chaetetid sponges. **E, F** – unit 4: coral-rudist-microbialite biostrome with phaceloid pachythealliine coral *Pleurophyllia* sp. (E) and monospecific, rudist cluster of *Mathesia darderi* (F). **G, H** – unit 5: rudist biostromes, built of dense aggregates of elongated, monopleurid rudists

mentary studies. Their detailed description, including microfascies analysis, is given in the next section.

(5) ~6 m. Rudist biostromes, built by dense aggregates of rudists, more diversified and larger in size than those in unit 4: large and small requieniids, elongated, monopleurids and possibly *Agriopleura* sp. (Fig. 4G, H; J.-P. Masse, pers. comm., 2012). Chaetetids occur subordinately.

PACHYTHECALIINE-RUDIST-MICROBIALITE BIOCONSTRUCTIONS

The simple term *biostrome* or *autobiostrome* (*sensu* Kershaw, 1994) can be used for the boundstones, analysed at Rusalya, because field observations indicate the bedded nature, although owing to difficult access to the upper part of the section at Rusalya, the occurrence of low-relief bioherms can not be excluded. The term *biostromal reef* is applied here, in accordance with the recent, broad approach to reef definition (Leinfelder *et al.*, 1994; Insalaco *et al.*, 1997; Wood, 1999; Stanley, 2001; Riding, 2002; Kiessling *et al.*, 2002; Kiessling, 2009; Flügel, 2010). The broad term *bioconstruction* represents a more inclusive term for more-or-less clearly recognizable structures (Höfling, 1997; Riding, 2002). This term is also used here, because determination of the exact morphology of reefal lithosomes (biostromes *versus* bioherms) was difficult, owing to the state of the outcrops at Zarapovo and Hotnitsa, and limited access for study in the Vishovgrad Quarry.

The descriptions of pachytheicaliine-rudists-microbialite bioconstructions are based on field observations and microscopic studies (including the taxonomy of corals and some other fossils) mainly of samples from Rusalya and Vishovgrad. Pachytheicaliines are clearly the most common and diversified corals in the bioconstructions studied (see Systematic Palaeontology, Figs 10–24): 14 species (six new, eight in open nomenclature), 12 genera (four new, three of them are distinguished formally). Large, phaceloid forms are often in growth position (Figs 4F, 5A, B). The most common (17 fragments of large coralla; 15 of them collected in Rusalya) is *Pleurophyllia bulgarica* sp. nov. reaching about 70 cm in height. In contrast to Rusalya, sampling at other sites was not extensive, which also reflected in the taxonomic diversity, namely at Rusalya: eight genera, nine species; at Vishovgrad: four genera, five species; at Hotnitsa: two genera, two species; at Zarapovo: one genus, one species. Analysis of 20 thin sections from three samples collected by the late Prof. R. Marcinowski at the unknown site at Veliko Tarnovo revealed three species of pachytheicaliines, *Calamophylliopsis* sp. and one undetermined coral, thus showing a general, “taxonomic pattern” as the rich coral assemblages from Rusalya. Associated corals, except for *Calamophylliopsis* sp. are rare. These corals are small, and only the phaceloid *Calamophylliopsis* sp. attains a larger size of up to 40 cm (Fig. 8).

Apart from corals, the monopleurid rudist *Mathesia darderi* and microbialite crusts provided additional, structural support for reef development. Rudist assemblage is nearly monospecific, dominated by *M. darderi*, with small, cylindrical shells (50% of the average diameters ranging

from 0.7 to 0.8 cm; Fenerci-Masse *et al.*, 2011) with the outer, calcitic shell layer and the inner one, aragonitic converted to neomorphic calcite (Fig. 6; Fenerci-Masse *et al.*, 2011, figs 5–7, 11; see also Masse and Fenerci-Masse, 2010). They frequently are clustered densely and occur between coral branches, or are attached directly to them (Figs 4F, 5C, 6, 12A, D, 14C, 16A; see also Fenerci-Masse *et al.*, 2011). Some small growth anomalies were observed in corals or rudists (Fig. 6C). Outside Rusalya, *M. darderi* is common at Vishovgrad and in samples from Hotnitsa (Fig. 5C). This species was also recognized in thin sections from samples, collected in Veliko Tarnovo by Prof. R. Marcinowski. They were not found at Zarapovo, but this may be due to sampling bias because, only limited sampling was performed at this site.

The corals and rudists are encrusted by microbialites, which occur also in semi-closed spaces (e.g., in interskeletal spaces between coral skeleton elements; Figs 7D, 23 A–B) and may partially (microbial “bridges”, e.g., Figs 14C, D, 18F) or completely fill space between skeletons. Microbialites show dense, non-laminated micritic microfabric (Figs 7A–E, 8A, B), thus at the scale of microstructure can be categorized as pure leiolites. Poorly marked microlamination (Fig. 18A) and micropeloidal (mostly within microcavities; Fig. 9C, D) or clotted microfabrics was observed only locally. The crusts are termed here *microbialite*, but the more inclusive term *automicrite* can also be applied (see Microbialites in the chapter Palaeoenvironmental interpretation).

Microbialites are “pure”, contain micrometre-scale “sparitic spots” of uncertain origin (small bioclasts?) or incorporate varying amounts of skeletal debris. In thin sections, microbialites are easy to distinguish from allomicrites by their dark colour (e.g., Figs 7A, D, 8A, B, 23B, C). Microbialites do not exhibit borings or encrusting microorganisms, either externally or internally, but fine burrowing traces are relatively common (Fig. 7B, C; compare with Fig. 9). Microbialite growth generated small cavities, locally with geopetal structures, filled with allochthonous micrite, calcite sparite cement, and rarely small grains (Figs 6A, 9). Bioclasts are rare in cavities, except for ostracods (Fig. 6E). Microbialite growth between skeletal elements (septa, wall, dissepiments) occurred only locally, “closed” interskeletal space, which prevented filling by sediment. Therefore, the space between skeletal elements is largely filled with calcite spar cement (e.g., Figs 5A, 10B, C, 12D, L, 13, 14C, 15D, 17A, 22B, C, H, 24B, C). Because septa are poorly preserved (usually only slightly micritized), observations of coral-lite morphology are difficult, particularly in vertical sections.

Small pyrite or orange-brown iron oxides (Fig. 7E), and locally also dolomite crystals are scattered within the microbialites, though very rarely within allomicrite in samples from Vishovgrad (Figs 8B, 15F). The surrounding sediment and biota appear unaffected by dolomitization, although in places the distinction between automicrite and allomicrite, if dolomitized, may be difficult. The presence of dolomite was supported by the standard staining of thin sections with Alizarin Red-S (see Adams *et al.*, 1984). In contrast to the calcite cement and allomicrite, dolomite remained unstained (Fig. 15F).

With regard to microfacies the pachytheicaliine-bearing limestones from all of the sampling sites are similar. The associated sediment is calcimudstone, bioclastic wackestone and packstone. Quartz was observed only within the tests of some agglutinated foraminifera. Most metazoan bioclasts are fragments of small rudists (Figs 6A, 7A, 12B, 248A, B). Large metazoans, apart of corals and rudists, include sponges (calcified and non-rigid; Figs 8, 22A, E), and complete shells or fragments of small gastropods, recognizable mostly in thin sections (Figs 12A, 17A, C, D). Sponge spicules occur locally within the microbialites (Fig. 9A, B). Sponges, both non-rigid and calcified, show differential preservation even in spatially close parts of the same specimen (Fig. 9C, D, E). Well preserved skeletons may pass laterally to micropeloidal aggregates (Fig. 9C, D). Some structures may be relicts of sponge spicules (Fig. 9G, H). Other biota includes echinoderm plates, decapod crustacean appendages, similar to *Carpathocancer? plassenensis* (Schlagintweit et Gawlick, 2002) (Fig. 22G; cf. Schlagintweit *et al.*, 2007), small brachiopods, benthic foraminifera, including encrusting/cryptic forms (Figs 7F, H, 9F) and a few orbitolinids, ostracods (relatively common, particularly in microcavities, Fig. 7D, E), encrusting bryozoans, serpulids (Fig. 22A, F, G), *Girvanella*-like tubes and rare algae. This latter group includes dasycladacean green algae: *Zittelina hispanica* Masse, Arias et Vilas, 1993 (Fig. 18E), *Neomeris cretacea* Steinmann, 1899 (Fig. 18F), but mostly *Terquemella* sp., that is reproductive structures of undetermined large algae (I. I. Bucur, pers. comm., 2010). Crusts of coralline red algae were recognized only in two samples at Zarapovo. In contrast to the lower part of the section at Rusalya (Fig. 4B), thin crusts of *Lithocodium aggregatum* (*sensu* Schlagintweit *et al.*, 2010) and microbial structures with “bacinellid” fabric (*sensu* Maurin *et al.*, 1985; Schlagintweit and Bover-Arnal, 2013) with poorly developed vesicular meshwork, are rare (except in samples from Zarapovo) in most samples with pachytheicaliines studied (Fig. 7F, G). Structures similar to and confused with *L. aggregatum*, recently reinterpreted as entobian borings (see section *Lithocodium/Bacinella* in the next chapter), associated with boring foraminifer *Troglotella incrustans* Wernli et Fookes, 1992 (Schmid and Leinfelder, 1996; Schlagintweit, 2012), were not recognized in the limestones studied. Large borings in metazoans, mostly *Entobia* sp., are moderate to sparse in abundance. Only two large bivalve borings were observed, cutting both metazoans and sediment.

PALAEOENVIRONMENTAL INTERPRETATION

General interpretation of sedimentary setting at Rusalya

Analysis of section at Rusalya, about 42 m thick, allows placement of biostromes discussed within a broader sedimentary and environmental context. The dominance of bioclastic packstones, the sparse rudists, the occurrence of small corals, the presence of encrustations of *L. aggregatum*, “bacinellid” structures and the subordinate role of microbialites indicate that the lower part of the section (unit 1)




Fig. 5. Field photographs of pachytheicaliine and *Mathesia*-bearing limestones from Vishovgrad Quarry (A, B) and Hotnitsaecotrack (C). **A** – *Pleurophyllia* sp. in growth position. Most coralites are in longitudinal section; note calcite spar cement filling most of interskeletal space, except uppermost part of coral branches. **B** – *Aulastrea* sp.; scale in millimeters. **C** – large massive colony of *Metaulastrea* sp. (*Met*) and densely clustered, small rudists *Mathesia darderi* (*Ma*)

represents the outer carbonate platform. Organisms and matrix sediment indicate a mildly mesotrophic environment and moderate hydrodynamics, but higher than during the development of the biostromes of unit 4 (see discussion below). Limestones, similar to those in unit 1, are common in the sedimentary succession of the Emen Formation at the Alexander Stambolijski Dam studied by Minkovska (1996) and by the present authors (unpublished data). Our studies of the section at the Alexander Stambolijski Dam revealed different facies, including limestones with corals, rudists, *L. aggregatum* and crusts with a “bacinellid” fabric. Pachythe-




Fig. 6. Small rudist *Mathesia darderi* from pachythealiine-rich boundstones. **A** – *M. darderi* aggregated in cluster and two branches of coral *Pleurophyllia* sp. **B** – rudist, attached to coral skeleton (longitudinal section). **C** – growth anomaly (arrow) of coral skeleton at boundary with rudist. **D** – juvenile rudist (arrow), closely attached to coral skeleton. Note two layers of rudist shell: inner layer originally aragonitic, and outer, calcitic layer. A – Vis 52/1, B – Ru 19-08/1, C – Ru, D – Vis 47/2

caliines, other branching corals and *Mathesia darderi* are absent there. Changes in lithology and biotic assemblages in units 2 and 3 and particularly the recurrent interlayers of rudist limestones, indicate changes in environmental parameters possibly controlled by changes of sea-level, but still remaining in the depositional setting of the outer carbonate platform.

It has been suggested previously that the pachythealiine and *Mathesia*-rich biostromes studied (unit 4) developed in the distal portion of the rudist-dominated part of the carbonate platform (Kołodziej *et al.*, 2009; Fenerci-Masse *et al.*, 2011). In the topmost part of the section, biostromes of unit 5 built by dense rudist assemblages, cover directly the pachythealiine-rich biostromes indicating an inner carbonate platform setting. Thus, the vertical, sedimentary and biotic succession in the section studied shows – with some fluctuations – a general shallowing trend, from the outer- to the inner-platform environment. Palaeoenvironmental inferences, based on an analysis of corals, rudists, microbialites and other biota from the pachythealiine-bearing bioconstructions, are discussed below.

Corals

The coral communities in the bioconstructions discussed are dominated by phaceloid forms: pachythealiines and less common *Calamophylliopsis* sp. In phaceloid corals, polyp tissue does not extend beyond the margin of the calice, so that such corals are considered pseudocolonies (Coates and Jackson, 1985; Roniewicz and Stolarski, 1999). Phaceloid corals were particularly well adapted to deal with high sedimentation rates in low-energy settings (e.g., Leinfelder *et al.*, 1994; Roniewicz and Stolarski, 1999; Dupraz and Strasser, 2002; Helm and Schülke, 2006). Geister (1995) calculated the growth rate of the Late Jurassic phaceloid *Aplosmilia* sp. as about 10 mm/year. However, a low rate of accumulation of allocthonous sediment is assumed for limestones discussed (but see discussion below on the

growth rate of microbialites). A low energy environment at the transition from the inner to outer platform may be inferred, on the basis of the abundant, branching corals in the growth position (even though microbialite crusts enhanced the rigidity of the corals) and matrix-supported background sediment (mostly bioclastic wackestone).

Phaceloid, epithecate corals, with simple polyp organization, were highly developed in the Late Triassic and Late Jurassic (Roniewicz and Stolarski, 1999, and references therein). Modern, phaceloid corals are rare and epithecate, solitary corals occur in deep-water and cryptic habitats. Since the Late Cretaceous the decline of epithecate corals (including phaceloid ones), and the proliferation of non-epithecate corals are observed. This evolutionary trend probably was mainly driven by increased bioerosion in coral environments (Roniewicz and Stolarski, 1999).

It is worth emphasizing that corals, representing *Calamophylliopsis* (relatively common in the limestones studied) were described in the literature from various Jurassic–Cretaceous sediments, which implies growth under a different set of conditions, including stressful ones, such as higher sediment and nutrient input (for the Jurassic, see Leinfelder *et al.*, 1994; and references to systematic papers in Roniewicz, 1976; Turnšek, 1997; Morycowa, 2012; for the Cretaceous, see citation lists in Löser *et al.*, 2002). In the Barremian–Albian of Romania, *Calamophylliopsis* is known from pure limestones containing diverse corals associated with phototrophic and oligotrophic microencrusts, from ‘*Lithocodium–Bacinella*’ facies, with poorly diversified corals, as well as from siliciclastic-dominated sequences (Kołodziej *et al.*, 2011a). It is also common in marls of the Lovech Urgonian Group (Idakieva and Ivanov, 2002; Idakieva, 2003).

During the Late Jurassic, the main period of development of phaceloid pachythealiines (see the next chapter, the section Spatial and temporal proliferation of pachythealiines), these corals (mostly amphistraeids) occurred in different types of reef structures. They are known from coral-microbialite reefs, with photophilic/oligotrophic micro-

encrusters (e.g., Eliášová, 1975, 1981; Insalaco *et al.*, 1997; Kołodziej, 2003), and from coral thickets without microbial and/or the crusts mentioned above (e.g., Roniewicz, 1966; Insalaco *et al.*, 1997). Of particular significance is the common occurrence of specimens of phaceloid *Pleurophyllia*. Among Late Jurassic assemblages it was one of the most common, amphistraeid corals, but except for in the Tithonian–Lower Berriasian Štramberg Limestone (see below), its record from the Lower Cretaceous is rare. In the list of citations on Cretaceous corals, Löser *et al.* (2002) mentioned only two species of *Pleurophyllia*, and sparse records worldwide.

A strictly actualistic approach to the palaeoecology of fossil corals cannot be applied, owing to evolutionary changes and the different, environmental preferences of many, modern reef corals (e.g., adaptation to high energy and an oligotrophic regime; see Wood, 1999). Coral communities from the biostromes studied are dominated by phaceloid forms. The question under debate is whether phaceloid corals were photosymbiotic or not. Some Late Triassic phaceloid corals (*Retiophyllia*, *Pachysolenia*) are considered to have been zooxanthellate, on the basis of stable isotope composition (Stanley and Swart, 1995). Studies of stable isotopes of the organic matrix from skeletons of the Upper Triassic solitary, pachythealiine coral *Pachythealis major*, recently performed by Muscatine *et al.* (2005), indicate that this species was photosymbiotic. The authors hypothesize that photosymbiosis may have played a role in scleractinian skeletogenesis, after the disappearance of the *Rugosa* in the Permian.

On the other hand, Recent, low-integrated or pseudocolonial corals are highly resistant to sedimentation and feed largely or entirely heterotrophically (Dryer and Logan, 1978, *vide* Sanders and Baron-Szabo, 2005 and Silvestri *et al.*, 2011). These observations have been used in the interpretation of fossil phaceloid corals. Kiessling *et al.* (2009) suggest that the occurrence of Early Jurassic corals from southern France, mostly phaceloid forms, within siliciclastic sediments may indicate, that these possibly were mostly azooxanthellate. Thus, the unresolved question of whether phaceloid corals were zooxanthellate or not represents obstacle to the attempt to unravel the palaeoenvironmental parameters, controlling growth of the corals, studied here. However, rapid calcification does not always correlate with a zooxanthellate status of corals (Marshall, 1996; Wood, 1999). Some branching azooxanthellate species, representing for example *Lophelia*, *Madrepora*, *Oculina* or *Tubastraea*, reveal a rapid growth rate, as much as 26 mm/year (Sabatier *et al.*, 2012, and references therein).

Phaceloid pachythealiines were common and highly diversified during sedimentation of the Tithonian–Early Berriasian Štramberg Limestone from the Czech Republic (17 genera, 35 species) and Štramberg-type limestones from Poland (see next chapter, section Spatial and temporal proliferation of pachythealiines). These reef limestones are characterized by the common occurrence of microbialites (typically laminated, with a micropeloidal microfabric), phototrophic microencrusts and diversified macrobiota (e.g., Morycowa, 1974; Eliášová, 1981; Eliáš and Eliášová, 1984; Kołodziej, 1997; Bucur *et al.*, 2005; for references

see also Vašíček and Skupien, 2004). However, the factors that governed prolific growth and diversification of pachythealiines during this time and at this part of the north Tethyan margin are unknown. Cuif and Stolarski (1999) have hypothesized that formation of the epithelial wall without septa (wall-based corals) in pachythealiines and Recent *Gyunia* might be an adaptation to a stressful environment. However, in contrast to pachythealiine corals from the Emen Formation, those from the Štramberg Limestone are associated with diversified corals and other biota, although short-term stressful events may be difficult to reveal.

Most of the Late Jurassic coral assemblages were described from the Upper Oxfordian and Kimmeridgian (Leinfelder *et al.*, 2002; Martin-Garin *et al.*, 2012). However, proliferation of pachythealiines in the Tithonian was not strictly time controlled. By comparison, among 42 species from coral-bearing limestones of the Carpathian Foreland, comparable in age, only one belongs to Pachythealiina (Morycowa, 2012). During that time, this area was located within a palaeolatitude position, similar to that of the carbonate platforms with sedimentation of the Štramberg Limestone. The possible reasons for environmental differences may be due to the fact that the former area was located slightly farther north, on the SW margin of the East European Craton, and was less affected by Late Jurassic/earliest Cretaceous tectonics. In another example, in the Kimmeridgian to Valanginian, biostromal reefs (similar to the Štramberg Limestone, with regard to lithofacies; Ivanova *et al.*, 2008) in SW Bulgaria, Roniewicz (2008) recognized diversified coral communities (50 genera, 72 species), but only four genera and five pachythealiine species.

Some cerioid/pseudocerioid amphistraeids, such as *Amphiastrea*, were opportunistic organisms. The Middle–Late Jurassic *Amphiastrea piriformis* Gregory, 1900 is known from various lithologies/environments, including turbid, siliciclastic environments, with fluctuating salinity. These corals were adapted to a high nutrient level and to the active removal of sediment (Fürsich *et al.*, 1994; Dupraz and Strasser, 2002). Similarly, cerioid/plocoid heterocoeniids are known from different sediments, including siliciclastics (e.g., Morycowa, 1964a; 1971; Beauvais, 1982; Morycowa *et al.*, 1994), implying broad, environmental adaptation.

By contrast to the Emen Formation in the study area, apart of the amphistraeid *Metaulastrea*, *Amphiastrea* and heterocoeniid *Latusastrea*, other pachythealiines are unknown within the diversified coral assemblages in marls and limestones of the Lovech Urganian Group and Vraca Urganian Group, as indicated by published results (see references in Geological setting and material), and the unpublished results of recent studies, including the analysis of a huge coral collection (coll. V. Zlatarski) housed at the National Museum of Natural History in Sofia.

Rudists

During the Cretaceous, corals coexisted with rudists in a range of environmental overlap. However, there is a general pattern in their distribution on the Urganian and other Cretaceous carbonate platforms. Rudist associations charac-




Fig. 7. Microbialites and some microfossils from pachytheicaliine-*Mathesia*-microbialite boundstones. **A** – ?*Paracarolastraea* sp. and geopetally filled growth cavity. Arrows indicate boundary between automicrite (*m*) and allochthonous micrite (*alm*); *r* indicates rudists and their fragments. **B** – microbialites, poorly laminated locally, developed on and between coral branches; arrows show burrowings or small growth cavities. **C** – small burrows (arrows) in microbialite, developed between coral branches. **D** – microbialite crusts on coral septa, mostly developed on their left side; arrows indicate ostracods. **E** – growth cavity within microbialites filled with ostracods and pyrite crystals. **F** – *Lithocodium aggregatum* (*La*), microbial crusts (*m*) and encrusting foraminifera (*f*). **G** – coral, encrusted by *Lithocodium aggregatum* (*La*). **H** – hyaline foraminifer, exhibiting fine spines (?*Ramulina* sp.) within microbialites. A – Vis 40/1, B – Ru 34/3, C – Ru 22/2, D – Vis 2/3, E – Ru 14-08/2, F – Ru 34/3, G – Ru 30-09, H – Ru 9-10

terized different settings, depending on morphotype, but occurred mainly in the inner carbonate platform environment; bioclastic limestones are referable to the outer platform, while coral facies to the transition between the inner and outer platform. The biotic distribution pattern was largely controlled by changes in environmental conditions, forced by external factors, such as water turbidity, nutrient level and hydrodynamic regime (e.g., Masse and Philip, 1981; Gili *et al.*, 1995a, b; Skelton *et al.*, 1997). In the mixed, car-

bonate-siliciclastic Urgonian, system in Bulgaria, rudists (poorly known in contrast to corals) may co-occur with corals, but usually they occur in separate lithosomes. Moreover, rudists are absent in the marls, while corals may be highly diversified there.

Apart from the limestones discussed, *Mathesia darderi* is unknown from other deposits of the Urgonian complex in Bulgaria. *M. darderi* is included in the rudist family Monopleuridae that, in general, played a limited role on the Early

Cretaceous carbonate platforms of the Mediterranean region. Locally, however, monopleurids were abundant and may have a significant, constructional potential, especially during the Early Valanginian and Late Aptian–Albian, with *M. darderi* as a key species. Relatively thick lithosomes (up to 3 m) are usually formed by dense, monospecific assemblages (Fenerci-Masse, 2006 *vide* Masse and Fenerci-Masse, 2010; Fenerci-Masse *et al.*, 2011; Skelton and Gili, 2012).

The monospecific genus *Mathesia* is known from the Late Barremian to Middle Albian. The first Late Barremian occurrences have recently been recognized in Bulgaria and Spain. During the Late Barremian to Early Aptian, this genus was present only locally and later was widespread (Fenerci-Masse *et al.*, 2011) escaping the mid-Aptian crisis, when 90% of species and 70% of the Mediterranean rudist genera disappeared (Masse, 1989; Skelton and Gili, 2012). The ecological changes, recorded in *Mathesia* through time, reflect a displacement from the distal, rudist-dominated part to the proximal part of carbonate platforms (Fenerci-Masse *et al.*, 2011). In the Albian of southern Spain, *M. darderi* build dense, monospecific assemblages, found interbedded with intertidal stromatolites and muddy sediments (Fenerci-Masse, 2006; see Fenerci-Masse *et al.*, 2011). In SE Spain, *M. darderi* occurs within the uppermost Barremian rudist-rich (Requieniidae, Monopleuridae) packstones and wackestones, with dasycladalean algae, above marls and dasyclad-rich limestones and below lower Aptian orbitolinid-rich marly limestones with quartz (Fenerci-Masse *et al.*, 2011). Thus, with regard to the sedimentary context, including biofacies (lack of corals), this occurrence differs from the *Mathesia*-bearing limestones of the Emen Formation in Bulgaria.

Rudists that are in contact with corals were attached to the skeleton of phaceloid corals, which, unlike branching ramose forms (like recent *Acropora*), were not covered by living tissue (except for the tips of branches). Growth anomalies, observed both in corals and in rudists, are very rare, and suggest rare, direct tissue contact (*in vivo* interaction). However, some kind of synecological relationship between rudists and corals is plausible. Synecological interactions in coral-rudist associations, such as the one from the Campanian of Spain (Götz, 2003), have been described only rarely in the literature. Protuberances of rudist shells were explained by Götz (2003) as possible defence reactions of the rudist against the coral cnidia. However, both groups might have benefited from this coexistence. Rudists might provide hard substrates for coral settlement, while rudists were stabilised by encrusting corals (Götz, 2003). A similar relationship may be assumed for the coral-rudist association studied, even though indications for *in vivo* association are only hypothetical.

Physical erosion during the growth of corals and rudists was rather negligible, because of the inferred, low energy level. Common, small rudist fragments are rather a result of intrinsic (shell structure/mineralogy), than extrinsic (hydrodynamics, bioerosion), taphonomical factors. Fragmentation of rudists, due to their composition, that is aragonite (originally) inner and calcitic outer shell layers, may be an important, syndimentary modifier of the original rudist bioconcretion (Sanders, 1999).




Fig. 8. *Calamophylloopsis* sp. Arrows indicate geopetally filled growth cavities or small burrows. ZPAL Bulg 1/1

Other macrobiota

In general, apart from pachytheicaliine corals and *M. darderi*, macrobiota and their remains are not common in the bioconstructions studied. Therefore, small gastropods are of special interest, as they are common components of the bioconcretion described. Taxonomy and feeding behaviour of these gastropods are unknown. They were possibly grazers or sediment feeders, basing on assumed abundant, organic matter in the microbialites. An increased nutrient level may also enhance primary production of fleshy algae, favouring herbivorous gastropods (see Dupraz and Strasser, 2002).

Microbialites

A characteristic feature of the bioconstructions analysed are dense, nearly always non-laminated, micritic crusts, occurring on and between skeletal metazoans, as well as within semi-closed microcavities. Their genesis may have inferences for the environmental interpretation of the depositional environment. These crusts are called here *microbialites*, the term used in a broad meaning, i.e. as a result of organomineralization *s.l.*, defined as microbially-induced and microbially-influenced mineralization (Dupraz *et al.*, 2009). Automicrite is a more inclusive term and can be applied both for 'classical', microbial fabrics, as well as for microcrystalline carbonates, which originated in association with non-living organic macromolecules, that is related to organomineralization *s.s.* (Trichet and Défarge, 1995). Some authors propose use of the term *automicrite* instead of *microbialite*, if the origin of autochthonous micrite is unknown (e.g., Reitner and Neuweiler, 1995; Bourque, 1997; Neuweiler *et al.*, 1999; Webb, 2001; Schlager, 2003). Automicrite that resulted from organomineralization *s.s.* was recognized by Neuweiler *et al.* (1999) in exceptionally well preserved, organic fractions in the Albian carbonate mud mounds of Spain. According to these authors, the Late Jurassic represents an important period of organomineralization *s.s.* and resulted in the precipitation of automicrite, which is frequently referred to as micro-

bial crusts, common in coral- and sponge-microbialite reefs (e.g., Leinfelder *et al.*, 1993, 1994; Leinfelder, 2001; Dupraz and Strasser, 2002; Olivier *et al.*, 2004; Matyszkiewicz *et al.*, 2012; Pleš *et al.*, 2013). Discrimination between these different processes and products in fossil material is a challenge for future research (Riding, 2000).

Attribution of microbialites from the Emen Formation to one of the major categories of stromatolites, thrombolites, dendrolites, or leiolites can not be strictly applicable here. These categories are based on macrofabrics, namely laminated, clotted, dendritic and aphanitic respectively (Kennard and James 1986; Schmid, 1996; Riding, 2000), which are not recognizable macroscopically in the material studied. These main categories are also used in the classification of microbialites, based on a combination of macrostructure and microstructure (Schmid, 1996; Flügel, 2010, fig. 9.1B). In such an approach, microbialite crusts in the biostromes studied can be classified largely as leiolite microbialites, i.e., microbialites with dense microstructure.

The presence, mostly on coral branches, of differentially preserved non-rigid and calcified sponges raises the question of possible contribution of non-rigid sponges to microbialite formation. Sponge spicules, occurring locally within microbialites, reticulate microfabric, similar to sponge structures recognized in Triassic sponge-microbial stromatolites (Szulc, 1997, 2000, fig. 21g), as well as a lateral transition from well preserved sponge to micritic/micropeloidal microfabric, indicate that some microbialites might be the result of sponge soft-tissue diagenesis. Microbially induced carbonate precipitation (microbialite formation) inside decaying Recent and fossil sponge tissue is well documented (e.g., Reitner, 1993; Reitner and Neuweiler, 1995; Delecat and Reitner, 2005; Reolid, 2007). Increased alkalinity induces dissolution of siliceous spicules and may be responsible for the lack of microscleres in some facies (Delecat and Reitner, 2005).

Foraminifera, occurring within microbialites may be an additional argument in the discussion of the possible contribution of sponges to microbialite formation. They are similar to those reported from fossil and modern sponge meshworks (e.g., Guilbault *et al.*, 2006; Reolid, 2007) or inhabiting cavities (e.g., Helm, 2005; Helm and Schülke, 2006, fig. 111; Schlagintweit and Velić, 2012, fig. 7g, h). Alternatively, these foraminifera may be interpreted as loosely attached to microbialite crusts. Automicrite occurs locally between coral skeletal elements, thus indicating development of microbialites also in aphotic conditions (compare Albian microbialites: Neuweiler, 1993, 1995). Apart from the calcification of decaying sponges (sponge 'container automicrites'), these organisms, as well as the soft tissue of other metazoans may be primary sources of organic material (Reitner and Neuweiler, 1995; Neuweiler *et al.* 1999).

Micropeloids, which are rare in the limestones studied, may be similar in origin to the dense microbialite crusts. Many authors regard peloids as *in situ* growth products, related to randomly distributed nucleation centers, or as a result of bacterial degradation of organic matter (e.g., Reid, 1987; Neuweiler, 1993; Riding and Tomás, 2006, and references therein).

Bryozoans, serpulids, rare *L. aggregatum* and 'bacinellid' crusts, sporadically red algae and juvenile rudists di-

rectly encrust corals or rudists. Except some foraminifera of uncertain mode of attachment and generic status, discussed above, microbialites are never encrusted by microencrusters. That suggests that they were only partially lithified and hard substrate was not available for larval settlement. Such an interpretation is supported by the lack of borings in microbialites (except for rare, large bivalve borings in metazoans, microbialites and sediment matrix). Similarly, a lack of metazoan encrusters on microbialites was observed by Webb (1999) in Carboniferous patch reefs, and by Neuweiler (1993) in Albian microbialites. According to Webb (1999), this implies that some automicrites (= microbialites *s.l.*) have not been lithified at the sediment-water interface, but were formed as synsedimentary precipitates within sediment. The presence of burrows also indicates that the studied microbialites were not rigid, but only semi-consolidated, when burrowing took place (cf. Riding, 2000; Riding and Tomás, 2006). In contrast to these examples, intergrowths of skeletal microencrusters and microbialites were recorded for example in the Late Jurassic reefs, indicating the presence of a hard substrate and interruptions of microbialite growth (Olivier *et al.*, 2003). Growth cavities in the material studied are small; their presence is marked by geopetal fillings and cryptic ostracods (cf. Aubrecht *et al.*, 2002; Shen and Webb, 2005).

In Recent reefs, enhanced microbialite growth is characteristic for environments with higher nutrient levels and elevated alkalinity (Camoin and Montaggioni, 1994; Camoin *et al.*, 1999, 2006; Sprachta *et al.*, 2001), which is also accepted for fossil coral reefs (e.g., Leinfelder *et al.*, 1994; Dupraz and Strasser, 2002; Olivier *et al.*, 2004). Sedimentation of allochthonous deposits and microbialite growth rate, in relation to the growth of metazoan constructors, is crucial for the development of a constrictal or superstratal growth fabric, which has architectural, palaeoecological, sedimentological and diagenetic implications (Insalaco, 1998). Microbialite growth is favoured by a low accumulation rate, but the growth rate is difficult to estimate, because there are no modern analogues for fossil, reefal microbialites. Microbialites from cryptic caves of Lizard Island display very low net growth rates of 10 to 15 mm/1000 years (Reitner, 1993). However, in shallow-water reefs, where corals are closely intergrown with microbialites, they might have grown as fast as the corals, about 1–2 mm/year, as postulated by some authors (see Schmid, 1996; Schmid *et al.*, 2001). Seard *et al.* (2011) observed microbialite growth rates, ranging from 7 to 12 mm/year. Such unusually rapid growth rates probably result from their development during a period of environmental change (sea-level rise), while in more stable environments, they are significantly lower (Seard *et al.*, 2011; Heindel *et al.*, 2012).

As discussed above, phaceloid corals were well adapted to high sedimentation rates, soft, muddy substrates and low-hydrodynamic regimes. A low background sedimentation rate is assumed for the biostrome formation analysed. However, the growth of phaceloid corals also might be favoured by a higher growth rate of microbialites. However, it was not a dominant influence on the development of phaceloid corals, because except for *Calamophylliopsis* (sub-order Faviina), only phaceloid pachythecaliines were recog-




Fig. 9. Differential preservation of sponges from pachytheicaliine-*Mathesia*-microbialite boundstones **A, B** – sponge spicules (s) within dark-coloured microbialites (m); alm allochthonous micrite, filling small growth cavities; d (in B) indicate dolomite crystals within microbialites (compare Fig. 15F). **C–E** – differential preservation of rigid (calcified) sponges. Micropeloids in C and D, possibly resulted from degradation of sponge tissue. Note micropeloids in intraskeletal space (arrow in D). **F** – non-rigid sponge, encrusted by nubeculariid foraminifera (n) and microbialites with *Ramulina*-like foraminifera (R). **G** – reticulate microfabric, and **H** – needle-like structures, possibly resulted from dissolution of sponge spicules. A – Vis 2/3, B – Vis 2, C, D – Ru 7257/1, E – Ru 34/3, F – Vis 47/2, G, H – Zar 996/3

nized. If coral accretion occurred at a rate, similar to that of microbialite growth, then limited, positive relief, that is, a constrictal coral growth fabric (*sensu* Insalaco, 1998) may be inferred.

Dolomite crystals, recognized in the samples from Vis-hovgrad, possibly are restricted to microbialites, which indicates the syngenetic nature of dolomite formation and a possible link with microbial activity or organomineralization *s.s.* (cf. Wright and Wacey, 2005; Mastandrea *et al.*, 2006; Bontognali *et al.*, 2010).

At elevated nutrient levels, bioerosion generally is considered to be enhanced (Hallock, 1988; and Sanders and Baron-Szabo, 2005 for review). However, increased bioerosion is not observed in the material studied. Macroboring in metazoan skeletons are moderate or scarce, and absent in microbialites. A higher nutrient level, constant or episodic, raises the question about water oxygenation. Episodes of low oxygen concentrations are generally invoked to explain microbialite development in Late Jurassic reefs (Leinfelder *et al.*, 1994; Betzler *et al.*, 2007). However, in the material

studied here, the fauna, although not highly diversified, does not support the concept of a poorly oxygenated bottom. Moreover, some Jurassic microbialites could have formed in oxic waters (Olivier and Boyet, 2006; Olivier *et al.*, 2011).

Lithocodium/Bacinella

Lithocodium aggregatum and associated, microbial structures, with a ‘bacinellid’ fabric (*Bacinella irregularis*) are an important issue in palaeoenvironmental interpretation of Upper Jurassic and Lower Cretaceous shallow water limestones. Difficulties in the interpretation of depositional environment are due to their unclear, systematic position and environmental demands. *Lithocodium aggregatum* Elliot, 1956 was variously interpreted, among others as algae, calcimicrobes or encrusting foraminifera, while *Bacinella irregularis* Radoičić, 1959 was mostly interpreted as microbial structures. Some authors assumed that both taxa represent different parts of one organism (for review see Schmid and Leinfelder, 1996; Rameil *et al.*, 2010; Schlagintweit *et al.*, 2010; Schlagintweit and Bover-Arnal, 2012, 2013). Recently, new interpretations have been proposed for taxa, traditionally labelled as *L. aggregatum* and *B. irregularis*. Schlagintweit *et al.* (2010) and Schlagintweit and Bover-Arnal (2013), based on studies of Aptian material, interpreted *L. aggregatum* as ulvophycean green alga with heterotrichale encrusting thallus, and *B. irregularis* as an euendolithic chlorophycean alga. Most of structures described as *B. irregularis* are, according to these authors, vesicular, microbial crusts with “bacinellid” fabrics. *L. aggregatum* may develop also a cryptic stage with net-like structure (Schlagintweit and Bover-Arnal, 2012) which can be misinterpreted with “bacinellid” structures. Most of the structures described as *B. irregularis* are, according to these authors, vesicular, microbial crusts with “bacinellid” fabrics. *L. aggregatum* may develop also a cryptic stage with a net-like structure (Schlagintweit and Bover-Arnal, 2012), which can be misinterpreted as “bacinellid” structures. For a summarizing diagram, showing different, morphological and taxonomic interpretations, see Huck *et al.* (2012, fig. 8). Further interpretative complications arise, because many crusts (in particular in the Upper Triassic and Upper Jurassic rocks), determined in the literature as *L. aggregatum*, are in fact sponge borings in microbialites or other carbonate substrates (Cherchi and Schroeder, 2010, 2013; Schlagintweit, 2010).

Occurrences of *L. aggregatum* s.s., *Lithocodium*-like entobian borings and “bacinellid” structures in the Late Jurassic–earliest Cretaceous, coral-microbialite reefs are commonly interpreted as an indicator of shallow-water, although data from the literature indicate that *L. aggregatum* s.s. was adapted also to outer-shelf/ramp environments (see discussion in Bover-Arnal *et al.*, 2011; Schlagintweit and Bover-Arnal, 2012; Huck *et al.*, 2012). These microencrusters are also assumed to be an indicator of oligotrophic or mildly mesotrophic, reefal and lagoonal environments (e.g., Leinfelder *et al.*, 1993; Dupraz and Strasser, 2002; Olivier *et al.*, 2004; Ivanova *et al.*, 2008). However such an interpretation seems to be appropriate, only when

they are associated with other microencrusters and occur as subordinate components in Late Jurassic–Early Cretaceous coral reefs and lagoonal facies (e.g., Rameil *et al.*, 2010). During the mid-Cretaceous time, in particular in the Early Aptian, a local “bloom” of so-called “*Lithocodium-Bacinella*” facies (a descriptive term) occurred (Immenhauser *et al.*, 2005; Huck *et al.*, 2010). It is assumed that the Aptian “*Lithocodium-Bacinella*” facies show a more complex, biotic pattern, controlled by fluctuating nutrient levels, sea-water alkalinity, elevated seawater temperatures, seawater acidification and low sedimentation rates, superimposed on global perturbations of neritic ecosystems (Neuweiler and Reitner, 1992; Immenhauser *et al.*, 2005; Huck *et al.*, 2010; Rameil *et al.*, 2010; Bover-Arnal *et al.*, 2011; Huck *et al.*, 2012).

In the pachythecalline-bearing limestones studied, corals and other metazoans are rarely encrusted by *L. aggregatum* and “bacinellid” structures, but instead they are mostly encrusted directly by microbialites or by heterotrophic microorganisms and small rudists, followed by microbial crusts. The studies of Late Jurassic coral reefs revealed that the lack of phototrophic-dominated organisms in the first layer of encrustation indicates more turbid waters and/or a higher nutrient level (Leinfelder *et al.*, 1993; Dupraz and Strasser, 2002; Olivier *et al.*, 2004; Helm and Schülke, 2006). Similar, environmental controls can be assumed for the Early Cretaceous coral reefs, which in contrast to the Late Jurassic reefs, are studied less extensively in this respect. Even though the taxonomic interpretation of some structures, described as *L. aggregatum* and *B. irregularis* (= “bacinellid” structures), have been recently re-interpreted, the environmental inferences appear to be still valid, when these organisms occur as a moderate component of coral reefs, and not as a dominant bioconstructor, as in some Aptian occurrences. However, as concluded by Schlagintweit *et al.* (2010, p. 541), “...the palaeoenvironmental significance of *Lithocodium aggregatum* occurrences must be carefully viewed in each case study”.

Concluding remarks

The unique association of pachythecaliines and *Mathesia* forming biostromes occurs within a palaeogeographically limited area. The resulting bioconstructions (biostromes and possibly also low-relief bioherms), unknown elsewhere, imply particular environmental factors driving their growth and microbialite formation. *Lithocodium* and ‘bacinellid’ structures are moderately common in unit 1 at Rusalya. Obviously the shift from the outer carbonate platform (unit 1) towards the more inner platform setting (unit 4) has led to the change of environmental factors, such as lowering of the hydrodynamic regime and possibly increasing nutrient level. These changes favoured growth of phaceloid pachythecaliine corals, nearly monospecific rudist assemblage (*Mathesia darderi*), facilitated growth of microbialites, but limited growth of *L. aggregatum* and “bacinellid” structures. Difficulties in deciphering environmental constraints are, among other factors, due to the lack of comparable, fossil biocoenoses. *Mathesia* up to now was known from deposits lacking, or with rare corals. Phaceloid pachy-

thecaliines were rare in the Early Cretaceous, whereas Late Jurassic taxa were relatively common, but occurred in various, sedimentary settings, as indicated by host lithology and associated biota. By comparison with the corals studied, highly diversified, phaceloid pachythecaliines, known from the Tithonian–Lower Berriasian Štramberk Limestone and Štramberk-type limestones (Czech Republic, Poland), are associated with diversified corals and other biota, including phototrophic/oligotrophic microorganisms.

In the best exposed section at Rusalya, the pachythecaliine-rich biostromes attain only a thickness of about 2.5 m. Thus, it is tempting to view development of this biofacies during a short time interval and as a potential, local, stratigraphic marker. However, stratigraphic data from Hotnitsa and limited sampling in Hotnitsa and Zarpovo do not permit unambiguous, stratigraphic implications. However, development of this biofacies appears to have occurred largely during the middle Late Barremian.

GENERAL REMARKS ON MORPHOLOGY, SYSTEMATICS AND DISTRIBUTION OF PACHYTHECALIINA

Morphology and microstructure

The suborder Pachythecaliina Eliášová, 1976, in particular the Late Triassic family Zardinophyllidae Montanaro Gallitelli, 1975 (= Pachythecaliidae Cuif, 1975), and the Jurassic–Cretaceous Amphistraeidae Alloiteau, 1952, contains skeletal features non-typical for scleractinian corals: a thick wall (pachytheca), and septa, commonly deep in the calice and arranged in bilateral symmetry (quasi-radial in the adult stage of some taxa). Pachythecallines, except most heterocoeniids, are solitary or their skeleton indicates polyps that originally were poorly integrated. They are mostly of the phaceloid, cerioid or pseudocerioid growth form. Amphistraeids, the most common pachythecaliines (Roniewicz and Stolarski, 2001; Stolarski and Russo, 2001), reproduced asexually, by unique Taschenknospung ('pocket'-budding). Other families, classified into Pachythecaliina, share only some features with their supposed relatives, the zardinophyllids (Montanaro Gallitelli, 1975; Cuif, 1975, 1981) and amphistraeids (Eliášová, 1975, 1976b, 1978; Melnikova and Roniewicz, 1976; Kołodziej, 1995, 2003; Roniewicz and Stolarski, 2001; Stolarski and Russo, 2001; see also Amphistraeina and Heterocoeniina, generally characterized in Baron-Szabo, 2002; Löser, 2009; and <http://www.coralosphere.org>).

The skeleton microstructure has been debated focussing on trabecular *versus* non-trabecular microstructure of septa and wall. Most post-Triassic pachythecaliines have badly preserved skeletons and diagnostic microstructural characters are not well recognized. The dominant view is that the wall (pachytheca) is built by horizontal modules with circumferential calcification centers (Cuif, 1975; Roniewicz and Stolarski, 2001; Stolarski, 2003; Cuif, 2010). Change in the style of biomineralization, from horizontal modules, having circumferential calcification centres, to a strictly trabecular one (vertical trabeculae, with axially arranged calci-

fication centers) was observed in the Upper Triassic amphistraeid *Quenstedtiphyllia fritschi* (Volz, 1896) (see Roniewicz and Stolarski, 2001). Septal microstructures of pachythecaliines *s.s.* were considered both as non-trabecular (Cuif, 1975, 2010; see also Stolarski, 2003), and minitrabecular (see discussion in Kołodziej, 1995; Roniewicz and Stolarski, 2001). Roniewicz *et al.* (2007, p. 593) defined the microstructure of Triassic pachythecaliines as "Wall epithelial, thick and modular in structure; septa thin and built of centripetally growing trabeculae, or septal microstructure fibronormal (in relation to the midseptal plane); septal faces not ornamented". However, if we take into account also pachythecaliines *s.l.* (*sensu* Stolarski and Russo, 2001), not all post-Triassic pachythecaliines, particularly heterocoeniids do fit with such definition. Wall can be thin (e.g., *Donacosmilia*), and in contrast to the zardinophyllids, even in amphistraeids, septal faces can show ornamentation (e.g., Fig. 16D, E). Differences in microstructures between amphistraeids and heterocoeniids are discussed below. Moreover, traditional terminology dealing with skeletal microstructure (thick-trabecular, minitrabecular; Roniewicz and Morycowa, 1993), possibly requires modification, owing to the new model of skeletal coral growth, proposed by Stolarski (2003).

The microstructure of the wall and septa in post-Triassic pachythecaliines has been interpreted variously. In Jurassic–Cretaceous amphistraeids, the wall is built up of "...well arranged modules of the shape of horizontal spines" (Roniewicz and Stolarski, 1999, pp. 144–145; see also Ogilvie, 1897; Morycowa, 1964b; Eliášová, 1975; Kołodziej, 1995; Roniewicz and Stolarski, 2001). The pachytheca – a wall diagnostic for Pachythecaliina – implies that it is built by fibrous microstructure. In fact, in most pachythecaliines, the microstructure of the septa and wall is unknown, owing to the poor state of preservation. Moreover, differentiation in diagenetic alterations of the wall in amphistraeids and related families might suggest an original variability in microstructure (Eliášová, 1976b; Roniewicz and Stolarski, 1999; Kołodziej, 2003). A wall in pachythecaliines (including heterocoeniids) is commonly termed as "wall developed in advance to septa" (Cuif, 1975, 2010; Cuif and Stolarski, 1999; Roniewicz and Stolarski, 2001; Stolarski and Russo, 2001) or "wall developed prior to septa" (see Kołodziej, 1995).

In particular, there is a controversy, concerning the microstructure of heterocoeniid skeletons, which was interpreted in different ways. Septa have been described as built by large trabeculae (Morycowa, 1971; Roniewicz and Morycowa, 1993) or minitrabeculae (Kołodziej, 1995). It is worth emphasizing that septa in heterocoeniids (even the primary septum) can be thinner than the lower limit of thick trabeculae. The diameter of thick trabeculae *sensu* Roniewicz and Morycowa (1993, p. 235) is "...over 50 μm (usually more than 100 μm) to ca. 300 μm ". Moreover, there are indications suggesting that the modular structures of amphistraeid *Amphiastrea* and heterocoeniid *Latusastrea* are similar (Kołodziej, 1995). Important difference between these two groups is a trabecular(?) microstructure of a peritheca, a skeletal structure, occurring in some heterocoeniids, but unknown in other pachythecaliines. Relics of

more diverse, than usually thought, septal and wall microstructures were observed in some pachythecaliines *s.l.* (Eliášová, 1976b; Kołodziej, 2003, figs 3, 4).

Classification and phylogeny

Following Roniewicz and Stolarski (2001), some authors distinguish the suborder Pachythecaliina instead of the suborder Amphistraeina (Stolarski and Roniewicz, 2001; Stolarski and Russo, 2001; Kołodziej, 2003; Roniewicz, 2008; Melnikova and Roniewicz, 2012; Morycowa, 2012). Others still accept the priority of Amphistraeina. However, the possible relationships of Triassic pachythecaliines with amphistraeids and related families are seldom discussed by these authors (e.g., Baron-Szabo, 2002, 2006; Turnšek *et al.*, 2003; Löser, 2008c, 2012; Löser *et al.*, 2009).

Usually the following families are included in the Pachythecaliina (= Amphistraeina): (1) the Late Triassic Zardinophyllidae (recent finding also in the Lower Jurassic; Melnikova, 1975; Melnikova and Roniewicz, 2012), and Late Triassic–Cretaceous (2) Amphistraeidae [two amphistraeid genera *Quenstedtiphyllia* Melnikova, 1976 and *Sichuanophyllia* Deng Zhanqiu et Zhang Yansheng, 1984 (= *Lubowastraea* Melnikova, 1986) are known from the Upper Triassic], and Jurassic–Cretaceous (3) Carolastraeidae Eliášová, 1975, (4) Donacosmiliidae Krasnov, 1970, and (5) Intersmiliidae Melnikova et Roniewicz, 1975. More problematic is the systematic position of the family Heterocoeniidae Oppenheim, 1930, which is usually classified in the separate suborder Heterocoeniina Beauvais, 1977. Previously heterocoeniids were classified in different suborders; some heterocoeniids were classified within the Amphistraeidae (Vaughan and Wells, 1943; Wells, 1956; see Kołodziej, 1995). L. Beauvais (1974, 1976) believed that there are similarities (also in microstructure) between Amphistraeidae and Heterocoeniidae and later L. Beauvais (1981) classified Amphistraeidae together with Heterocoeniidae in a new suborder Distichophylliina, a suborder not recognized in current coral classifications. By contrast, M. Beauvais (1977, 1982) included Heterocoeniidae in the new suborder Heterocoeniina. Eliášová (1976b) accepted both Amphistraeina and Heterocoeniina. However, she classified them together with the new suborder Carolastraeina (not recognized in current coral classifications), and Triassic Pachythecaliina in the separate order Hexanthiniaria (see below). Hence, she assumed morphological similarities and phylogenetic relationships for these coral groups. Kołodziej (1995) proposed to include Heterocoeniidae (superfamily Heterocoenioidae) in Amphistraeina, and later into Pachythecaliina (Kołodziej, 2003), a view, which is also held in the present account. Stolarski and Russo (2001) considered heterocoeniids as “supposed pachythecaliines”. Baron-Szabo first classified heterocoeniids into Heterocoeniina (e.g., Baron-Szabo and Steuber, 1996; Baron-Szabo, 1998), later into Amphistraeina (Baron-Szabo, 2002, 2006). Taxonomical difficulties, concerning heterocoeniids, are well exemplified by *Latusastrea* Orbigny, 1849, a genus, which between 1940–1990 was classified within five suborders (Turnšek and Löser, 1989). The suborder Heterocoeniina was accepted recently in scleractinian systematics by Morycowa and Marcopoulou-Diacantoni

(2002), Idakieva (2003), Morycowa and Decrouez (2006), Roniewicz (2008), and by Löser (e.g., 2008a, b, c, 2009, 2010).

Co-occurrence (in the limestones studied) of pachythecaliines *s.s.* (*sensu* Stolarski and Russo, 2001), that is, amphistraeids (five genera, six species), and pachythecaliines *s.l.*, that is, carolastraeids (one genus, two species), intersmiliids (one genus, one species), donacosmiliids (one genus, one species), and heterocoeniids (four genera, four species), indicates similar environmental preferences. The occurrence of four heterocoeniid genera, displaying the phaceloid growth form, which is rare in this family, should be emphasized. The co-occurrence with other four pachythecaline families is not conclusive evidence of phylogenetic relationships; however, it is one more piece of evidence, supporting such an interpretation. A similar co-occurrence of five pachythecaline families is known from the Tithonian–Lower Berriasian Štramberg Limestone (Czech Republic) and its equivalent in the Polish Outer Carpathians. Only heterocoeniids are rare there, but in general they were rare and poorly diversified in the Jurassic and the earliest Cretaceous. Certainly, taxonomic variability of pachythecaliines in the limestones studied is higher, as some of findings were not described here, owing to a poor state of preservation or not enough, available thin sections. Further sampling in the active quarry at Rusalya should provide more data on these unique anthozoans during their Cretaceous acme.

Taschenknospung, a peculiar type of budding, is only well recognized in amphistraeids. Possibly it occurs also in some heterocoeniids, as indicated by the relationship of parent and daughter corallites in *Thecidiosmilia morycowae* (see Kołodziej, 1995). On the other hand, strongly developed, lateral, septal ornamentation or even septal outgrowths are an important feature of heterocoeniids, but only for some genera, while others can have septa with poor ornamentation. In *Heterosmilia* gen. nov., a heterocoeniid genus established in this paper, only the primary septum has strong, septal outgrowths, while other septa lack ornamentation. Strong, septal outgrowths are not common in scleractinians, but occur also in other coral groups (e.g., Morycowa, 1971; Stolarski *et al.*, 2004), including Rhipidogyrina, which have true, aphophysal septa (Eliášová, 1973).

In summary, intersmiliids, carolastraeids, donacosmiliids and heterocoeniids are “...much easier clustered with pachythecaliines than with other coeval scleractinians” (Stolarski and Russo, 2001, p. 253), even taking into account microstructural differences. In particular, some heterocoeniid genera are morphologically closer to amphistraeids than to end members of a spectrum of Heterocoeniidae. Heterocoeniids include some colonial genera with high integration level, which distinguishes them from other pachythecaliines. However, progressive trends of increased integration are observed in several groups of reef-building, modular invertebrates, including corals (e.g., Coates and Oliver, 1973; Coates and Jackson, 1985; Roniewicz and Stolarski, 1999; Wood, 1999). Some of the diagnostic characteristics for pachythecaliines might have been lost in some lineages (see Stolarski and Russo, 2001). A good example is bilateral symmetry of the septal apparatus. Such a septal pattern is characteristic for amphistraeids. However, *Quenstedtiphyllia* Melnikova (subfamily Quenstedtiphylliinae) has quasi-ra-

dial symmetry in the adult stage (Roniewicz and Stolarski, 2001). Recently, Löser (2012) has distinguished a new, amphistraeid genus *Hexamphistrea*, similar to *Metaulastraea* (= *Amphistrea*), but showing radial symmetry (see also *Amphistrea suprema* in Morycowa and Marcopoulou-Diacantoni, 1997, 2002). Differences in morphology of the septa between the new genus and other amphistraeids, as well as the lack of information about Taschenknospung budding, raise a question about the systematic position of *Hexamphistrea*. Still, it is morphologically closest to amphistraeids than to other corals.

Even assuming differences in the microstructure between different pachytheicaliine groups, the possibility of phylogenetic relationships is not excluded (cf. Roniewicz and Stolarski, 2001, p. 36), because evolution of the skeleton microstructure of pachytheicaliines also should be taken into account. Therefore, it seems reasonable to consider a combination of features, when phylogenetic relationships are studied. Molecular data on modern corals support the usefulness of skeleton microstructure and morphology in classifications at the family level, but on the other hand, found their grouping into suborders to be debatable (e.g., Romano and Palumbi, 1996; see also Stolarski and Roniewicz, 2001). It is particularly important, if we consider classifying pachytheicaliines within a separate order (see below).

Position of Pachytheicaliina within Anthozoa

The other debatable question is the position of pachytheicaliines within the class Anthozoa and their possible relationship to Palaeozoic corals. These corals were usually, or still are, classified within the order Scleractinia. However, recognition of the unique, skeletal features of pachytheicaliines led to their also being considered as rugosan corals (Koby, 1888; Ogilvie, 1897) or survivors of the Rugosa (Alloiteau, 1957; Cuif, 1975, 1977, 1981, 2010; Melnikova and Roniewicz 1976; Stolarski, 1996; Cuif and Stolarski, 1999). Montanaro Gallitelli (1975) distinguished the new order Hexanthiniaria, intermediate between the Rugosa (plerophyllines) and the order Scleractinia, and classified here a new Upper Triassic family Zardinophyllidae. Later Eliášová (1976b) included in the Hexanthiniaria in the suborder Pachytheicaliina (with the family Pachytheicalidae Cuif, 1975, a younger synonym of Zardinophyllidae), Amphistraeina (Pachytheicaliina *sensu* Roniewicz and Stolarski, 2001), Carolastraeina and Heterocoeniina. Recently, this systematic position of Pachytheicaliina was accepted by Roniewicz (2008), Morycowa (2012) and Melnikova and Roniewicz (2012). It is also accepted in the present paper. However, the concept of Pachytheicaliina should be re-examined with regard to its diagnostic features, both in terms of skeletal morphology, septal pattern and significance of the microstructure as diagnostic feature of the pachytheicaliines. The broad range of septal pattern in pachytheicaliines, from (1) “rugosan”-like in *Zardinophyllum*, and (2) corallites with few or only one recognizable septum (*Pachythe-cophyllia*, *Monoaulastraea*; see Kołodziej, 2003; Löser *et al.*, 2009) to (3) scleractinian-like septal pattern as in the adult stage of *Carolastraea* and *Paracarolastraea* at the end of the spectrum is a challenge in establishing the diagnostic

criteria for pachytheicaliines. However, as discussed above, some “typical” pachytheicaliine skeletal features could have been lost in some lineages.

Further research is needed, particularly in dealing with septal insertion. Montanaro Gallitelli (1974, 1975) and Stolarski (1996) observed in *Zardinophyllum* a rather broad, intraspecific variability in metaseptal insertion, however, without a typical rugosan, septal pattern. Interestingly, the drawings, presented by Lebanidze (1991), show septal insertion in amphistraeid *Mitrodendron ogilviae* Geyer, 1955 similar to the one in the Rugosa, but it was never verified and documented in photographs. Stolarski (1996) hypothesized that some rugosans might have survived the P/T extinction in refuges and realized their potential, with modifications to their skeleton mineralogy and septal insertion. Thus some Permian, scleractiniamorphs might be the ancestors of some Triassic corals (see references above), whereas other scleractinian lineages might have evolved from soft-bodied (corallimorpharian-like) ancestors, as is commonly assumed (e.g., Stanley, 2003). The discovery of Late Cretaceous, solitary corals with an original calcitic skeleton indicates that some scleractinian corals indeed may secrete skeletons of different carbonate polymorphs (Stolarski *et al.*, 2007). The long lasting discussion on Triassic corals, representing the family Zardinophyllidae (= Pachytheicaliidae), and their supposed relationship to some Late Palaeozoic corals is still not expressed in the commonly accepted, higher-rank classification. Recently, Cuif (2010) summarized some ideas on the possible relationships between some scleractinians and rugosans and called for a re-examination of the present concept of the Scleractinia. The hypothesis on the phylogenetic relationships between the skeletal Palaeozoic corals and corals from the suborder Pachytheicaliina/Amphistraeina (or in general Scleractinia) was challenged by researchers of rugosans and other Palaeozoic corals, including Permian scleractiniamorphs (Oliver, 1980a, b, 1996; Fedorowski, 1997; Scrutton, 1997; Ezaki, 2004).

Spatial and temporal proliferation of phaceloid pachytheicaliines

The first pachytheicaliines – the Zardinophyllidae and two genera from the Amphistraeidae – are known from the Upper Triassic (Montanaro Gallitelli, 1974, 1975; Cuif, 1975; Roniewicz and Stolarski, 2001). Since the Late Jurassic, pachytheicaliines became more common, but still were poorly diversified. Usually only some species occur in Oxfordian–Kimmeridgian coral assemblages (e.g., Roniewicz, 1966: two genera, two species; Roniewicz, 1976: five genera, seven species). However, they are highly diversified and abundant (and largely phaceloid) in the Tithonian–Lower Berriasian Štramberg Limestone in Czech Republic, Outer Carpathians (17 genera, 35 species; Ogilvie, 1897; Geyer, 1955; Eliášová, 1974, 1975, 1976a, b, 1978) and in pebbles/boulders (so called exotics) of the Štramberg-type limestones from Poland (14 genera, 22 species; Ogilvie, 1897; Geyer, 1955; Morycowa, 1964b, 1974; Kołodziej, 1995, 2003). Owing to the displaced character of these limestones (huge olistoliths in Štramberg, pebbles and boul-

ders in the Polish Carpathians), the precise age of particular coral specimens/taxa from these limestones cannot be specified. On the basis of calpionellids, a Tithonian–Early Berriasian age is commonly accepted for the coral-bearing limestones (Morycowa, 1968, 1988; Houša, 1990; Ciborowski and Kołodziej, 2001), but it is possible that reefs were developed mainly during the Late Tithonian. Locally sedimentation of Štramberg-type limestones in the Polish segment of northern part of Tethys (mostly in lagoonal facies) continued until the Valanginian (see Morycowa, 1988; Ivanova and Kołodziej, 2010). In most papers on corals from the Štramberg Limestone, the age was determined as Tithonian (see Vašíček and Skupien, 2004 for references). For the reasons mentioned above, corals from the Štramberg Limestone are usually recorded in the literature and databases as Jurassic, not Cretaceous fauna. However, for the discussion of evolutionary trends it is important that until the Valanginian, coral assemblages have a “Late Jurassic character” (Roniewicz and Morycowa, 1993), recently exemplified by a detailed study of corals from the continuous, Kimmeridgian–Valanginian succession in SW Bulgaria (Roniewicz, 2008).

During the Barremian and Aptian, a time span with the highest coral development in the Early Cretaceous (Löser, 1998), phaceloid pachythecaliines were rare. Only a widespread proliferation of heterocoeniids, mostly with a cerioid-plocoid type of colony, is observed in the Cretaceous (see citation lists in Löser *et al.*, 2002). The richest Early Cretaceous assemblage of pachythecaliines, except for heterocoeniids, was described by Baron-Szabo and Steuber (1996) from the Aptian of Greece. They recognized 10 genera and 13 species (including two heterocoeniid genera and two species), but only two genera and three species are phaceloid.

In summary, a prolific development of phaceloid pachythecaliines in the Tithonian and Berriasian, possibly mainly in the Late Tithonian (Czech Republic, Poland), in the Late Barremian (Bulgaria), and to a smaller extent in the Aptian (Greece), occurred in palaeogeographically restricted areas, with poorly known, environmental constraints.

SYSTEMATIC PALAEOLOGY

(by Bogusław Kołodziej)

Abbreviations: UJ 225 P – x 1 – x refers to the sampling site (Ru – Rusalya, Vis – Vishovgrad, Zar – Zarapovo, Hot – Hotnitsa) and the specimen number from the collection UJ 225 P (Institute of Geological Sciences, Jagiellonian University, Kraków); UJ 225 P – x/1 – numbers 1, 2, 3 *etc.* refer to the thin section number from a given sample; ZPAL – Bułg 1/1 – number of the sample (Bułg 1/x) and thin section (Bułg 1/1), housed at the Institute of Palaeobiology, Polish Academy of Sciences, Warszawa.

Measurements used in descriptions are abbreviated as follows: **d** – corallite diameter (in mm); **c–c** – distance between corallite axes (in mm); **s** – number of septa; **S1... Sn** – septa of succeeding size orders; **number of the sample in bold** – samples presented on figures; () – less frequent values are given in brackets.

Order HEXANTHINIARIA Montanaro-Gallitelli, 1975
Suborder PACHYTHECALIINA Eliášová, 1976
Family AMPHIASTRAEIDAE Ogilvie, 1897
Subfamily AMPHIASTRAEINAE Ogilvie, 1897

Remarks: Roniewicz and Stolarski (2001) established the new subfamily Quenstedtiphyllinae containing the genus *Quenstedtiphyllia* Melnikova, 1975 from the Upper Triassic. This subfamily differs from the subfamily Amphiastracinae by one-zonal (tabular) endotheca and corallites with quasi-radial symmetry in the adult stage. In the case of poorly preserved specimens, skeletal structures in longitudinal section (e.g., type of endotheca) are difficult to decipher. Recently, Löser (2012) established the genus *Hexamphiastrea* with the type species *Amphiastrea suprema* Morycowa et Marcopoulou-Diacantoni (Morycowa and Marcopoulou-Diacantoni, 1997, 2002). *Hexamphiastrea* is close to *Metaulaestrea* Dietrich and *Amphiastrea* Étallon but has hexamerous symmetry and thick rhopaloid septa. However, owing to the septal morphology of *Hexamphiastrea*, its classification in the Amphiastracidae is under question.

Genus *Metaulaestrea* Dietrich, 1926

Type species: *Aulaestrea pompeckji* Dietrich, 1926

Remarks: Löser (2008c) replaced *Amphiastrea* Geyer, 1955 with *Metaulaestrea*. *Metaulaestrea* was provisionally established by Dietrich (1926), but it is valid, according to the International Code of Zoological Nomenclature (Ride *et al.*, 1999; see Löser, 2008c). According to Löser (2008c), the type specimen of the type species (*Aulaestrea conferta* Ogilvie, 1897), selected by Geyer (1955), does not represent *Amphiastrea*. From the Cretaceous, seven species of *Metaulaestrea* (= *Amphiastrea*) are known (see citation lists in Löser *et al.*, 2002). Thus, apart from *Amphiastrea* (about ten species), it is the most diversified genus among the Cretaceous amphiastreaeids.

Metaulaestrea cf. *rarauensis* (Morycowa, 1971)

Fig. 10

Material: Six samples, six thin sections.

Measurements: Morphometric measurements are presented in Table 1.

Table 1

Dimensions (in mm) of *Metaulaestrea* cf. *rarauensis* (Morycowa, 1971)

Sample number UJ 225 P -	d	c–c	s
Vis 5	4.5 × 7–10	6–9	up to 45
Vis 10	6.7 × 7–11	(5) 6–8 (9)	up to 50
Vis 26	6.9 × 9–12	7–10	?
Vis 44	4.6 × 6–9	5–7	up to 43
Ru 3-10	4.5 × 5–7	5–6	up to 40
Ru 14	5–11	5–7	up to 35

Remarks: The samples studied are determined in open nomenclature as *Metaulaestrea* cf. *rarauensis* (Morycowa). *M. rarauensis*, described from Lower Aptian of Romania (Morycowa, 1971), was also recognized from the Valanginian to Aptian of Greece, Ukraine, Turkmenistan and Azerbaidjan, but mostly was not illustrated (see Löser *et al.*, 2002). New species, established by Baron-Szabo (Baron-Szabo and Steuber, 1996) as *Amphiastrea keuppi*, is considered as a younger synonym of *M. rarauensis*. This species is also known from Barremian–Lower Aptian marls of the Lovech




Fig. 10 *Metaulastrea cf. rarauensis* (Morycowa). **A, B** – corallites with well developed marginarium. **C** – corallite with poorly developed marginarium. UJ 225 P; A – Vis 10/1, B – Vis 26/1, C – Vis 44/1

Urgonian Group (Idakieva, 2003). The peripheral part of the calicular space (containing shortest septa) in the samples studied is commonly filled with calcite spar cement and the septa are recrystallized. As a result, the exact counting of septa is difficult. Differences in measurements revealed in the specimens seem to reflect intraspecific variability, rather than the presence of more species. However, more thin sections are required to obtain detailed measurements.

Distribution: Lower part of middle Upper Barremian (Rusalya, Vishovgrad).

Genus *Amphiastrea* Etallon, 1859

Type species: *Amphiastrea basaltiformis* Etallon, 1859

Amphiastrea sp.

Fig. 11

Material: One sample (UJ 225 P – Ru 9-08), one thin section.

Measurements (in mm): $d = 6-7$ (8), $c-c = 5-7$, $s =$ up to 23.

Remarks: Locally, the minute interstices between adjacent corallites are filled with sediment. These interstices may be due to a poor state of preservation or may reflect subcerioid (with closely packed *basaltiformes* corallites). *Amphiastrea*, usually described as a cerioid coral, displays in fact a pseudocerioid growth form, with each corallite having its own wall (Morycowa and Lefeld, 1966; Melnikova and Roniewicz, 1976; Roniewicz and Stolarski, 1999, 2001). Such a growth form can be recognized, only if the lateral surfaces of corallites can be observed as covered by epitheca (see Ogilvie, 1897; Geyer, 1955; Morycowa, 1964b).

Distribution: Lower part of middle Upper Barremian (Rusalya).

Genus *Pleurophyllia* de Fromentel, 1856

Type species: *Pleurophyllia trichotoma* de Fromentel, 1856

Remarks: *Pleurophyllia* is rarely reported from the Cretaceous, whereas in the Late Jurassic it was a relatively common genus among amphiastreaeids. Löser *et al.* (2002) cited only a few records of this genus in the Cretaceous and only two species, *P. trichotoma* de Fromentel and *P. skuviensis* Turnšek. According to Roniewicz (2008), *P. skuviensis* described by Turnšek (in Turnšek and Mihajlović, 1981) from the Barremian–Lower Aptian of Serbia represents the genus *Cladophyllia* Milne Edwards et Haime, 1851. *P. trichotoma*, described by Sikhuralidze (1979) from Georgia, is possibly from the Upper Jurassic, not the Albian. Other Lower Cretaceous specimens are determined in open nomenclature, but they are too poorly documented to confirm their taxonomic determinations. In the limestones studied, *Pleurophyllia bulgarica* sp. nov. is the most common coral.

Although a phaceloid corallum is characteristic for this genus, a densely packed to subcerioid growth form may occur locally in




Fig. 11. *Amphiastrea* sp. Arrow indicates inter-corallite interstice filled with sediment. UJ 225 P – Ru 9-08/1

the specimens studied. This is in accordance with the recent redescription of the syntype of the type species *Pleurophyllia trichotoma* by Lathuilière (2012a): “Probably phaceloid corallum (the polished syntype shows on one side a single corallite and on the other face, in correspondence, a set of three corallites not yet separated)”. It is worth emphasizing that a mixed growth form of corallum occurs in some other pachythecaliine taxa. A dendroid-phaceloid to subcerioid growth form occurs in amphiastreaeids *Aulastrea* Ogilvie, 1897, *Hykeliphyllum* Eliášová, 1975 (e.g., Geyer, 1955; Eliášová, 1975), and in *Pleuroaulastrea variabilis* gen. et sp. nov., described below, and in particular in *?Pleurophyllia* sp. These examples give evidence of the high potential of some pachythecaliines for variability of growth form, which might be favoured by low, biological integration in these corals. According to Coates and Oliver (1973), who discussed rugosan corals, cerioid corallum is only a compact, phaceloid corallum. Many Recent and fossil corals show a response of growth forms to environmental factors (e.g., Young, 1999a and literature therein). However, mixed, cerioid-phaceloid growth forms are not common in scleractinian corals (Lathuilière, 1989, 1996). A changing growth pattern may have no systematic significance, but may be interpreted as a partial death of the colony, due to sedimentation or predation (Lathuilière, 1989), or reflects only close packing, which resulted from space compaction. Mixed growth forms of corals occur more commonly in the Rugosa. Partly phaceloid/partly cerioid coralla occur in rugosan Lithostrotionidae (e.g., Nudds, 1979) and Phillipsastreaeidae (e.g., Wrzolek, 2007). Some rugosan corals show changes in colony type – from cerioid to fasciculate (branching) – during regeneration (Fedorowski, 1980; Poty, 1981; Young, 1999b).

Pleurophyllia bulgarica new species, Kołodziej

Fig. 12

Holotype: UJ 225 P – Ru 14-08.**Paratypes:** UJ 225 P – Ru 19-08, Ru, 21, Ru 30, Ru 7257, Ru 8837, Ru 8860.**Etymology:** *bulgarica* – from the country name Bulgaria.**Type locality:** Rusalya, Bulgaria.**Type level:** Lower part of middle Upper Barremian, Emen Formation, Bulgaria.**Stratigraphic distribution:** Lower part of middle Upper Barremian.**Diagnosis:** *Pleurophyllia* of corallite diameters ranging from (4) 5 to 9 (12) mm and number of septa up to 49.**Material:** 17 samples, 30 thin sections (13 large), one polished slab.**Measurements:** Morphometric measurements are presented in Table 2.**Table 2**Dimensions (in mm) of *Pleurophyllia bulgarica* sp. nov.

Sample number UJ 225 P -	d	s
Holotype Ru 14-08	(5) 6–7 × 6–8 (11)	up to 32
Paratype Ru 19-08	(4) 5–8 × 7–9 (10)	up to 38
Paratype Ru 21	(4) 5–7 × 6–9	up to 32
Paratype Ru 30	4–6 × 4–8 (9)	up to 39
Paratype Ru 7257	(4) 5–6 (7) × 6–9 (12)	up to 49
Paratype Ru 8837	(4) 5–7 × 7–9	up to 41
Paratype Ru 8860	4–7 × 6–8 (9)	up to 38
Ru 1-10	4–8	?
Ru 2	(4) 5–7 × (6) 7–10	up to 34
Ru 4-08	5–6, longest up to 10	up to 32
Ru 7-08	4–6 × 5–10	up to 32
Ru 13-08	5–9 × (5) 8–12	up to 41
Ru 15-08	5–8 × 6–10	up to 40
Ru 17-08	4–8 × 6–10	up to 32
Ru 249	5–10	up to 35
Vis 1673	6–11	up to 34
Hot 1	6–10	up to 35

Description: Phaceloid corallum. Locally, corallites are close to each other, resulting in a subcerioid growth form. Corallites in transverse section are more or less elongated, rarely round. Septa arranged in bilateral symmetry, defined by longer septa on one side of corallite, but with the primary septum usually only slightly longer and thicker than others. Septa smooth and thin, although thick septa may occur in some corallites. Lonsdaleoid septa present. Endotheca two-zonal, built by large tabuloid dissepiments in the central part and vesicular dissepiments in the narrow peripheral zone. Marginarium in corallites rarely observed. Discontinuous “exfoliation” of wall recognized in one corallite. Taschenknospung budding.

Remarks: As in many other species, described in the present paper, differences in the number of septa depend partly on the state of preservation. At the peripheral part of corallites, septa are difficult to count also, because they are poorly and irregularly developed there (possibly also as septal spines like in heterocoeniids; cf. Morycowa, 1971; Kołodziej, 1995). *Pleurophyllia bulgarica* sp. nov. differs from *P. trichotoma* (known from the Upper Jurassic and Berriasian) in larger corallite diameters and more abundant septa. For example, *P. trichotoma*, described by Roniewicz (1966): d 6–8, s 20 (24); Eliášová (1975): d 5–7 (8.5), s 20 + S3; Kołodziej (2003): d (5) 6–8, s 20–30. Other known species of *Pleurophyllia* have smaller diameters and smaller numbers of septa. *P. trichotoma* shows bilateral symmetry, defined by a well developed, primary septum, which in the new species is poorly developed, although bilateral symmetry is well marked. On the other hand, a “main sector”, as in *Hykeliphyllum* Eliášová and *Pseudopistophyllum* Geyer does not occur. A skeletal structure, resulting from “exfoliation” (poorly preserved marginarium?) of a wall, similar to the one observed in *Paracarolastraea zlatarskii* gen. et sp. nov., was recognized only in one corallite (Fig. 12K). The greatest differences are in the specimen from Hotnitsa, where the “main sector” is more clearly developed (Fig. 12F, G).

Distribution: Lower and middle parts of Upper Barremian (Hotnitsa, Rusalya, Vishovgrad).

?Pleurophyllia sp.

Fig. 13

Material: One sample (UJ 225 P – Ru 7262), one large thin section.

Measurements (in mm): d = (4) 5–7 (8), s = up to 31.

Remarks: The coral shows mixed phaceloid (*Pleurophyllia*-like) to subcerioid (*Amphiastrea*-like) growth forms. The small piece of the sample does not permit determination of whether it is a fragment of a pseudoceroid-dominated or phaceloid-dominated corallum. Corallites are particularly poorly preserved in the phaceloid part. The interskeletal space is filled here with sparite cement, thus analysis of the septal apparatus is not possible, but the marginarium can be recognized in some corallites. The relation between corallites of the subcerioid and phaceloid parts of the corallum excludes the possibility of fusion of two different corals. As described above, the subcerioid growth form may locally occur in *Pleurophyllia bulgarica* sp. nov. and in the type species of *Pleurophyllia*.


Distribution: Lower part of middle Upper Barremian (Rusalya).

Genus *Aulastrea* Ogilvie, 1897Type species: *Aulastrea schäferi* Ogilvie, 1897*?Aulastrea toulis* new species, Kołodziej

Fig. 14

Holotype: UJ 225 P – Vis 2.**Etymology:** *touli* – in honour of Franz Toula, 19th Austrian coral researcher of Cretaceous corals in Bulgaria.**Type locality:** Vishovgrad, Bulgaria.**Type level:** Lower part of middle Upper Barremian, Emen Formation, Bulgaria.

Fig. 12. *Pleurophyllia bulgarica* sp. nov. **A–K** – transverse sections, **L** – longitudinal section. **A, B, D** – *r* indicates small rudists, mostly *Mathesia darderi* (Astre), particularly numerous in D. **B** – densely packed corallites. **C** – two corallites with septa of variable thickness. **F, G** – corallites with quasi main sector. **H, K** – corallites with ‘exfoliation’ of wall (compare Fig. 17). **J** – corallites with Taschenknospung budding. UJ 225 P; **A** – Ru 14-08/2, **B** – Ru 8860/1, **C** – Ru 8837/1, **D** – Ru 7257/3, **E, I** – Ru 30/1, **F, G** – Hot 1/2, **H, J** – Ru 19-08/2, **K** – Ru 4-08/1, **L** – Ru 249/1. **A** – holotype, **B, C, E, H, I, J** – paratypes






Fig. 13. ?*Pleurophyllia* sp., displaying, mixed subcerioid (A) to phaceloid growth form (B). Arrows indicate corallite, linking subcerioid and phaceloid parts of corallum. UJ 225 P – Ru 7262/1

Stratigraphic distribution: Lower part of middle Upper Barremian.

Diagnosis: Dominantly phaceloid, locally phaceloid-dendroid to subcerioid corallum. Corallites from about 7 mm to 21 mm in diameters, and up to 50 septa.

Material: One sample (UJ 225 P – Vis 2), four large thin sections, three polished slabs.

Measurements (in mm): $d = 7\text{--}21$, $s =$ highly variable, up to at least 50.

Description: Phaceloid, locally a phaceloid-dendroid to subcerioid corallum. Locally corallites form compact clusters. Corallites oval. Septa arranged in bilateral symmetry. Septa on one side of corallite are longer, but a primary septum is only slightly longer and slightly thicker. Lonsdaleoid septa present. Complete marginarium developed around the septal apparatus only in some corallites; in others, marginarium is poorly developed (“marginal pockets”). Endotheca in peripheral zone built of vesicular dissepiments; in central part poorly preserved, but possibly built of tabuloid dissepiments. Taschenknospung budding.

Remarks: The new species is tentatively assigned to the genus *Aulastrea*. It differs from *Aulastrea* in its growth form. *Aulastrea* has a massive subcerioid to dendroid-phaceloid corallum, with short branches and a conical lower part (see Ogilvie, 1897; Geyer, 1955; Eliášová, 1975; Schäfer and Senowbari-Daryan, 1980; Lathuilière, 2012b). In contrast to *Aulastrea*, there is a lack of a distinct, thick, primary septum. The septal pattern is similar to the one, occurring in *Pleurophyllia bulgarica* sp. nov. *Aulastrea* is poorly known from the Lower Cretaceous (see citation lists in Löser *et al.*, 2002). The best documented in the Cretaceous are *Aulastrea schaeferi* Ogilvie, 1897 and *Aulastrea* cf. *macer* Eliášová, 1975 described by Baron-Szabo (in Baron-Szabo and Steuber, 1996) from the Aptian of Greece, previously known only from the Upper Jurassic/earliest Cretaceous.

Distribution: Lower part of middle Upper Barremian (Vishovgrad).

Genus *Pleuroaulastrea* new genus, Kołodziej
Type species: *Pleuroaulastrea variabilis* sp. nov.,
Kołodziej

Etymology: In relation to *Pleurophyllia* and *Aulastrea*.

Diagnosis: Corallum phaceloid, locally subcerioid. Corallites round or slightly oval, and highly variable in diameter. Septa arranged in bilateral symmetry, defined by long axial primary sep-

tum. Lateral septal faces covered by small granules. Marginarium present. Lonsdaleoid septa present. Endotheca unknown. Taschenknospung and parricidal budding.

Stratigraphic distribution: Lower part of middle Upper Barremian.

Remarks: *Pleuroaulastrea* gen. nov. differs from *Aulastrea* in the general corallum growth form (see remarks on ?*Aulastrea touli* sp. nov.). In the new genus, the marginarium is poorly developed and not complete. Corallites are highly variable in size. The septal pattern in small corallites is close to *Pleurophyllia*. Apart from Taschenknospung, parricidal budding occurs, resulting in a partial, subcerioid growth form, built of small polygonal corallites. Figure 16A, B shows recrystallized corallites with parricidal budding. This budding type is similar to the one, occurring in *Mitrodendron* Quenstedt, 1880 (Roniewicz, 1966, text-fig. 15; pl. 16, fig. 1c; Melnikova and Roniewicz, 1976, fig. 4; Roniewicz, 2008, fig. 3c), in which new buds appear on the endothecal elements. It differs from the parricidal budding in *Intersmilia* Eliášová, 1974, where septa of a parent individual continue into a daughter one (Melnikova and Roniewicz, 1976, fig. 4; see also remarks on *Intersmilia* aff. *diaboli* Eliášová, 1974 in the present paper).

Pleuroaulastrea variabilis new species, Kołodziej
Figs 15, 16

Holotype: UJ 225 P – Vis 1.

Etymology: *variabilis* – for its high variability in corallite diameter and corallum growth form.

Type locality: Vishovgrad, Bulgaria.

Type level: Lower part of middle Upper Barremian, Emen Formation, Bulgaria.

Stratigraphic distribution: Lower part of middle Upper Barremian.

Diagnosis: *Pleuroaulastrea* with highly variable corallite size, from 4–8 mm in the subcerioid/densely phaceloid part of the corallum to 10–27 mm in the phaceloid part. Number of septa relatively small, up to 29.

Material: One sample (UJ 225 P – Vis 1), seven thin sections (two large), eight polished slabs.

Measurements (in mm): $d = 4\text{--}8$ in subcerioid/densely phaceloid part of corallum 10–27 in phaceloid part of corallum, $s =$ up to 29 (in large corallites).

Description: Phaceloid corallum. Locally corallites, resulting from parricidal budding, are close to each other, resulting in sub-




Fig. 14. ?*Aulastrea touli* sp. nov., holotype UJ 225 P – Vis 2. **A** – polished slab of holotype specimen. Corallites strongly recrystallized; interskeletal space commonly filled with calcite spar cement, in particular in peripheral parts of corallites. **B** – dendro-phaceloid to subcerioid part of corallum with Taschenknospung budding (Vis 2/2). **C, D** – corallites with poorly developed marginarium; arrows indicate small rudists (*r*) and small microbialite ‘bridges’ (*m*) between coral branches (C – Vis 2/2, D – Vis 2/1). **E** – corallite with broad marginarium (Vis 2/2). **F** – longitudinal section showing large, vesicular dissepiments at the peripheral part of endotheca. At left side, transverse to slightly oblique section displays small parricidal bud (arrow) (Vis 2/4)

cerioid growth form. Corallites round or slightly oval. Diameter of corallites is highly variable in diameter; they are significantly smaller in the subcerioid part of corallum. Septa arranged in bilateral symmetry, defined by long, axial, primary septum. Lateral faces of larger septa are covered by small granules. Marginarium well or poorly developed. Lonsdaleoid septa present. Endotheca unknown. Taschenknospung and parricidal budding.

Distribution: Lower part of middle Upper Barremian (Vishovgrad).

Family CAROLA STRAEIDAE Eliášová, 1976

Remarks: Until now Carolastraeidae contained only one genus *Carolastraea* Eliášová, 1976 with three species. Löser (2009) included this genus in the Amphistraeidae.

Genus *Paracarolastraea* new genus, Kołodziej

Type species: *Paracarolastraea zlatarskii* sp. nov., Kołodziej

Etymology: *Paracarolastraea* – for similarity to *Carolastraea* Eliášová, 1976.




Fig. 15. *Pleuroaulastrea variabilis* gen. et sp. nov, holotype UJ 225 P – Vis 1. **A–C** – polished slabs of holotype specimen, showing high variability in corallite arrangement and size. **B** – polished slab, showing (black arrow) local, densely, packed polygonal corallites (subcerioid growth form). White arrows show densely packed rounded corallites. **C** – transverse section, made five millimeters above the surface on Figure B, with densely packed rounded corallites (black arrow). White arrows show loosely packed branches (compare with B). **D** – two corallites with well developed marginarium and bilateral symmetry, defined by long primary septum (Vis 1/2). **E** – corallite with destroyed, inner parts of septa (Vis 1/1). **F** – dolomite crystals in microbial automicrite (see Fig. 9B) in sample UJ 225 P – Vis 2/2. After staining of the thin section with Alizarine Red-S, only coral skeleton (c) and automicrite show reddish color




Fig. 16. *Pleuroaulastrea variabilis* gen. et sp. nov; holotype UJ 225 P – Vis 1. **A, B** – corallites with parricidal budding. Arrows in **A** show rudist shell (*r*). **C–E** – corallite with short septa possibly due to state of preservation; in **D** and **E** (enlargement of **C**), septa show granular ornamentation. **B–E** – thin section Vis 1/4

Diagnosis: Phaceloid corallum. Septa arranged in six subequal systems, in radio-bilateral symmetry. Larger septa are rhopaloid. A slightly enlarged, primary septum is present in some corallites. The wall is thick. Around the outer wall surface of some adult corallites there occur structures, resulting from a wall “exfoliation”. These consist of one or more “layers”, more or less parallel to the wall, and usually do not continuously surround it. Endotheca poorly preserved (one-zonal?). Budding extracalicular marginal and by septal division.

Stratigraphic distribution: Lower part of middle Upper Barremian.

Remarks: The new genus is similar to *Carolastraea* (Eliášová, 1976a). It differs in the presence of septal division and structures, resulting from a wall “exfoliation” (Fig. 17A–J) and in showing some analogy with marginarium in amphiastreaids. However, in contrast to the interseptal space in many corallites, the space between the wall and these structures is filled by sediment, not by sparite cement, implying that there were no dissepiments, which might close the space. In this respect, it differs from marginarium, which commonly contains dissepiments or septa (compare *Metaulastrea* cf. *rarauensis*, Fig. 10). Melnikova and Roniewicz (1976, p. 98) discussing the wall structure in *Mitrodendron ogilviae* Geyer and *Pleurophyllia trichotoma* de Fromental from the Kimmeridgian of Poland, stated: “The wall develops in result of successive actions of skeletal secretion (pl. XXVIII, figs 1, 2), it is ‘multilamellar’ as was assumed by Beauvais (1974)”. Melnikova and Roniewicz (1976, pl. 23, fig. 1) showed (vertical section) structures termed as “rudiments of calicular border” similar to those described in the preset paper and named here as structures resulted from wall “exfoliation”.

Paracarolastraea zlatarskii new species, Kołodziej
Figs 17, 18 A–D

Holotype: UJ 225 P – Ru 34.

Paratypes: UJ 225 P – Ru 20-08, Ru 32, Ru 33.

Etymology: Named in honour of Dr. Vassil Zlatarski, a researcher of fossil and recent corals.

Type locality: Rusalya, Bulgaria.

Type level: Lower part of middle Upper Barremian, Emen Formation, Bulgaria.

Stratigraphic distribution: Lower and middle parts of Upper Barremian.

Material: Six samples UJ 225 P – Ru 20/08, Ru 31, Ru 32, Ru 33, Ru 34, Hot 1171; 16 thin sections (two large).

Diagnosis: *Paracarolastraea* with corallite diameter of 3–5 (6) mm, and 24 septa arranged in three size orders; very rare septa of the fourth order.

Description: Phaceloid corallum, only in places corallites are closely packed. In all samples corallite diameter is similar, ranging from 3 to 5 (6) mm. Septa arranged in six subequal systems, in radio-bilateral symmetry. Usually four of six S1 are longer and thicker than others. Larger septa (S1 and some S2) are commonly thickened at the inner margin (rhopaloid septa). A slightly enlarged primary septum occurs in some corallites (Fig. 17B, J). Septa S2 are S3 usually well developed, septa S4 rare. Septal faces are smooth or show faint, irregular granulations. Many adult corallites (up to 30% per thin section) contain structures resulting from “exfoliation” of the outer wall. These structures usually discontinuously surround the wall. Endotheca poorly preserved (uni-zonal?). Budding extracalicular marginal and by septal division. Septal budding results in formation of two (Fig. 18A) or more daughter corallites (Fig. 18B–D).




Fig. 17. *Paracarolastraea zlatarskii* gen. et sp. nov. **A–J** – transverse section. **K** – longitudinal section. Most of corallites on **A–J** show singular (**A**, **B**, **J**) or multiple (**C–I**) wall structures, resulting from ‘exfoliation’ of corallite wall. In **G** and **H**, these structures continuously surround corallite resulting in formation of some kind inner corallite. In **B**, corallites are highly variable in size and densely packed locally. UJ 225 P; **A**, **C**, **D** – Ru 34/4, **B** – Ru 34/3, **E**, **F** – Ru 34/4a, **G** – Ru 34/7, **H**, **I**, **K** – Ru 34/2. **A–G**, **J** – holotype, **H**, **I** – paratype

Remarks: With respect to diameter of corallites and number of septa the new species is similar to *Carolastraea gracea* Baron-Szabo, 1996 [corallites diameter (2.5) 3–4; number of septa (20) 24] from the Aptian of Greece (Baron-Szabo and Steuber, 1996). However, other features (‘exfoliation’ of a wall, septal increase) justify classification of the new species in the new genus *Paracarolastraea*. The septal division shows similarity to the one, re-

ported in some Upper Triassic scleractinian corals (Roniewicz, 1989), and in the middle Permian scleractiniomorph, *Numidia-phyllum* (Ezaki, 2004), where both bipartite, as well as hexapartite and tripartite increases occur.

Distribution: Lower and middle parts of Upper Barremian (Rusalya, Hotnitsa).




Fig. 18. A–D. Septal increase in *Paracarolastraea zlatarskii* gen. et sp. nov. In C and D, septal increase resulted in origin of four daughter corallites (numbered from 1 to 4); *m* in A indicates poorly laminated, microbialite crust. UJ 225 P; A – Hot 1171/1, B, D – Ru 32/1, C – Ru 32/2. E–I. *?Paracarolastraea* sp. E–H – septal increase resulted in origin of a few daughter corallites. In H densely packed corallites are surrounded locally by common wall (arrow). *a* and *m* in E and F show dasyclad algae (E – *Zittelina hispanica* Masse, Arias et Vilas, F – *Neomeris* cf. *cretacea* Steinmann) and microbialite crust respectively. I – corallite with thick wall. Arrow possibly indicates early stage of blastogeny. UJ 225 P; E, G – Vis 40/1, F – Vis 30/1, I – Vis 30/2

?Paracarolastraea sp.
Figs 6A, 18E–G

Material: Two samples (UJ 225 P – Vis 30, Vis 40), three thin sections.

Remarks: Two specimens show similarities to *Paracarolastraea zlatarskii* sp. nov., with respect to diameter and number of septa, but they lack structures resulting from ‘exfoliation’ of the wall. Moreover, septal budding, resulting in the formation of even five

daughter corallites (Fig. 18E–H), is more common, although only three thin sections were studied. In one thin section, a unique aggregation of corallites occurs. Densely packed corallites are surrounded locally by a common wall (possibly the wall of the parent individual). More serial sections are needed to reveal the exact increase pattern in this specimen, as well as its taxonomy (*Paracarolastraea zlatarskii* or a new species).

Distribution: Lower part of middle Upper Barremian (Vishovgrad).




Fig. 19. ?*Donacosmilia* sp. A–C – transverse sections of corallites; note lack of marginarium and slightly bilateral symmetry. UJ 225 P–Ru 7270/1




Fig. 20. *Intersmilia* aff. *diaboli*. A–C – transverse sections of corallites; note dark color of wall. C – parricidal budding (arrow). ZPAL Bułg 3; A – 3/6, B – 3/4, C – 3/2

Family DONACOSMILIIDAE Krasnov, 1970

Genus *Donacosmilia* de Fromentel, 1861

Type species: *Donacosmilia corallina* de Fromentel, 1861

?*Donacosmilia* sp.

Fig. 19

Material: One sample (UJ 225 P – Ru 7270), one large thin section.

Measurements (in mm): $d = (7) 8–10$ (11), $s = 41–47$ (60) in adult corallites.

Remarks: Phaceloid corallum with some corallites, showing slightly marked, bilateral symmetry. Rare, lonsdaleoid septa present. The specimen described differs from the best known, Late Jurassic species *Donacosmilia corallina* de Fromentel and *D. etalloni* (Koby, 1888) (e.g., Turnšek, 1972, 1997; Melnikova and Roniewicz, 1976; Buzcu and Babayiğit, 1998) in having smaller corallite diameters and a greater number of septa. *D. massaliensis* Morycowa et Masse, 1998, from the Upper Barremian of Provence in France, shows smaller corallite diameters (6–7.5 mm) and less numerous septa (32 S1–S3 plus S4) (Morycowa and Masse, 1998). In the available fragment of corallum (15 corallites), no marginarium was observed in corallites. Thus, the generic attribution of this species is uncertain.

Distribution: Lower part of middle Upper Barremian (Rusalya).

Family INTERSMILIIDAE Melnikova et Roniewicz, 1976

Genus *Intersmilia* Eliášová, 1974

Type species: *Intersmilia malevola* Eliášová, 1974

Remarks: *Intersmilia* is a rare genus, known to occur since the Lower Jurassic (?Hettangian–Sinemurian; Melnikova and Roniewicz, 1976, 2002), but mostly from the Upper Jurassic/earliest Cretaceous (Eliášová, 1974; Roniewicz, 1976; Kołodziej, 2003; Roniewicz, 2008).

Intersmilia aff. *diaboli* Eliášová, 1974

Fig. 20

Material: One sample (ZPAL Bułg 3), eight thin sections.

Measurements (in mm): $d = 6–8$ (9), $s = 12$.

Remarks: With respect to corallite diameters, the specimen is similar to *Intersmilia diaboli* Eliášová [$d = (6) 8–9$ (10)]. However, the number of septa in *I. diaboli* is from 12 to 24 (S1–S3, sporadically S4). It is possible that septa S3 occur also in the specimen studied, but they are unrecognizable, owing to the state of preservation. The wall is dark in color, particularly in its outer part. The wall shows a rather complex, inner structure, but no double-layer structure as in *I. diaboli* was recognized. According to Eliášová (1974) *Intersmilia* displays intracalicular marginal budding. However the published pictures do not support this. It appears rather, as stated by Melnikova and Roniewicz (1976), that the budding is lateral. Parricidal budding was not observed by Eliášová (1974), neither in *I. diaboli*, nor in *I. malevola* Eliášová, 1974. It was recognized in one corallite of the sample, studied here, and in the Kimmeridgian *I. irregularis* Roniewicz, 1976 (see Melnikova and Roniewicz, 1976). In the specimen studied (Fig. 20C), as in *I. irregularis*, the septa of the parent individual continue into the daughter one (Melnikova and Roniewicz, 1976, fig. 4). Symmetry of the septal apparatus in *Intersmilia* was described by Eliášová (1974) as radial. However, pictures of *I. malveola* and particularly of *I. diaboli* (Eliášová, 1974, pl. 1, fig. 1, pl. 2, fig. 2, pl. 3, fig. 1, pl. 4, figs 1, 2) permit recognition of a longer, primary septum. Accordingly, the septal symmetry should be rather characterized as radio-bilateral.

Distribution: The sampling site can not be exactly located (Veliko Tarnovo or its close vicinity), therefore a general Barremian age is assumed for the studied specimen. *I. diaboli* is known from Tithonian–Lower Berriasian of the Štramperk Limestone.

Family HETEROCOENIIDAE Oppenheim, 1930

Remarks: The four species, described below, represent four phaceloid genera. In contrast to other pachythealiines, the phaceloid growth form is rare in Heterocoeniidae. The systematic position of phaceloid genera *Cuneiphyllia* (one species; Eliášová, 1978) and *Pachythecephyllia* (one species; Kołodziej, 2003) from the Štramberk Limestone and Štramberk-type limestones respectively, originally attributed to the Heterocoeniidae, appears to be problematic, but certainly they represent pachythealiines.

Genus “*Pseudopistophyllum*” Geyer, 1955

Type species: *Pseudopistophyllum berckhemeri* Geyer, 1955

Remarks: The new species, described below, is tentatively assigned to the genus “*Pseudopistophyllum*” Geyer, 1955. This species represents a new genus and was for the first time illustrated by Kołodziej *et al.* (2011b) and determined informally as Gen. nov. 2. sp. nov. 1. Recently Löser *et al.* (in press) proposed establishment of a new genus, based on a new species, recognized in the Upper Albian of Spain (see also Löser, 2009, fig. 101; Löser *et al.*, 2011). Apart from the Spanish and Bulgarian species, the new genus includes also the species, described from Slovenia by Turnšek (in Turnšek and Buser, 1976), as *Pseudopistophyllum quinquesepatum*. This species was recognized in a limestone block of possible Late Jurassic age, occurring in Senonian breccias (Turnšek and Buser, 1976). The new genus differs from *Pseudopistophyllum* in having a different, septal pattern. In *Pseudopistophyllum* septa are developed on all sides of the wall, although the septa on one side are much longer. In the new genus, in specimens from Bulgaria, Slovenia and Spain, 3–5 septa are strongly developed on one side of corallites; other septa are absent, or if present, very rare and short (septal spines?). The genus is classified here in the Heterocoeniidae. In general, the septal pattern and poor developed septa of the new genus are similar to those, occurring in some heterocoeniids, such as *Latusastrea* and *Thecidiosmilia*. New buds were established in the wall of the parent corallite, but in contrast to amphistraeids, further growth was outside of a calice of parent individuals.

“*Pseudopistophyllum*” *triseptatum* new species, Kołodziej
Fig. 21

2011b. Gen. nov. 2. sp. nov. 1. – Kołodziej *et al.*, fig. 1g, h.

Holotype: UJ 225P – Zar 996.

Paratype: UJ 225P – Zar 2-08.

Etymology: *triseptatum* – named after occurrence of three dominated septa.

Type locality: Zarapovo, Bulgaria.

Type level: Lower part of middle Upper Barremian, Emen Formation, Bulgaria.

Stratigraphic distribution: Lower part of middle Upper Barremian.

Diagnosis: “*Pseudopistophyllum*” with longer diameter ranging from (2) 2.5 to 3 (4) mm and a shorter one from (1.5) 2 to 2.5 (3) mm. Three, rarely four septa occur on one side of a corallite. Other septa are absent or very rare and short (septal spines?).

Material: Two samples (holotype UJ 225P – Zar 962, paratype UJ 225P – Zar 2-08), eight thin sections (two large).

Measurements (in mm): Long d = (2) 2.5–3 (4), small d = (1.5) 2–2.5 (3), s = 3–4; other septa rare or absent.

Description: Phaceloid corallum. Locally corallites are densely packed resulting in a pseudocerioid growth form. Septa rare, arranged in bilateral symmetry defined by 3–4 long septa on one side of the corallite wall. On the opposite side, septa (septal spines?) are absent or very rare and, if present, very short (septal

spines?). The primary septum is usually strongest of all. In young corallites, only one septum can be recognized. A marginarium was recognized only in a few corallites. Endotheca unknown. New buds were established in a wall, but their growth, in contrast to Taschenknospung, was outside of the parent corallite.

Remarks: The new species differs from the one described by Löser *et al.* (in press; see also Löser, 2009, fig. 101) in having smaller corallite diameter, and in the number and character of the septa. In the species, described by Löser *et al.* (in press) there are 4–5 distinct septa, which have rhopaloid, T-shaped tips. *Pseudopistophyllum quinquesepatum* has larger corallites (3–6 mm) and a unique, septal pattern, with 5 long septa, reaching almost the opposite side of the wall. As in some other pachythealiines, in particular heterocoeniids, the number of septa is difficult to counted (e.g., Morycowa, 1971; Kołodziej, 1995, 2003; Löser, 2009; Löser *et al.*, 2009).

Distribution: Lower part of middle Upper Barremian (Zarapovo).

Genus *Heterosmilia* new genus, Kołodziej

Type species: *Heterosmilia spinosa* sp. nov., Kołodziej

Etymology: *Heterosmilia* – indicating its similarity to heterocoenid corals.

Stratigraphic distribution: Lower part of middle Upper Barremian.

Diagnosis: Phaceloid corallum. Bilateral, septal symmetry, with mostly distinct septa of subsequent orders. Primary septum displays strong, lateral outgrowths. Other septa lacking ornamentation. Lonsdaleoid septa present. Large dissepiments are clearly visible in transverse sections, but endotheca unknown. Budding extracalicular, lateral.

Remarks: The new genus shows similarities both with Heterocoeniidae as well as with Amphistraeidae and even Intersmiliidae. Strong septal outgrowths are characteristic for many heterocoeniids (e.g., Kołodziej, 1995; Schöllhorn, 1998; Löser, 2008a, b, 2010), but are absent in amphistraeids and intersmiliids. Transverse sections through the type species, i.e. *Heterosmilia spinosa* show some similarities to the solitary *Hexasmiliopsis* Löser, 2008 (see below remarks on *H. spinosa*).

Heterosmilia spinosa new species, Kołodziej
Fig. 22

2011b. Gen. nov. 1. sp. nov. 1. – Kołodziej *et al.*, fig. 1f.

Holotype: UJ 225P – Ru 20-09.

Paratype: UJ 225P – Ru 6-08.

Etymology: *spinosa* – named for spiny-like lateral outgrowths in the primary septum.

Type locality: Rusalya, Bulgaria.

Type level: Lower part of middle Barremian, Emen Formation, Bulgaria.

Stratigraphic distribution: Lower part of middle Upper Barremian.

Diagnosis: *Heterosmilia* with corallites of (7) 8–10 (11) in diameter. Septal pattern mostly regular: 6 S1 and 6 S2 in adult corallites. One or two sectors in some corallites are much wider than others. Septa S3, if developed, present only in two sectors on both sides of the primary septum.

Material: Four samples (UJ 225P – Ru 6-08, Ru 11-08, Ru 20-8, Ru 22-09), six thin sections (three large).

Measurements: Morphometric measurements are presented in Table 3.

Description: Phaceloid corallum. Septa arranged in bilateral symmetry defined by the presence of the primary septum, and in some corallites the presence of two wider sectors. In adult corallites, septal apparatus shows six septa S1 and six S2. Only two S3 are




Fig. 21. “*Pseudopistophyllum*” *triseptatum* sp. nov. **A** – loose, phaceloid corallum. **B** – densely packed corallites. **C** – row of four coralites with 3–4 recognizable septa, arranged in same direction. **D** – coralites with thick wall and one prominent septum and two other poorly developed septa. **E**, **F** – coralites with poor marginarium. **G** – coralites with ‘pockets’ (arrows) in wall, which possibly correspond to early stage of new calice formation; further growth is external, as shown in **H** and **I** (arrow). UJ 225 P; A–E, H–Zar 996/6; F, I–Zar 2/08/1, G–Zar 996/5. A–E, G–H–holotype, F, I–paratype

developed in two wider sectors (if present) on both sides of the primary septum. Septa S2 in these two sectors are longer, giving impression that there are eight septa S1. Septa are thicker at the wall and thinner at the inner margin. A primary septum shows strong, septal outgrowths. Ornamentation on lateral faces of other septa is lacking. Large dissepiments present and may form regular zones (Fig. 22G). Endotheca unknown. On basis of relations of parent and daughter corallites, budding possibly is extracalicular, lateral.

Remarks: Dissepiments, observed in transverse section, are well

developed. If the interseptal space is filled with sediment (not by calcite cement), it provides a better image of septal pattern and septal ornamentation. Transverse sections through *Heterosmilia spinosa* sp. nov. display similar corallite morphology and septal pattern, as in some transverse sections through the central part of the solitary heterocoeniid *Hexasmiliopsis saldanai* (Löser, 2008b, fig. 4a). Apart from the type specimens, two others (Ru 11-08, Ru 22-09) are attributed to this species. However, the poor state of preservation does not permit detailed measurements.

Distribution: Lower part of middle Upper Barremian (Rusalya).

Table 3

Dimensions (in mm) of *Heterosmilia spinosa* sp. nov.

Sample number	d	s
Holotype UJ 225P - Ru 20-09	(8) 9–10 (11)	12 +S3
Paratype UJ 225P - Ru 6-08	(7) 8–10	12? +S3

Genus *Hexasmilia* de Fromentel, 1870

Type species: *Hexasmilia ferryi* de Fromentel, 1870

Remarks: This poorly known genus (Upper Barremian–Santonian) was recently reviewed by Löser (2008a). *Hexasmilia* differs from *Heterocoenia* Milne Edwards and Haime, 1848 in its phaceloid growth form and the presence of strong septal outgrowths.




Fig. 22. *Heterosmilia spinosa* gen. et sp. nov.; A–E – holotype, UJ 225 P – Ru 20; F–H – paratype, UJ 225 P – Ru 6-08. A – corallite with 6 septa S1, 6 S2; note that only in two sectors (arrows) septa S3 are developed. See also different size of sectors on E. B–H – transverse sections, showing corallites with well preserved (B, F, G, H) and not preserved (C–E) dissepiments. Arrows on H show lonsdaleoid septa. Microfossils: *s* sponge, *se* serpulids, *a* dasyclad alga *Zittelina hispanica* Masse, Arias et Vilas, *c* remain of crustacean *Carpathocancer? plassenensis* (Schlagintweit et Gawlick). A–B – Ru 20/1, C–D – Ru 20/2, E – Ru 20/3, F–H – Ru 6-08/1

Three species were included by Löser (2008a): *H. ferryi* de Fromentel, *H. pachythealia* (Kuzmicheva, 1980), and new species *H. elmari*.

?*Hexasmilia* sp.

Fig. 23

Material: One sample (UJ 225P – Ru 7862), three thin sections (one large).

Dimensions (in mm): $d = (5) 6-7$, $s = 12 + S3$.

Remarks: The specimen from Bulgaria is similar to *Hexasmilia elmari* Löser, 2008, which shows some variability, expressed in differentially developed, septal apparatus. The specimen studied displays transverse sections, similar to *H. elmari* (compare Löser, 2008a, fig. 4, fig. 5.6, 5.8; and Fig. 23A, B in this paper) with six S1, including a primary septum. In other sections of *H. elmari*

(Schöllhorn, 1998, pl. 21, fig. 8; Löser, 2008a, fig. 5.1, 5.4, 5.5), there is a prominent, primary septum with strong septal outgrowths, while other septa are poor and less regularly developed. It is possible that more thin sections through the Bulgarian specimen will reveal greater, morphological variability. The septal division, observed by Löser (2008a) in *H. elmari*, was not observed in the specimen studied.

Distribution: Lower part of middle Upper Barremian (Rusalya).

Genus *Hexapetalum* Eliášová, 1975

Type species: *Hexapetalum impium* Eliášová, 1975

Remarks: The poorly known genus *Hexapetalum* was originally placed by Eliášová (1975) among the Amphistraeidae, later in the separate family Hexapetalidae (Eliášová, 1976b; Kuzmicheva, 1980). Kołodziej (2003) transferred this genus to the Heterocoeniidae.




Fig. 23. ?*Hexasmilia* sp. A – transverse section of corallite, with thick septa S1 and septal outgrowths, particularly strongly developed on primary septum (B). C – corallite with different, septal pattern (lack internal invaginations of the wall); *m* on B, C indicate microbialite crusts. UJ 225 P; A, B – Ru 7862/1, C – Ru 7862/2




Fig. 24. ?*Hexapetalum* sp.; A – transverse section of corallite with thick septa S1 and distinct, granular, septal ornamentation; UJ 225 P – Ru 2-08/1. B, C – corallites, showing external concavities of wall (arrows on C); UJ 225 P – Ru 40-08/2

?*Hexapetalum* sp.
Fig. 24

Material: Three samples, six thin sections.

Measurements: Morphometric measurements are presented in Table 4.

Table 4

Dimensions (in mm) of ?*Hexapetalum* sp.

Sample number	d	s
UJ 225P - Ru 40-08	4 (5)	6S1 + S2 + S3
UJ 225P - Ru 2	(4) 5 (6)	6 S1 + S2 + S3
ZPAL Bulg 2	(3) 4–5 (6)	6 S1 + S2

Remarks: Six septa S1 are well developed. Septa S2 are poorly and irregularly developed, septa S3 very rare or absent. One septum S1 is more or less longer. In a poorly preserved specimen Bulg 2, only six septa S1 are recognizable. Septa S1, if well preserved, show septal outgrowths or spiny ornamentation. Budding lateral, extracalicular. Wall morphology in some corallites display external concavities and internal invaginations, corresponding to septa (Fig. 23B, C). In this respect, the specimens described are similar to *Hexapetalum impium* Eliášová, *H. pium* Eliášová (Eliášová, 1975, pl. 9, fig. 2a; Eliášová, 1976b, pl. 2, figs 2, 3; Kołodziej, 2003, fig. 21), *H. pachythecallium* Kuzmicheva and *Heteroconia* spp. (Beauvais, 1982; Baron-Szabo, 1998; fig. 2; Löser, 2009, fig. 174). In particular, in *Hexapetalum pium*, the septa and wall show clear, structural continuity. The specimen, described here, has strong, septal, granular ornamentation (in well preserved coralli-

tes), in which it differs from other corals, described as *Hexapetalum*.
Distribution: lower part of middle Upper Barremian (Rusalya); Barremian (Veliko Tarnovo or its close vicinity).

CONCLUSIONS

1. The Upper Barremian limestones of the Emen Formation of the Lovech Urgonian Group in the Veliko Tarnovo area (Fore-Balkan, northern central Bulgaria) contain bioconstructions (possibly mostly biostromes) with a distinctive coral community, dominated by diversified corals of the extinct suborder Pachythecaliina. Its higher systematic position is debatable, but following some other coral specialists, these corals are classified here not in the Scleractinia, but in the order Hexanthiniaria. The corals described are mostly of the phaceloid growth form, with *Pleurophyllia bulgarica* sp. nov. particularly frequent. Other corals, except for the phaceloid *Calamophylliopsis*, are rare.

2. Small, monopleurid, cylindrical rudists *Mathesia darderi*, frequently densely clustered, and non-laminated microbialites provided additional, structural support for bioconstruction growth. Other macrofauna are only moderately common and diversified. *Lithocodium aggregatum* (possibly chlorophycean alga) and microbial, ‘bacinellid’ structures are rare, although they are common in many other coral-bearing limestones of the Lovech Urgonian Group.

3. The association of phaceloid pachythecaliine corals and *M. darderi*, accompanied by microbialites, is unique,

worldwide. The resulting bioconstructions are known only from the Emen Formation and show limited, regional extent, from Veliko Tarnovo to about 20 km NW. Biostratigraphic data indicate that this biofacies was developed, mainly during the early middle Late Barremian (Gerhardtia sartousiana Zone).

4. The section at the Rusalya Quarry, about 42 m thick provides, the sedimentary and environmental context of the reefal biostromes. Bioclastic packstones of unit 1 contain small boundstone patches, with corals (but not pachythealiines or other branching corals), calcified sponges (mainly chaetetids) and encrustations of *L. aggregatum*. Units 2 and 3 consist of bioclastic limestones (rare corals) interlayered with rudist limestones. Pachythealiine-*Mathesia*-microbialite biostromes (unit 4) developed in a narrow 2.5-m-thick interval, are covered by biostromes, built mostly of large, monopleurid rudists (unit 5). The sedimentary succession shows, with some fluctuations, a general shallowing trend, from the outer to inner carbonate platform. Pachythealiine-rich biostromes were developed on the distal part of the rudist-dominated area of the carbonate platform, in a low-energy setting, with low rates of net, background sedimentation and possibly a higher nutrient level. This environment favoured growth of phaceloid pachythealiines, a nearly monospecific rudist community (*M. darderi*), and microbialite development, but limited growth of phototrophic/oligotrophic microencrusters.

5. It is hypothesized that some of microbialites (*sensu lato*) resulted from decaying sponges. Dolomite, recognized in the microbialites (Vishovgrad site), is interpreted as resulting from precipitation, linked to microbial activity or organomineralization *s.s.*

6. Taxonomic diversity and abundance of phaceloid pachythealiines show a spatially and temporally restricted pattern. They are only more common in the Tithonian–Lower Berriasian Štramberg Limestone (Czech Republic) and its equivalent in the Polish Carpathians. However, the lithology and associated biota imply that pachythealiines from the Emen Formation developed in a different, environmental setting. Other coral-bearing deposits of the mixed, siliciclastic-carbonate Urganian complex in Bulgaria contain diversified coral communities, but pachythealiines are very poorly diversified and phaceloid pachythealiines are absent.

7. Fourteen species (including six new and eight in open nomenclature), representing twelve genera (four new, three of them are distinguished formally) and five families (Amphistraeidae, Carolastraeidae, Intersmiliidae, Donacosmiliidae, Heterocoeniidae), are described. Four heterocoeniid species (two new) from four genera (one new) are phaceloid, thus displaying a growth form, generally rare in this family. This suggests similar, environmental demands for phaceloid heterocoeniids and other phaceloid pachythealiines and may support phylogenetic relationships between them.

Acknowledgements

These studies were financed by a research grant from the Polish Ministry of Science and Higher Education (No. 2P04D 028 29). The referees, Ewa Roniewicz (Warszawa) and Felix Schlag-

intweit (München), are thanked for their corrections and valuable comments that have led to improvements in this paper. Elżbieta Morycowa (Kraków) is thanked for providing helpful discussion on coral taxonomy and comments on the manuscript. We are grateful to Ioan Bucur (Cluj-Napoca) for determinations of dasyclad algae, Hannes Löser (Hermosillo) for discussion of some coral taxa, Jean-Pierre Masse (Marseille) for rudist determinations, and Felix Schlagintweit for discussion concerning microproblematica. E. Roniewicz kindly provided thin sections, taken from samples collected by the late Prof. Ryszard Marcinowski. Antoaneta Ilcheva and Neda Motchurova-Dekova (Sofia), as well as Vassil Zlatarski (Bristol, Rhode Island) helped during the study of his coral collection housed at the National Museum of Natural History in Sofia. V. Zlatarski accompanied us in the first season of field studies. The authorities of the Rusalya Quarry are thanked for permission to conduct studies in the quarry. We are indebted to Frank Simpson (Windsor, Ontario) for English corrections and Michał Gradziński and Alfred Uchman (Kraków) for editorial help. Special thanks go to Ewa Roniewicz and Vassil Zlatarski for establishing the Polish–Bulgarian collaboration on studies of Urganian corals in Bulgaria.

REFERENCES

- Adams, A. E., MacKenzie, W. S. & Guilford, C., 1984. *Atlas of Sedimentary Rocks under the Microscope*. Longman, Burnt Mill, Harlow, 104 pp.
- Alloiteau, J., 1957. *Contribution à la Systématique des Madréporaires Fossiles*. Centre National de la Recherche Scientifique, Paris, 462 pp.
- Aubrecht, R., Szulc, J., Michalik J., Schlögl, J. & Wagneich, M., 2002. Middle Jurassic stromatactis mud-mound in the Pieniny Klippen Belt (Western Carpathians). *Facies*, 47: 113–126.
- Baron-Szabo, R. C., 1998. A new coral fauna from the Campanian of Northern Spain (Torallola Village, Prov. Lleida). *Geologisch-Paläontologische Mitteilungen Innsbruck*, 23: 127–198.
- Baron-Szabo, R. C., 2002. *Scleractinian Corals of the Cretaceous. A Compilation of Cretaceous Forms with Descriptions, Illustrations and Remarks on their Taxonomic Position*. Baron-Szabo, Knoxville, 539 pp.
- Baron-Szabo, R. C., 2006. Corals of the K/T-boundary: Scleractinian corals of the suborders Astrocoeniina, Faviina, Rhipidogyrina, and Amphistraeina. *Journal of Systematic Palaeontology*, 4: 1–108.
- Baron-Szabo, R. C. & Steuber, T., 1996. Korallen und Rudisten aus dem Apt im tertiären Flysch des Pannass-Gebirges bei Delphi-Arachowa (Mittelgriechenland). *Berliner Geowissenschaftliche Abhandlungen*, E18: 3–75.
- Beauvais, L., 1974. Studies upon the characters of the suborder Amphistraeida Alloiteau. In: Sokolov, B. S. (ed.), *Drevnye Cnidaria*, Izdatelstvo Nauka, Novosibirsk, pp. 238–248.
- Beauvais, L., 1976. Étude morphologique, taxonomique et phylogénétique du sous-ordre Amphistraeida Alloiteau (Madréporaires jurassiques). *Mémoires de la Société Géologique de France, n.s.*, 126: 5–42.
- Beauvais, L., 1981. On the taxonomy of the Mesozoic Madreporearia. *Acta Palaeontologica Polonica*, 25: 345–360.
- Beauvais, M., 1977. Le nouveau sous-ordre des Heterocoeniida. *Mémoires du Bureau de Recherches Géologiques et Minières*, 89: 271–282.
- Beauvais, M., 1982. Révision Systématique des Madréporaires des couches de Gosau. *Travaux du Laboratoire de Paléontologie des Invertébrés, Paris*, 5 vols 1: 1–256; 2: 1–278; 3: 1–177; 4 and 5: atlas.

- Betzler, C., Pawellek, T., Abdullah, M. & Kossler, A., 2007. Facies and stratigraphic architecture of the Korallenoolith Formation in north Germany (Lauensteiner Pass Ith Mountains). *Sedimentary Geology*, 194: 61–75.
- Bontognali, T. R. R., Vasconcelos, C., Warthmann, R. J., Bernasconi, S. M., Dupraz, C., Strohmenger, C. J. & McKenzie, J. A., 2010. Dolomite formation within microbial mats in the coastal sabkha of Abu Dhabi (United Arab Emirates). *Sedimentology*, 57: 824–844.
- Bourque, P. A., 1997. Paleozoic finely crystalline carbonate mud mounds: cryptic communities, petrogenesis and ecological zonation. *Facies*, 36: 250–253.
- Bover-Arnal, T., Salas, R., Martin-Closas, M., Schlagintweit, F. & Moreno-Bedmar, J. A., 2011. Expression of an oceanic anoxic event in a neritic setting: Lower Aptian coral rubble deposits from the western Maestrat Basin (Iberian Chain, Spain). *Palaios*, 26: 18–32.
- Bucur, I. I., Hoffmann, M. & Kołodziej, B., 2005. Late Jurassic–lowermost Cretaceous benthic algae from Tethys and European platform: A case study from Poland. *Revista Española de Micropaleontología*, 37: 105–129.
- Buzcu, S. & Babayiğit, S., 1998. The occurrence of *Donacosmilia corallina* de Fromentel (Upper Jurassic) from the Kastamonu region in Turkey. *Geological Bulletin of Turkey*, 41: 99–107. [In Turkish, English abstract].
- Camoin, G., Cabioch, G., Eisenhauer, A., Braga, J.-C., Hamelin, B. & Lericolais, G., 2006. Environmental significance of microbialites in reef environments during the last deglaciation. *Sedimentary Geology*, 185: 277–295.
- Camoin, G. F., Gautret, P., Montaggioni, L. F. & Cabioch, G., 1999. Nature and environmental significance of microbialites in Quaternary reefs: the Tahiti paradox. *Sedimentary Geology*, 60: 15–49.
- Camoin, G. F. & Montaggioni, L. F., 1994. High-energy coralgastromatolite frameworks from Holocene reefs (Tahiti, French Polynesia). *Sedimentology*, 41: 655–676.
- Cherchi, A. & Schroeder, R., 2010. Boring sponges (*Entobia*) in Mesozoic *Lithocodium* calcimicrobial crusts. *Rivista Italiana di Paleontologia e Stratigrafia*, 116: 351–356.
- Cherchi, A. & Schroeder, R., 2013. Revision of the holotype of *Lithocodium aggregatum* Elliott, 1956 (Lower Cretaceous, Iraq): new interpretation as sponge–calcimicrobe consortium. *Facies*, 59: 49–57.
- Cheshitev, G., Kânčev, I., Vâlkov, V., Marinova, R., Shiljafova, J., Russeva, M. & Iliev, K., 1989. *Geological Map of P.R. Bulgaria, 1:500 000*. Committee of Geology, Department of Geophysical Prospecting and Geological Mapping, Sofia.
- Ciborowski, T. & Kołodziej, B., 2001. Tithonian–Berriasian calpionellids from the Štramberg-type limestones, Polish Flysch Carpathians. *Geologica Carpathica*, 52: 343–348.
- Coates, A. G. & Jackson, J. B. C., 1985. Morphological themes in the evolution of clonal and aclonal marine invertebrates. In: Jackson, J. B. C., Buss, L. W. I. & Cook, R. E. (eds), *Biology and Evolution of Colonial Organisms*. Yale University Press, New Haven, pp. 67–106.
- Coates, A. G. & Oliver, W. A., Jr., 1973. Coloniality in zoantharian corals. In: Boardman, R.S., Cheetham, A. H. & Oliver, W. A., Jr. (eds), *Animal Colonies Development and Function Through Time*, 3–27. Stroudsburg, Pennsylvania.
- Cuif, J.-P., 1975. Caractères morphologiques, microstructuraux et systématiques des Pachythecalidae nouvelle famille de Madréporaires Triassiques. *Géobios*, 8: 157–180.
- Cuif, J.-P., 1977. Arguments pour une relation phylétique entre les Madréporaires paléozoïques et ceux du Trias. Implications systématiques de l'analyse microstructurale des Madréporaires triassiques. *Mémoires de la Société Géologique de France*, 129: 1–54.
- Cuif, J.-P., 1981. Microstructure versus morphology in the skeleton of Triassic scleractinian corals. *Acta Palaeontologica Polonica*, 25: 361–374.
- Cuif, J.-P., 2010. The converging results of microstructural analysis and molecular phylogeny: Consequence for the overall evolutionary scheme of post-Paleozoic corals and the concept of Scleractinia. *Palaeoworld*, 19: 357–367.
- Cuif, J.-P. & Stolarski, J., 1999. Origin and paleobiology of wall-based corals. In: *Abstracts of the 8th International Symposium on Fossil Cnidaria and Porifera, Sendai, September 12-16, 1999*. p. 95.
- Delecat, S. & Reitner, J., 2005. Sponge communities from the Lower Liassic of Adnet (Northern Calcareous Alps, Austria). *Facies*, 51: 385–404.
- Dietrich, W. O., 1926. Steinkorallen des Malms und der Unterkreide im südlichen Deutsch-Ostafrika. *Palaeontographica* (suppl. 7), 1: 43–62.
- Dryer, S. & Logan, A., 1978. Holocene reefs and sediments of Castle Harbour, Bermuda. *Journal of Marine Research*, 36: 399–425.
- Dupraz, C., Reid, P. R., Braissant, O., Decho, A. W., Norman, S. R. & Visscher, P. T., 2009. Processes of carbonate precipitation in modern microbial mats. *Earth-Science Reviews*, 96: 141–162.
- Dupraz, C. & Strasser, A., 2002. Nutritional modes in coral-microbialite reefs (Jurassic, Oxfordian, Switzerland): evolution of trophic structure as a response to environmental change. *Palaios*, 17: 449–471.
- Eliáš, M. & Eliášová, H., 1984. Facies and palaeogeography of the Jurassic in the western part of the Outer Flysch Carpathians in Czechoslovakia. *Sborník Geologických Věd*, 39: 105–170.
- Eliášová, H., 1973. Sous-famille Rhipidogyrinae Kobay, 1905 (Hexacorallia) des calcaires de Štramberg Tithonien, Tchécoslovaquie. *Časopis pro Mineralogii a Geologii*, 18: 267–287.
- Eliášová, H., 1974. Genre nouveau *Intersmilia* (Hexacorallia) du Tithonien des calcaires de Štramberg (Tchécoslovaquie). *Časopis pro Mineralogii a Geologii*, 19: 415–417.
- Eliášová, H., 1975. Sous ordre Amphipora Alloiteau 1952 (Hexacorallia) des calcaires de Štramberg (Tithonien, Tchécoslovaquie). *Časopis pro Mineralogii a Geologii*, 20: 1–23.
- Eliášová, H., 1976a. Nouvelle famille du sous-ordre Amphipora, 1952 (Hexacorallia), Tithonien de Tchécoslovaquie. *Věstník Ústředního Ústavu Geologického*, 51: 177–178.
- Eliášová, H., 1976b. Les coraux de l'ordre Hexanthinaria Montanaro-Gallitelli, 1975, Zoantharia de Blanville, 1830, dans les calcaires de Štramberg (Tithonien, Tchécoslovaquie). *Věstník Ústředního Ústavu Geologického*, 51: 357–366.
- Eliášová, H., 1978. La rédefinition de l'ordre Hexanthinaria Montanaro Gallitelli, 1975 (Zoantharia). *Vestník Ústředního Ústavu Geologického*, 53, 89–101.
- Eliášová, H., 1981. The Tithonian reef of Štramberg Limestone (Czechoslovakia, West Carpathians). *Časopis pro Mineralogii a Geologii*, 26: 113–124.
- Ezaki, Y., 2004. Paleoecological and phylogenetic implications of asexual reproduction in the Permian scleractiniamorph *Numidiaphyllum*. *Journal of Paleontology*, 75: 84–97.
- Fedorowski, J., 1980. Some aspects of coloniality in corals. *Acta Palaeontologica Polonica*, 25: 429–437.
- Fedorowski, J., 1997. Rugosa and Scleractinia – a commentary methods of phylogenetic reconstructions. *Acta Palaeontologica Polonica*, 42: 446–456.
- Fenerci-Masse, M., 2006. *Les communautés à rudistes du Crétacé inférieur de la marge ouest européenne de la Téthys*. Unpub-

- lished PhD thesis, Université de Provence (Aix-Marseille I), 436 pp.
- Fenerci-Masse, M., Masse, J.-P., Kołodziej, B., Ivanov, M. & Idakieva V., 2011. *Mathesia darderi* (Astre) (Bivalvia, Hipuritoidea, Monopleuridae): morphological, biogeographical and ecological changes in the Mediterranean domain during the late Barremian–Albian. *Cretaceous Research*, 32: 407–421.
- Flügel, E., 2010. *Microfacies of Carbonate Rocks. Analysis, Interpretation and Application*. Springer, Berlin, 984 pp.
- Fürsich, F. T., Pandey, D. K., Oschmann, W., Jaitly, A. K. & Singh, I. B., 1994. Ecology and adaptive strategies of corals in unfavourable environments: Examples from the Middle Jurassic of the Kachchh Basin, Western India. *Neues Jahrbuch für Geologie und Paläontologie – Abhandlungen*, 194: 269–303.
- Geister, J., 1995. Sclerochronology of Jurassic corals: geological applications. *Publications du Service Géologique du Luxembourg*, 29: 150–152.
- Geyer, O. F., 1955. Beiträge zur Korallen-Fauna des Stramberger Tithon. *Paläontologische Zeitschrift*, 29: 177–216.
- Gili, E., Masse, J.-P. & Skelton, P. W., 1995a. Rudists as gregarious sediment-dwellers, not reef-builders, on Cretaceous carbonate platforms. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 118: 245–267.
- Gili, E., Skelton, P. W., Vicens, E. & Obrador, A., 1995b. Corals to rudists – an environmentally induced assemblage succession. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 119: 127–136.
- Götz, S., 2003. Biotic interaction and synecology in a Late Cretaceous coral-rudist biostrome of southeastern Spain. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 193: 125–138.
- Guilbault, J.-P., Krautter, M., Conway, K. W. & Barrie, J. V., 2006. Modern foraminifera attached to Hexactinellid sponge meshwork on the West Canadian Shelf: Comparison with Jurassic counterparts from Europe. *Palaeontologia Electronica*, 9: 1–48.
- Hallock, P., 1988. The role of nutrient availability in bioerosion: consequences to carbonate buildups. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 63: 275–291.
- Heindel, K., Birgel, D., Brunner, B., Thiel, V., Westphal, H., Gischler, E., Ziegenbalg, S. B., Cabioch, G., Sjövall, P. & Peckmann, J., 2012. Post-glacial microbialite formation in coral reefs of the Pacific, Atlantic, and Indian Oceans. *Chemical Geology*, 304–305: 117–130.
- Helm, C., 2005. Riffe und fazielle Entwicklung der *florigemma*-Bank (Korallenoolith, Oxfordium) im Süntel und östlichen Wesergebirge (NW-Deutschland). *Geologische Beiträge Hannover*, 7: 3–339.
- Helm, C. & Schülke, I., 2006. Patch reef development in the *florigemma*-Bank Member (Oxfordian) from the Deister Mts (NW Germany): a type example for Late Jurassic coral thrombolite thickets. *Facies*, 52: 441–467.
- Houša, V., 1990. Stratigraphy and calpionellid zonation of the Štramberk Limestone and associated Lower Cretaceous beds. *Atti del secondo convegno internazionale Fossili, Evoluzione, Ambiente, Pergola 25-30 ottobre 1987*, pp. 365–370.
- Höfling, R., 1997. Eine erweiterte Riff-Typologie und ihre Anwendung auf kretazische Biokonstruktionen. *Bayerische Akademie der Wissenschaften. Mathematisch-naturwissenschaftliche Klasse, Abhandlungen*, 169: 1–127.
- Huck, S., Heimhofer, U. & Immenhauser, A., 2012. Early Aptian algal bloom in a neritic proto–North Atlantic setting: Harbinger of global change related to OAE 1a? *Geological Society of America Bulletin*, 124: 1810–1825.
- Huck, S., Rameil, N., Korbar, T., Heimhofer, U., Wieczorek, T. D. & Immenhauser, A., 2010. Latitudinally different responses of Tethyan shoal-water carbonate systems to the Early Aptian oceanic anoxic event (OAE 1a). *Sedimentology*, 57: 1585–1614.
- Idakieva, V., 2002. Taxonomy of Lower Cretaceous corals from Lovech Urganian Group (Lower Cretaceous) in Central Fore-Balkan, Bulgaria. *Annuaire de l'Université de Sofia "St. Kliment Ohridski", Faculte de Géologie et Géographie*, 95: 5–16.
- Idakieva, V., 2003. *Dolnokredni skleraktinii v chast ot Tsentralnina Predbalkan (taksonomia i organogenni postroiki)*. Disertatsia, Sofiiski Universitet "Sveti Kliment Ohridski", 327 pp. [In Bulgarian].
- Idakieva, V., 2007. Taxonomy of scleractinian corals from the Barremian–Lower Aptian of Central North Bulgaria (Lovech Urganian Group). *Annuaire de l'Université de Sofia "St. Kliment Ohridski", Faculte de Géologie et Géographie*, 100: 29–66.
- Idakieva, V., 2008. Taxonomy of scleractinian corals from the Barremian–Lower Aptian of Central North Bulgaria (Lovech Urganian Group). *Annuaire de l'Université de Sofia "St. Kliment Ohridski", Faculte de Géologie et Géographie*, 101: 5–32.
- Idakieva, V. & Ivanov, M., 2002. Coral buildups in the Smochan Formation (Urganian, Lower Cretaceous) around Lovech (Central North Bulgaria). *Annuaire de l'Université de Sofia "St. Kliment Ohridski", Faculte de Géologie et Géographie*, 95: 31–46. [In Bulgarian, English abstract].
- Ilcheva, A. & Motchurova-Dekova, N., 2011. Catalogue of type collections of Early Cretaceous corals (Scleractinia, Anthozoa) at the National Museum of Natural History, Sofia. *Review of the Bulgarian Geological Society*, 72: 129–140.
- Immenhauser, A., Hillgärtner, H. & van Bentum, E., 2005. Microbial-foraminiferal episodes in the Early Aptian of the southern Tethyan margin: ecological significance and possible relation to oceanic anoxic event 1a. *Sedimentology*, 52: 77–99.
- Insalaco, E., 1998. The descriptive nomenclature and classification of growth fabrics in fossil scleractinian reefs. *Sedimentary Geology*, 118: 159–186.
- Insalaco, E., Hallam, A. & Rosen, B., 1997. Oxfordian (Upper Jurassic) coral reefs in Western Europe: reef types and conceptual depositional model. *Sedimentology*, 44: 707–734.
- Ivanov, M., 1995. Stratigraphy of Barremian and Aptian Stage in the southern part of Moesian platform in Central North Bulgaria. *Review of the Bulgarian Geological Society*, 56: 37–46. [In Bulgarian, English summary].
- Ivanov, M. & Idakieva, V., 2009. Upper Barremian ammonite biostratigraphy in Central-North and Northeastern Bulgaria. In: *Proceedings of National Conference Geosciences 2009*. Bulgarian Geological Society, Sofia, pp. 55–56. [In Bulgarian, English summary].
- Ivanov, M. & Nikolov, T., 1995. New data on the ammonitic occurrence in the Lovech Urganian Group (Lower Cretaceous) of the Central Fore-Balkan (Bulgaria). *Comptes Rendus de l'Académie bulgare des Sciences*, 48: 59–61.
- Ivanova, D. & Kołodziej, B., 2010. Late Jurassic–Early Cretaceous foraminifera from Štramberk-type limestones, Polish Outer Carpathians. *Studia Universitatis Babeş-Bolyai, Geologia*, 55: 3–31.
- Ivanova, D., Kołodziej, B., Koleva-Rekalova, E. & Roniewicz, E., 2008. Oxfordian to Valanginian palaeoenvironmental evolution on the Western Moesian Carbonate Platform: A case study from SW Bulgaria. *Annales Societatis Geologorum Poloniae*, 78: 65–90.

- Kennard, J. M. & James, N. P., 1986. Thrombolites and stromatolites: two distinct types of microbial structures. *Palaios*, 1: 492–503.
- Kershaw, S., 1994. Classification and geological significance of biostromes. *Facies*, 31: 81–92.
- Khrischev, K., 1966. Lithostratigraphy of the Lovech Urgonian Group. *Bulletin of the "Strachimir Dimitrov" Institute of Geology*, 15: 231–240. [In Bulgarian, English summary].
- Khrischev, K., 1992. Lower Cretaceous. In: Khrischev, K. & Nedyalkova, L. (eds), *Explanation to Geological Map of N. R. Bulgaria, 1:000 000, Sheet Sevlievo*, Publishing House Bulgarian Academy of Sciences, Sofia, pp. 20–33. [In Bulgarian, English summary].
- Khrischev, K. & Bakalova, D., 1974. Distribution of the algae in the Emen Limestone Formation (the Lovech Urgonian Group). *Bulletin of the Geological Institute, series Stratigraphy and Lithology, Bulgarian Academy of Sciences*, 23: 65–89.
- Kiessling, W., 2009. Geologic and biologic controls on the evolution of reefs. *Annual Review of Ecology, Evolution, and Systematics*, 40: 173–192.
- Kiessling, W., Flügel, E. & Golonka, J., eds, 2002. *Phanerozoic Reef Patterns*. Society of Economic Paleontology and Mineralogy Special Publications, vol. 72, Tulsa, OK, 775 pp.
- Kiessling, W., Roniewicz, E., Villier, L., Léonide, P. & Struck, U., 2009. An early Hettangian coral reef in southern France: Implications for the end-Triassic reef crisis. *Palaios*, 24: 657–671.
- Koby, E., 1888. Monographie des polypiers jurassique de la Suisse. *Memoires de la Société Paléontologique Suisse*, 25: 401–456.
- Kołodziej, B., 1995. Microstructure and taxonomy of Amphistraeina (Scleractinia). *Annales Societatis Geologorum Poloniae*, 65: 1–17.
- Kołodziej, B., 1997. Boring Foraminifera from exotics of Štramberg-type limestones (Tithonian–lower Berriasian, Polish Carpathians). *Annales Societatis Geologorum Poloniae*, 67: 249–256.
- Kołodziej, B., 2003. Scleractinian corals of suborders Pachythecaliina and Rhipidogyrina (Scleractinia): discussion on similarities and description of species from Štramberg-type limestones, Polish Outer Carpathians. *Annales Societatis Geologorum Poloniae*, 73: 193–217.
- Kołodziej, B., Bucur, I. I., Lazar, D. & Săsăran, E., 2011a. Barremian–Aptian coral facies from Romania. In: Aretz, M., Declulée, S., Denayer, J. & Poty, E. (eds), *11th Symposium on Fossil Cnidaria and Sponges, Liège, August 19–29, 2011, Abstracts. Kölner Forum für Geologie Paläontologie*, 119: 77–79.
- Kołodziej, B., Idakieva, V. & Ivanov, M., 2011b. Unique Early Cretaceous development of phaceloid pachythecaliines (Scleractinia?, Hexanthiniaria?): Upper Barremian, Bulgaria. In: Aretz, M., Declulée, S., Denayer, J. & Poty, E. (eds), *11th Symposium on Fossil Cnidaria and Sponges, Liège, August 19–29, 2011, Abstracts. Kölner Forum für Geologie Paläontologie*, 119: 80–82.
- Kołodziej, B., Idakieva, V., Ivanov, M. & Masse, J.-P., 2009. Unusual association of corals (Pachythecaliina) and small rudists from the Upper Barremian, Bulgaria. In: Hart, M. B. (ed.), *8th International Symposium on the Cretaceous System, Plymouth, 6–12 September, 2009, Abstracts*. University of Plymouth, Plymouth, pp. 160–161.
- Kołodziej, B., Idakieva, V., Ivanov, M. & Zlatarski, V., 2007. Coral growth forms and growth-interruptions in turbid-water bioconstructions: examples from Lower Cretaceous, Bulgaria. In: Kossovaya, O., Somerville, I. & Evdokimova, I. (eds), *10th International Symposium on Fossil Cnidaria and Porifera, Sankt Petersburg, August 12–16, 2007, Abstracts*. A.P. Karpinsky Russian Geological Research Institute, St. Petersburg, pp. 54–55.
- Kołodziej, B., Idakieva, V., Ivanov, M. & Zlatarski, V., 2011c. Clear- and turbid-water coral assemblages from Barremian–Lower Aptian of Bulgaria. In: Aretz, M., Declulée, S., Denayer, J. & Poty, E. (eds), *11th Symposium on Fossil Cnidaria and Sponges, Liège, August 19–29, 2011, Abstracts. Kölner Forum für Geologie Paläontologie*, 119: 83–85.
- Kuzmicheva, E. I., 1980. Corals. In: Chernov, V. G., Yanin, B. T. & Golovinova, M. A. (eds), *Urgonian sediments of the Soviet Carpathians*. Nauka, Moskva, pp. 90–108. [In Russian].
- Lathuilière, B., 1989. *Isastrea*, polypier branchu! *Comptes Rendus de l'Académie des Sciences, Paris*, 308, Serie 2: 887–892.
- Lathuilière, B., 1996. Itinéraires astogeniques chez des coraux simples et coloniaux montlivaltiides du Bajocien de France. *Géobios*, 29: 577–603.
- Lathuilière, B., 2012a. *Pleurophyllia*. In: Cairns, S. D., Baron-Szabo, R., Budd, A. F., Lathuilière, B., Roniewicz, E., Stolarski, J., and Johnson, K. G. (eds), *Corallosphere*. <http://corallosphere.org> [01.12.2012]
- Lathuilière, B., 2012b. *Aulastrea*. In: Cairns, S. D., Baron-Szabo, R., Budd, A. F., Lathuilière, B., Roniewicz, E., Stolarski, J., and Johnson, K. G. (eds), *Corallosphere*. <http://corallosphere.org> [01.12.2012]
- Lebanidze, Z. M., 1991. Pozdnejurskie korally zapadnoy Gruzii (Abhazija). *Trudy, Geologičeskij Institut, Akademija Nauk Gruzii, Novaja Serija*, 105: 1–64. [In Russian].
- Leinfelder, R. R., 1993. Upper Jurassic reef types and controlling factors – a preliminary report. *Profil*, 5: 1–45.
- Leinfelder, R. R., 2001. Jurassic reef ecosystems. In: Stanley, G. D., Jr. (ed.), *The History and Sedimentology of Ancient Reef Systems*. Kluwer Academic/Plenum Publishers, New York, pp. 251–302.
- Leinfelder, R. R., Krautter, M., Laternser, R., Nose, M., Schmid, G., Schweigert, G., Werner, W., Keupp, H., Brugger, H., Herrmann, R., Rehfeld-Kiefer, U., Schroeder, J. H., Reinhold, C., Koch, R., Zeiss, A., Schweizer, V., Christmann, H., Menges, G. & Luterbacher, H., 1994. The origin of Jurassic reefs: current research developments and results. *Facies*, 31: 1–56.
- Leinfelder, R. R., Nose, M., Schmid, D. U. & Werner, W., 1993. Microbial crusts of the Late Jurassic: Composition, palaeoecological significance and importance in reef construction. *Facies*, 29: 195–230.
- Leinfelder, R. R., Schmid, D. U., Nose, M. & Werner, W., 2002. Jurassic reef patterns – the expression of a changing globe. In: Kiessling, W., Flügel, E. & Golonka, J. (eds), *Phanerozoic Reef Patterns. Society of Economic Paleontologist and Mineralogists, Special Publication*, 72: 465–520.
- Löser, H., 1998. Cretaceous corals – state of knowledge and current research. *Zentralblatt für Geologie und Paläontologie*, 1: 1475–1485.
- Löser, H., 2008a. Remarks on the genus *Hexasmilia* (Scleractinia; Cretaceous) and description of a new species from the Aptian of Spain. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 250: 45–52.
- Löser, H., 2008b. A new solitary coral genus of the suborder Heterocoeniina (Scleractinia) from the Aptian (Cretaceous) of Spain. *Paläontologische Zeitschrift*, 82: 279–284.
- Löser, H., 2008c. Early Cretaceous coral faunas from East Africa (Tanzania, Kenya; Late Valanginian–Aptian) and revision of the Dietrich collection (Berlin, Germany). *Palaeontographica A*, 285: 23–75.

- Löser, H., 2009. *Fossile Korallen aus Jura und Kreide. Aufbau, Klassifikation, Bestimmung und Fundmöglichkeiten*. CPress Verlag, Dresden, 206 pp.
- Löser, H., 2010. Revision of the Cretaceous coral genus *Tiarasmilia* Wells, 1932 (Scleractinia). *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, 258: 157–165.
- Löser, H., 2012. Revision of the Amphistraeidae from the Monti d'Ocre area (Scleractinia; Early Cretaceous). *Rivista Italiana di Paleontologia e Stratigrafia*, 118: 461–502.
- Löser, H., Barattolo, F., Badia, S. C., Chikhi-Aouimeur, F., Dhondt, A., Erlich, R. N., Fözy, I., Geister, J., Hiss, M., Kołodziej, B., Leloux, J., Lewy, Z., Minor, K. P., Mitchell, S., Moosleitner, G., Peza, L., Remane, J., Romana, R., Sikharulidze, G. Y., Sinnyovski, D., Steuber, T., Tröger, K.-A., Turnšek, D., Vecchio, E., Vilella i Puig, J. & Zitt, J., 2002. *Catalogue of Cretaceous Corals, Vol. 2, List of Citations*. CPress Verlag, Dresden, 784 pp.
- Löser, H., Castro, J. M. & Nieto, L., 2011. An exotic Late Albian coral fauna from the Prebetic Platform (Betic External Zone, South Spain). In: Aretz, M., Declucée, S., Denayer, J. & Poty, E. (eds), *11th Symposium on Fossil Cnidaria and Sponges, Liège, August 19-29, 2011, Abstracts. Kölner Forum für Geologie Paläontologie*, 119: 99.
- Löser, H., Castro, J. M. & Nieto, L., in press. Late Albian Scleractinian corals from the Prebetic Zone (SE Spain). *Palaeontographica A*.
- Löser, H., Stemann, T. A. & Mitchell, S. F., 2009. Oldest Scleractinian fauna from Jamaica (Hauterivian, Benbow Inlier). *Journal of Paleontology*, 83: 333–349.
- Marshall, A. T., 1996. Calcification in hermatypic and ahermatypic corals. *Science*, 271: 637–639.
- Martin-Garin, B., Lathuilière, B. & Geister, J., 2012. The shifting biogeography of reef corals during the Oxfordian (Late Jurassic). A climatic control? *Palaeogeography, Palaeoclimatology, Palaeoecology*, 365: 136–153.
- Masse, J.-P., 1989. Relations entre modifications biologiques et phénomènes géologiques sur les plates-formes carbonatées du domaine périméditerranéen au passage Bédoulien Gargasien. *Géobios, Mémoire Spécial*, 11: 279–294.
- Masse, J.-P. & Fenerci-Masse, M., 2010. *Mathesia Mainelli* (Hippuritoidea, Monopleuridae) from the late Aptian-Albian of the Mediterranean region, a revision. *Turkish Journal of Earth Sciences*, 19: 543–556.
- Masse, J.-P. & Philip, J., 1981. Cretaceous coral-rudist buildups of France. In: Toomey, D. F. (ed.), *European Reef Models. Society of Economic Paleontology and Mineralogy Special Publication*, 30: 399–426.
- Mastandrea, A., Perri, E., Russo, F., Spadafora, A. & Tucker, M., 2006. Microbial primary dolomite from a Norian carbonate platform: northern Calabria, southern Italy. *Sedimentology*, 53: 465–480.
- Matyszkiewicz, J., Kochman, A. & Duś, A., 2012. Influence of local sedimentary conditions on development of microbialites in the Oxfordian carbonate buildups from the southern part of the Kraków–Częstochowa Upland (South Poland). *Sedimentary Geology*, 263–264: 109–132.
- Maurin, A. F., Bernet-Rollande, M. C., Monty, C. L. V. & Nazhat, S., 1985. The microbial nature of bacinellid textures. Sedimentological bearings. In: *9th European Regional Meeting of Sedimentology, Leiden, Abstracts*. pp. 285–287.
- Melnikova, G. K., 1975. *Late Triassic Scleractinians of the South-Western Pamirs*. Donish, Dushanbe, 234 pp. [In Russian].
- Melnikova, G. K. & Roniewicz, E., 1976. Contribution to the systematics and phylogeny of Amphistraeina (Scleractinia). *Acta Paleontologica Polonica*, 21: 97–114.
- Melnikova, G. K. & Roniewicz, E., 2012. Early Jurassic corals of the Pamir Mountains – a new Triassic–Jurassic transitional fauna. *Geologica Belgica*, 15: 376–38.
- Minkovska, V., 1996. Microfacies of the Emen Formation (Lo-vech Urgonian Group, Lower Cretaceous, Fore-Balkan). *Geologica Balcanica*, 26: 21–38.
- Minkovska, V., Peybernès, B., Nikolov, T. & Ivanov, M., 2002. Paleogeographic reconstruction of a segment of the North-Tethyan margin in Bulgaria from Barremian to Albian. *Eclogae Geologicae Helvetiae*, 95: 183–195.
- Montanaro Gallitelli, E., 1974. Microstructure and septal arrangement in a primitive Triassic coral. *Bollettino della Società Paleontologica Italiana*, 12: 8–22.
- Montanaro Gallitelli, E., 1975. Hexanthinaria, a new order of Zoantharia (Anthozoa, Coelenterata). *Bollettino della Società Paleontologica Italiana*, 14: 21–25.
- Morycowa, E., 1964a. Hexacoralla des couches de Grodziszczce (Neocomien Carpathes). *Acta Palaeontologica Polonica*, 9: 1–114.
- Morycowa, E., 1964b. Polypiers de la klippe de Kruhel Wielki près de Przemyśl (Tithonique, supérieur, Carpathes polonaises). *Rocznik Polskiego Towarzystwa Geologicznego*, 34: 489–508.
- Morycowa, E., 1968. Sur les calcaires exotique a Madréporaires dans les Lac Rożnów (Carpathes polonaises de Flysch). *Rocznik Polskiego Towarzystwa Geologicznego*, 38: 19–32.
- Morycowa, E., 1971. Hexacorallia et Octocorallia du Crétacé inférieur de Rarau (Carpathes orientales roumaines). *Acta Palaeontologica Polonica*, 16: 1–142.
- Morycowa, E., 1974. Hexacorallia d'un bloc exotique de calcaire a Woźniki pres Wadowice (Carpathes polonaises occidentales). *Acta Geologica Polonica*, 24: 457–484.
- Morycowa, E., 1988. Klippes of Kruhel Wielki. In: Kotlarczyk, J. (ed.), *Przewodnik LIX Zjazdu Polskiego Towarzystwa Geologicznego, Przemyśl, 16–18.09.1988*, Wydawnictwo AGH, Kraków, pp. 259–266. [In Polish].
- Morycowa, E., 2012. Corals from the Tithonian carbonate complex in the Dąbrowa Tarnowska–Szcuczin area (Polish Carpathian Foreland). *Annales Societatis Geologorum Poloniae*, 82: 1–38.
- Morycowa, E. & Decrouez, D., 2006. Early Aptian scleractinian corals from the Upper Schrattealk of Hergiswil (Lucerne region, Helvetic Zone of central Switzerland). *Revue de Paléobiologie*, 25: 791–838.
- Morycowa, E., Decrouez, D. & Schenk, K., 1994. Présence de *Latusastraea exiguis* (Scléactiniaire) dans le Schrattealk du Rawil (Helvétique, Suisse) et quelques sur les espèces créacées du genre *Latusastraea* d'Orbigny, 1849. *Annales Societatis Geologorum Poloniae*, 64: 15–22.
- Morycowa, E. & Lefeld, J., 1966. Les Madréporaires des calcaires urgoniens de la série haut-tatrique dans la Tatra Polonaise. *Rocznik Polskiego Towarzystwa Geologicznego*, 36: 519–542. [In Polish, French summary].
- Morycowa, E. & Marcopoulou-Diacantoni, A., 1997. Cretaceous scleractinian corals from the Parnassos area (central Greece) (Preliminary note). *Bulletin of the Geological Society of Greece*, 30 [for 1994]: 249–273.
- Morycowa, E. & Marcopoulou-Diacantoni, A., 2002. Albian corals from the Subpelagonian Zone of Central Greece (Agrostylia, Parnassos Region). *Annales Societatis Geologorum Poloniae*, 72: 1–65.
- Morycowa, E. & Masse, J.-P., 1998. Les scléactiniaires du Barremien–Aptien inférieur de Provence (SE de la France). *Géobios*, 31: 725–766.
- Muscatine, L., Goiran, C., Land, L., Jaubert, J., Cuif, J.-P. &

- Allemand, D., 2005. Stable isotopes ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) of organic matrix from coral skeleton. *Proceedings of the National Academy of Sciences of the United States of America*, 102: 1525–1530.
- Neuweiler, F., 1993. Development of Albian microbialites and microbialite reefs at marginal platform areas of the Vasco-Cantabrian Basin (Soba Reef Area, Cantabria, N. Spain). *Facies*, 29: 231–250.
- Neuweiler, F., 1995. Dynamische Sedimentationsvorgänge, Diagenese und Biofazies unterkretazischer Plattformränder (Apt/Alb; Soba Region, Prov. Cantabria, N-Spanien). *Berliner Geowissenschaftliche Abhandlungen*, E17: 1–235.
- Neuweiler, F., Gautret, P., Thiel, V., Longe, L., Michaelis, W. & Reitner, J., 1999. Petrology of Lower Cretaceous carbonate mud mounds (Albian, N. Spain): insights into organomineralic deposits of the geological record. *Sedimentology*, 46: 837–859.
- Neuweiler, F. and Reitner, J., 1992. Karbonatbänke mit *Lithocodium aggregatum* Elliott/*Bacinella irregularis* Radoicic. *Berliner Geowissenschaftliche Abhandlungen*, E3: 273–293.
- Nikolov, T., 1969. On the age of the urgonian sediments in the Fore-Balkan. *Bulletin Géologique Institut Bulgare de l'Académie des Sciences, Série Stratigraphie-Lithologie*, 18: 73–82.
- Nikolov, T., Ruskova, N., Ivanov, M. & Minkovska, V., 2007. Main features of the Lower Cretaceous in Bulgaria. *Comptes Rendus de l'Académie Bulgare des Sciences*, 60: 407–418.
- Nudds, J. R., 1979. Coloniality in Lithostrotionidae (Rugosa). In: Larwood, G. & Rosen, B. R. (eds), *Biology and Systematics of Colonial Organisms*. Academic Press, London, pp. 155–171.
- Ogilvie, M. M., 1897. Die Korallen der Stramberger Schichten. *Palaeontographica, A*, (Suppl. 2), 7: 73–284.
- Oliver, W. A., Jr., 1980a. On the relationship between Rugosa and Scleractinia (Summary). *Acta Palaeontologica Polonica*, 25: 395–402.
- Oliver, W. A., Jr., 1980b. The relationship of the scleractinian corals to the rugose corals. *Paleobiology*, 6: 146–160.
- Oliver, W. A., Jr., 1996. Origin and relationships of Paleozoic coral groups and the origin of the Scleractinia. *Paleontological Society Papers*, 1: 107–134.
- Olivier, N. & Boyet, M., 2006. Rare earth and trace elements of microbialites in Upper Jurassic coral- and sponge-microbialite reefs. *Chemical Geology*, 230: 105–123.
- Olivier, N., Carpentier, C., Martin-Garin, B., Lathuilière, B., Gaillard, C., Ferry, S., Hantzpergue, P., Gaillard, C. & Geister, J., 2004. Coral-microbialite reefs in pure carbonate versus mixed carbonate-siliciclastic environments: the example of the Pagny-sur-Meuse section (Upper Jurassic, northeastern France). *Facies*, 50: 229–255.
- Olivier, N., Colombié, C., Pittet, B. & Lathuilière, B., 2011. Microbial carbonates and corals on the marginal French Jura platform (Late Oxfordian, Molinges section). *Facies*, 57: 469–492.
- Olivier, N., Hantzpergue, P., Gaillard, C., Pittet, B., Leinfelder, R., Schmid, D. U. & Werner, W., 2003. Microbialite morphology, structure and growth: a model of the Upper Jurassic reefs of the Chay Peninsula (western France). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 193: 383–404.
- Peybernès, B., Ciszak, R., Nikolov, T., Ivanov, M. & Stoykova, K., 1998. Géométrie des formations et des séquences de dépôt éocétacées (intervalle Hauterivien–Aptien) de Bulgarie nord-orientale. *Comptes Rendus de l'Académie des Sciences, Paris, Sciences de la Terre et des Planètes*, 327: 555–562.
- Pleş, G., Mircescu, C. V., Bucur, I. I. & Săsăran, E., 2013. Encrusting micro-organisms and microbial structures in Upper Jurassic limestones from the Southern Carpathians (Romania). *Facies*, 59: 19–48.
- Poty, E., 1981. Recherches sur les Tétracoralliaires et les Hétérocoralliaires du Viséen de la Belgique. *Mededelingen Rijks Geologische Dienst, N.S.*, 35: 1–161.
- Rameil, N., Immenhauser, A., Warrlich, G., Hillgärtner, H. & Droste, H. J., 2010. Morphological patterns of Aptian *Lithocodium–Bacinella* geobodies: relation to environment and scale. *Sedimentology*, 57: 883–911.
- Reid, R. P., 1987. Nonskeletal peloidal precipitates in Upper Triassic reefs, Yukon Territory (Canada). *Journal of Sedimentary Petrology*, 57: 893–900.
- Reitner, J., 1993. Modern cryptic microbialite/metazoan-facies from Lizard Island (Great Barrier Reef, Australia) – formation and concepts. *Facies*, 29: 3–40.
- Reitner, J. & Neuweiler, F., 1995. Mud mounds: a polygenetic spectrum of fine-grained carbonate buildups. *Facies*, 32: 1–70.
- Reolid, M., 2007. Taphonomy of the Oxfordian-Lowermost Kimmeridgian siliceous sponges of the Prebetic Zone (Southern Iberia). *Journal of Taphonomy*, 5: 71–90.
- Ride, W. D. L., Cogger, H. G., Dupuis, C., Kraus, O., Minelli, A., Thompson, F. C. & Tubbs, P. K., eds, 1999. *International Code of Zoological Nomenclature*. International Commission on Zoological Nomenclature, London, 306 pp.
- Riding, R., 2000. Microbial carbonates: the geological record of calcified bacterial-algal mats and biofilms. *Sedimentology*, 47 (Supplement 1): 179–214.
- Riding, R., 2002. Structure and composition of organic reefs and carbonate mud mounds: concepts and categories. *Earth-Science Reviews*, 58: 163–231.
- Riding, R. & Tomás, S., 2006. Stromatolite reef crusts, Early Cretaceous, Spain: bacterial origin of in situ-precipitated peloid microspar? *Sedimentology*, 53: 23–34.
- Romano, S. L. & Palumbi, S. R., 1996. Evolution of scleractinian corals inferred from molecular systematics. *Science*, 271: 640–642.
- Roniewicz, E., 1966. Les Madréporaires du Jurassique supérieur de la bordure des Monts de Sainte-Croix, Pologne. *Acta Palaeontologica Polonica*, 11: 157–264.
- Roniewicz, E., 1976. Les Sclérectiniaires du Jurassique supérieur de la Dobrogea Centrale, Roumanie. *Palaeontologia Polonica*, 34: 17–121.
- Roniewicz, E., 1989. Triassic scleractinian corals of the Zlambach Beds, northern Calcareous Alps, Austria. *Österreichische Akademie der Wissenschaften, Wien*, 126: 1–152.
- Roniewicz, E., 2008. Kimmeridgian–Valanginian reef corals from the western Moesian Carbonate Platform from Bulgaria. *Annales Societatis Geologorum Poloniae*, 78: 91–134.
- Roniewicz, E., Mandl, G. W., Ebli, O. & Lobitzer, H., 2007. Early Norian scleractinian corals and microfacies data of the Dachstein Limestone of Feisterscharte, southern Dachstein Plateau (Northern Calcareous Alps, Austria). *Jahrbuch der Geologischen Bundesanstalt*, 147: 577–594.
- Roniewicz, E. & Morycowa, E., 1993. Evolution of the Scleractinia in the light of microstructural data. *Courier Forschungsinstitut Senckenberg*, 164: 233–240.
- Roniewicz, E. & Stolarski, J., 1999. Evolutionary trends in the epithecate scleractinian corals. *Acta Palaeontologica Polonica*, 44: 131–166.
- Roniewicz, E. & Stolarski, J., 2001. Triassic roots of amphistraeid scleractinian corals. *Journal of Paleontology*, 75: 34–45.
- Sabatier, P., Reyss, J. L., Hall-Spencer, J. M., Colin, C., Frank, N., Tisnérat-Laborde, N., Bordier, L. & Douville, E., 2012. ^{210}Pb – ^{226}Ra chronology reveals rapid growth rate of *Madrepora oculata* and *Lophelia pertusa* on world's largest cold-

- water coral reef. *Biogeosciences*, 9: 1253–1265.
- Sanders, D., 1999. Shell disintegration and taphonomic loss in rudist biostromes. *Lethaia*, 32: 101–112.
- Sanders, D. & Baron-Szabo, R. C., 2005. Scleractinian assemblages under sediment input: their characteristics and relation to the nutrient input concept. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 216: 139–181.
- Schäfer, P. & Senowbari-Daryan, B., 1980. *Aulastraea conica*, eine neue Koralle (Scleractinia: Amphiastreidae) aus Tithon-Riffkalcken der Madonie-Berge (Mittel-Sizilien) *Senckenhergiana Lethaea*, 61: 1–11.
- Schlager, W., 2003. Benthic carbonate factories of the Phanerozoic. *International Journal of Earth Sciences*, 92: 445–464.
- Schlagintweit, F., 2010. Taxonomic revision of *Lithocodium aggregatum* Elliott sensu Schmid & Leinfelder, 1996. *Jahrbuch der Geologischen Bundesanstalt*, 156: 393–406.
- Schlagintweit, F., 2012. New insights into *Troglotella incrustans* Wernli & Fookes, 1992, a fascinating Upper Jurassic–Upper Cretaceous foraminifer. *Studia Universitatis Babeş-Bolyai, Geologia*, 57: 17–26.
- Schlagintweit, F. & Bover-Arnal, T., 2012. The morphological adaption of *Lithocodium aggregatum* Elliott (calcareous green alga) to cryptic microhabitats (Lower Aptian, Spain): an example of phenotypic plasticity. *Facies*, 58: 37–55.
- Schlagintweit, F. & Bover-Arnal, T., 2013. Remarks on *Bacinnella Radoičić*, 1959 (type species *B. irregularis*) and its representatives. *Facies*, 59: 59–73.
- Schlagintweit, F., Bover-Arnal, T. & Salas, R., 2010. New insights into *Lithocodium aggregatum* Elliott 1956 and *Bacinnella irregularis* Radoičić 1959 (Late Jurassic–Lower Cretaceous): two ulvophyceean green algae (?Order Ulotrichales) with a heteromorphic life cycle (epilithic/euendolithic). *Facies*, 56: 509–547.
- Schlagintweit, F., Gawlick, H.-J. & Auer, M., 2007. *Carpathocancer* nov. nom. based on new findings of *Carpathocancer triangulatus* (Misik, Sotak & Ziegler, 1999) (type-species) nov. nom. in Triassic and Jurassic reefal carbonates of the Northern Calcareous Alps and their taxonomic re-interpretation as decapod crustacean appendages. *Journal of Alpine Geology (Mitteilungen der Gesellschaft der Geologie- und Bergbaustudenden in Österreich, Wien)*, 48: 23–35.
- Schlagintweit, F. & Velić, I., 2012. Foraminiferan tests and dasycladalean thalli as cryptic microhabitats for thaumatoporellacean algae from Mesozoic (Late Triassic–Late Cretaceous) platform carbonates. *Facies*, 58: 79–94.
- Schmid, D. U., 1996. Marine Mikrobolithe und Mikroinkrustierer aus dem Oberjura. *Profil*, 9: 101–251.
- Schmid, D. U. & Leinfelder, R. R., 1996. The Jurassic *Lithocodium aggregatum*–*Troglotella incrustans* foraminifer consortium. *Palaeontology*, 39: 21–52.
- Schmid, D. U., Leinfelder, R. R. & Nose, M., 2001. Growth dynamics and ecology of Upper Jurassic mounds, with comparisons to Mid-Palaeozoic mounds. *Sedimentary Geology*, 145: 343–376.
- Schöllhorn, E., 1998. Geologie und Paläontologie des Oberapt im Becken von Organyf (Nordspanien). *Coral Research Bulletin*, 6: 1–139.
- Scrutton, C. T., 1997. The Palaeozoic corals, I: origins and relationships. *Proceedings of the Yorkshire Geological Society*, 51: 177–208.
- Seard, C., Camoin, G. F., Yokoyama, Y., Matsuzaki, H., Durand, N., Bard, E., Sepulcre, S. & Deschamps, P., 2011. Microbialite development patterns in the last deglacial reefs from Tahiti (French Polynesia; IODP expedition #310): implications on reef framework architecture. *Marine Geology*, 279: 63–86.
- Shen, J. W. & Webb, G. E., 2005. Metazoan-microbial framework fabrics in a Mississippian (Carboniferous) coral-sponge-microbial reef, Monto, Queensland, Australia. *Sedimentary Geology*, 178: 113–133.
- Sikhuralidze, G. Y., 1979. Albskije korally siela Tschanar (Zapadnaja Gruzija). *Trudy Geologicheskogo Instituta AN Gruzinskoy SSR*, 63: 1–49. [In Russian].
- Silvestri, G., Bosellini, F. R. & Nebelsick, J. H., 2011. Microtaphofacies analysis of lower Oligocene turbid-water coral assemblages. *Palaaios*, 26: 805–820.
- Skelton, P. W. & Gili, E., 2012. Rudists and carbonate platforms in the Aptian: a case study on biotic interactions with ocean chemistry and climate. *Sedimentology*, 59: 81–117.
- Skelton, P. W., Gili, E., Rosen, B. R. & Valdeperas, F. X., 1997. Corals and rudists in the late Cretaceous: a critique of the hypothesis of competitive displacement. *Boletín de la Real Sociedad Española de Historia Natural, Sección Geológica*, 92: 225–239.
- Sprachta, S., Camoin, G., Golubic, S. & Le Campion, T., 2001. Microbialites in a modern lagoonal environment: nature and distribution (Tikehau atoll, French Polynesia). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 175: 103–124.
- Stanley, G. D., Jr., (ed.), 2001. *The History and Sedimentology of Reef Systems*. Kluwer Academic/Plenum Publishers, New York, 458 pp.
- Stanley, G. D., Jr., 2003. The evolution of modern corals and their early history. *Earth-Science Reviews*, 60: 195–225.
- Stanley, G. D., Jr. & Swart, P. K., 1995. Evolution of the coral-zooxanthellae symbiosis during the Triassic: A geochemical approach. *Paleobiology*, 21: 179–199.
- Stolarski, J., 1996. *Gardineria* – a scleractinian living fossil. *Acta Palaeontologica Polonica*, 41: 339–367.
- Stolarski, J., 2003. Three-dimensional micro- and nanostructural characteristics of the scleractinian coral skeleton: A biocalcification proxy. *Acta Palaeontologica Polonica*, 48: 497–530.
- Stolarski, J., Meibom, A., Przeniosło, R. & Mazur, M., 2007. A Cretaceous scleractinian coral with a calcitic skeleton. *Science*, 318: 92–94.
- Stolarski, J. & Roniewicz, E., 2001. Towards a new synthesis of evolutionary relationships and classification of Scleractinia. *Journal of Paleontology*, 75: 1090–1108.
- Stolarski, J., Roniewicz, E. & Grycuk, T., 2004. A model for furcate septal increase in a Triassic scleractiniomorph. *Acta Palaeontologica Polonica*, 49: 529–542.
- Stolarski, J. & Russo, A., 2001. Evolution of post-Triassic pachytheicaliine corals. *Bulletin of the Biological Society of Washington*, 10: 242–256.
- Stoykova, K. & Ivanov, M., 2000. Upper Barremian–Lower Aptian ammonite and nannofossil biostratigraphy in North Bulgaria. *Annual of the Sofia University. Geology-Geographical Faculty*, 92: 5–12.
- Szulc, J., 1997. Middle Triassic (Muschelkalk) sponge-microbial stromatolites, diplopores and *Girvanella*-oncoids from the Silesian-Cracow Upland. In: *3rd IFAA Regional Symposium & 3rd IGCP 380 International Meeting, Kraków, 1997, Guidebook & Abstracts*. pp. 10–15.
- Szulc, J., 2000. Middle Triassic evolution of the northern Peri-Tethys area as influenced by early opening of the Tethys Ocean. *Annales Societatis Geologorum Poloniae*, 70: 1–48.
- Tchechmedjieva, V., 2001. Scleractiniaires coloniaux du Barrémien près du village Pouchevo, l'arrondissement de Veliko Târnovo (Prébalkan Central). *Annuaire de l'Université de Sofia "St. Kliment Ohridski", Faculté de Géologie et Géographie*, 93 (for 2000): 5–20. [In Bulgarian, French abstract].

- Toula, F., 1889. Geologische Untersuchungen im centralen Balkan. *Denkschriften der Kaiserlichen Akademie Wissenschaften, Mathematisch-Physikalische Klasse*, 55: 1–108.
- Trichet, J. & Défarge, C., 1995. Non-biologically supported organomineralization. *Bulletin de l'Institut Océanographique (Monaco), Numéro Spécial*, 14: 203–236.
- Turnšek, D., 1972. Upper Jurassic Corals of Southern Slovenia. *Razprave, Razreda Slovenska Akademija Znanosti in Umetnosti, IV*, 15: 1–121.
- Turnšek, D., 1997. *Mesozoic Corals of Slovenia*. Znanstvenoraziskovalni center, Slovenska Akademija Znanosti in Umetnosti, 16, Ljubljana, 512 pp.
- Turnšek, D. & Buser, S., 1974. The Early Cretaceous Corals, Hydrozoans and Chaetetids of Banja Planota and Trnovski Gozd. *Razprave, Slovenska Akademija Znanosti in Umetnosti, IV*, 17: 283–124.
- Turnšek, D. & Löser, H., 1989. The history of Mesozoic coral research after 1940. *Courier Forschungsinstitut Senckenberg*, 164: 37–46.
- Turnšek, D. & Mihajlović, M., 1981. Lower Cretaceous Cnidarians from Eastern Serbia. *Razprave, Slovenska Akademija Znanosti in Umetnosti, Razred za Naravoslovne Vede, IV*, 23: 1–54.
- Turnšek, D., Buser, S. & Debeljak, I., 2003. Liassic coral patch reef above the “Lithiotid Limestone” on Trnovski Gozd Plateau, west Slovenia. *Razprave Slovenska Akademija Znanosti in Umetnosti, Razred za Naravoslovne Vede, IV*, 44: 285–331.
- Vašíček, Z. & Skupien, P., 2004. The Štramberk fossil site (Uppermost Jurassic/Lower Cretaceous, Outer Western Carpathians) – two centuries of the geological and paleontological research. *Sbornik Vědeckých Prací Vysoké Školy Báňské – Technické Univerzity Ostrava. Řada Hornicko-Geologická*, 50: 83–102. [In Czech, English summary].
- Vaughan, T. W. & Wells, J. W., 1943. *Revisions of the Suborders, Families and Genera of the Scleractinia*. *Geological Society of America, Special Paper*, 44, 363 pp.
- Webb, G. E., 1999. Youngest Early Carboniferous (late Viséan) shallow-water patch reefs in eastern Australia (Rockhampton Group, Queensland): combining quantitative micro- and macro-scale data. *Facies*, 41: 111–139.
- Webb, G. E., 2001. Biologically induced carbonate precipitation in reefs through time. In: Stanley, G. D. Jr. (ed.), *The History and Sedimentology of Ancient Reef Systems*. Kluwer Academic/Plenum Publishers, New York, pp. 159–203.
- Wells, J. W., 1956. Scleractinia. In: Moore, R. C. (ed.), *Treatise of Invertebrate Paleontology, part F*, Geological Society of America & University of Kansas Press, Lawrence, pp. 328–444.
- Wood, R., 1999. *Reef Evolution*. Oxford University Press, Oxford, 414 pp.
- Wright, D. T. & Wacey, D., 2005. Precipitation of dolomite using sulphate-reducing bacteria from the Coorong Region, South Australia: significance and implications. *Sedimentology*, 52: 987–1008.
- Wrzosek, T., 2007. A revision of the Devonian rugosan phillipsastroid genus *Smithicyathus*. *Acta Palaeontologica Polonica*, 52: 609–632.
- Young, G. A., 1999a. Fossil colonial corals: colony type and growth form. In: Savazzi, E. (ed.), *Functional Morphology of the Invertebrate Skeleton*. John Wiley & Sons, New York, pp. 647–666.
- Young, G. A., 1999b. Fossil colonial corals: growth patterns and coral-substrate relationships. In: Savazzi, E. (ed.), *Functional Morphology of the Invertebrate Skeleton*. John Wiley & Sons, New York, pp. 667–687.
- Zlatarski, V., 1967. *Dimorphocoeniopsis beauvaisorum*, une nouvelle espèce de Madreporaria du Crétacé inférieur de la Bulgarie du Nord. *Doklady Bolgarskoj Akademii Nauk*, 20: 1317–1320.
- Zlatarski, V., 1968a. Diplocteniopsidae, une nouvelle famille de Madreporaria de l'Aptien de la Bulgarie du Nord. *Bulletin of the Geological Institute, Series Paleontology*, 17: 49–106.
- Zlatarski, V., 1968b. *Paraclausastraea*, un nouveau genre de madreporaire de l'Aptien de la Bulgarie du Nord. *Review of the Bulgarian Geological Society*, 29: 159–171.
- Zlatarski, V., 1968c. *Paleobiologični izsledvanja varhu madrepori ot urgona v Srednia Predbalkan*. Disertatsia. Geologičeski Institut “Strashimir Dimitrov”, Balgarska Academia na Naukite, Sofia, 264 pp. [In Bulgarian].